

Collapsing steady states of the Keller-Segel System

Manuel del Pino¹, Juncheng Wei²

¹Departamento de Ingeniería Matemática and CMM,
Universidad de Chile, Casilla 170 Correo 3, Santiago, Chile.
E-mail address: delpino@dim.uchile.cl ²Department of Mathematics,
The Chinese University of Hong Kong, Shatin, Hong Kong.
E-mail address: wei@math.cuhk.edu.hk

Abstract

We consider the boundary value problem

$$\Delta u - au + \varepsilon^2 e^u = 0, \quad u > 0 \text{ in } \Omega, \quad \frac{\partial u}{\partial \nu} = 0 \text{ on } \partial\Omega,$$

which is equivalent to the stationary Keller-Segel system from chemotaxis. Here $\Omega \subset \mathbb{R}^2$ is a smooth and bounded domain. We show that given any two non-negative integers k, l with $k + l \geq 1$, for ε sufficiently small, there exists a solution u_ε for which with $\varepsilon^2 e^{u_\varepsilon}$ develops asymptotically k interior Dirac deltas with weight 8π and l boundary deltas with weight 4π . Location of blow-up points is characterized explicitly in terms of Green's function of the Neumann problem.

1 Introduction

Chemotaxis is one of the simplest mechanisms for aggregation of biological species. The term refers to a situation where organisms, for instance bacteria, move towards high concentrations of a chemical which they secrete. A basic model in chemotaxis was introduced by Keller and Segel [27]. They considered an advection-diffusion system consisting of two coupled parabolic equations for the concentration of the considered species and that of the chemical released, represented respectively by positive quantities $v(x, t)$ and $u(x, t)$ defined on a bounded, smooth domain Ω in \mathbb{R}^N under no-flux boundary conditions. The system reads as follows.

$$(1) \quad \begin{cases} v_t = \Delta v - \nabla(v \nabla u), \\ \tau u_t = \Delta u - u + v, \\ u, v > 0 \text{ in } \Omega, \quad \frac{\partial u}{\partial \nu} = \frac{\partial v}{\partial \nu} = 0 \text{ on } \partial\Omega. \end{cases}$$

Here τ is a positive constant. After the seminal works by Nanjudiah [32], and Childress and Percus [8], many contributions have been made to the understanding of different analytical aspects of this system and its variations. We refer the reader for instance to [3, 4, 7, 9, 15, 17], [22]-[26], [30]-[44]. It is well known that if space dimension is $N = 2$ then classical solutions may blow-up in finite time. Structure of this phenomenon has been widely treated in the literature for the two-dimensional case. It is known that blow-up for the quantity v (whose mass is clearly preserved in time) takes place as a finite sum of Dirac measures at points with masses greater than or equal to respectively 8π or 4π depending on whether they are located inside the domain or at the boundary. This phenomenon, commonly referred to as “chemotactic collapse”, has become fairly well understood. Asymptotic local profiles, forms of stability of blow-up and dynamics post blow-up have also been analyzed. Relatively less is known about steady states of the problem, namely solutions of the elliptic system

$$(2) \quad \begin{cases} \Delta v - \nabla(v \nabla u) = 0 & \text{in } \Omega, \\ \Delta u - u + v = 0 & \text{in } \Omega, \\ u, v > 0 \text{ in } \Omega, \quad \frac{\partial u}{\partial \nu} = \frac{\partial v}{\partial \nu} = 0 & \text{on } \partial\Omega. \end{cases}$$

Steady states are of basic importance for the understanding of the global dynamics of the system since, as pointed out in [33], a Lyapunov functional for (1) is present, see [21, 39]. This problem was first studied by Schaaf, [36] in the one-dimensional case. Biler [3] established existence of nontrivial solutions of (2) in higher dimensions in the radially symmetric case. In the general two-dimensional domain case, Wang and Wei ([44]), independently Senba and Suzuki ([37]), proved the following result: given any positive number λ with $\lambda \in (0, \frac{1}{|\Omega|} + \lambda_1) \setminus \{4\pi m\}_{m=1, \dots}$ (where λ_1 is the first positive eigenvalue of $-\Delta$ with Neumann boundary condition), there exists a non-constant solution to (2) with $\int_{\Omega} v = \lambda|\Omega|$.

The purpose of this paper is to construct non-trivial solutions to (2) with masses in the v coordinate close to $4\pi m$ for each given $m \geq 1$. More precisely, if $2k + l = m$, we are able to find solutions which exhibit in the limit l Dirac measures on the boundary and k inside the domain, with respective weights 4π and 8π . Our main result reads as follows.

Theorem 1.1 *Given non-negative integers k, l with $k + l \geq 1$, there exists a family of non-constant solutions $(u_{\varepsilon}, v_{\varepsilon})$, $\varepsilon > 0$ to problem (2) such that*

$$\lim_{\varepsilon \rightarrow 0} \int_{\Omega} v_{\varepsilon} = 4\pi(2k + l).$$

More precisely, up to subsequences, there exist k points $\xi_1, \dots, \xi_k \in \Omega$ and l points $\xi_{k+1}, \dots, \xi_{k+l} \in \partial\Omega$ such that as $\varepsilon \rightarrow 0$,

$$v_{\varepsilon} \rightarrow \sum_{i=1}^k 8\pi \delta_{\xi_i} + \sum_{i=1}^l 4\pi \delta_{\xi_{k+i}}.$$

This limiting phenomenon for steady states is a form of chemotactic collapse which would be interesting to relate with that present in finite time blow-up. Stability of these solutions opens as a basic question for a better understanding of global dynamics of (1). Hints toward a global phase portrait in a simplified radially symmetric model have been found in [4], in particular connections between blow-up and steady state solutions are conjectured.

As we will state in Theorem 1.2 below, much more accurate information on these solutions, in particular location of points ξ_i is explicitly described in terms of Green's function of the domain.

A basic feature of Problem (2) is that it can be reduced to a scalar equation as follows. It is easy to check that solutions of (2) satisfy the relation

$$\int_{\Omega} v |\nabla(\log v - u)|^2 = 0,$$

so that $v = \varepsilon^2 e^u$ for some positive constant ε . Thus system (2) is equivalent to the boundary value problem

$$(3) \quad \Delta u - u + \varepsilon^2 e^u = 0, u > 0 \text{ in } \Omega, \quad \frac{\partial u}{\partial \nu} = 0 \text{ on } \partial\Omega.$$

In what follows, we assume that $N = 2$ and look for non-trivial solutions of this problem when $\varepsilon > 0$ is a small number.

In [37, 40], Senba and Suzuki characterized the asymptotic behavior of families of solutions to (3) with uniformly bounded mass as $\varepsilon \rightarrow 0$. For $y \in \bar{\Omega}$ we denote by $G(x, y)$ Green's function of the problem

$$(4) \quad \Delta_x G - G + \delta_y = 0 \text{ in } \Omega, \quad \frac{\partial G}{\partial \nu_x} = 0 \text{ on } \partial\Omega.$$

The *regular part* of $G(x, y)$ is defined depending on whether y lies in the domain or on its boundary as

$$(5) \quad H(x, y) = \begin{cases} G(x, y) + \frac{1}{2\pi} \log|x - y|, & \text{if } y \in \Omega, \\ G(x, y) + \frac{1}{\pi} \log|x - y|, & \text{if } y \in \partial\Omega. \end{cases}$$

In this way, $H(\cdot, y)$ is of class $C^{1,\alpha}$ in $\bar{\Omega}$. In [37] and [40], the following fact was established: if u_ε is a family of solutions to problem (3) such that

$$\lim_{\varepsilon \rightarrow 0} \varepsilon \int_{\Omega} e^{u_\varepsilon} = \lambda_0 > 0$$

then there exist non-negative integers $k, l, m \geq 1$ for which $\lambda_0 = 4\pi(2k + l)$. Moreover, up to subsequences, there exist points ξ_i , $i = 1, \dots, m$ with $\xi_i \in \Omega$ for $i \leq k$ and $\xi_i \in \partial\Omega$ for $k < i \leq m$ for which

$$u_\varepsilon(x) \rightarrow \sum_{i=1}^k 8\pi G(x, \xi_j) + \sum_{i=k+1}^m 4\pi G(x, \xi_i),$$

uniformly on compact subsets of $\bar{\Omega} \setminus \{\xi_1, \dots, \xi_m\}$. Moreover, the m -tuple (ξ_1, \dots, ξ_m) is a critical point of the functional,

$$(6) \quad \varphi_m(x_1, \dots, x_m) = \sum_{i=1}^m c_i^2 H(x_i, x_i) + \sum_{i \neq j} c_i c_j G(x_i, x_j).$$

where $c_i = 8\pi$ for $i = 1, \dots, k$ where $c_i = 4\pi$ for $i = k + 1, \dots, m$, defined on $\Omega^k \times (\partial\Omega)^l$ where with no ambiguity we set $\varphi_m(x_1, \dots, x_m) = +\infty$ if $x_i = x_j$ for some $i \neq j$.

Our main result then establishes the reciprocal of this property: Let k, l be any non-negative integers such that $k + l \geq 1$. Then for any ε sufficiently small, problem (3) admits a solution u_ε satisfying the above properties. In order to make this statement more precise, let us denote

$$(7) \quad U_{\mu, a} = \log \frac{8\mu^2}{(\mu^2 + |x - a|^2)^2}, \quad \mu > 0, \quad a \in \mathbb{R}^2.$$

It is well-known that these functions correspond to all solutions to the problem

$$(8) \quad \Delta u + e^u = 0 \quad \text{in } \mathbb{R}^2, \quad \int_{\mathbb{R}^2} e^u < +\infty.$$

Our main result can be precised as follows:

Theorem 1.2 *Let k, l be non-negative integers with $k + l \geq 1$. Then for all sufficiently small ε there is a solution u_ε to problem (4) with the following properties:*

(1) u_ε has exactly $k+l$ local maximum points $\xi_i^\varepsilon, i = 1, \dots, k+l$ such that $\xi_i^\varepsilon \in \Omega$, for $i \leq k$ and $\xi_i^\varepsilon \in \partial\Omega$ for $k + 1 \leq i \leq k + l$. Furthermore

$$\lim_{\varepsilon \rightarrow 0} \varphi_m(\xi_1^\varepsilon, \dots, \xi_m^\varepsilon) = \min_{\Omega^k \times (\partial\Omega)^l} \varphi_m.$$

(2) There are constants $\mu_j > 0$ such that

$$u_\varepsilon(x) = \sum_{j=1}^m U_{\varepsilon\mu_j, \xi_j^\varepsilon}(x) + O(1).$$

(3) $\varepsilon^2 \int_\Omega e^{u_\varepsilon} \rightarrow 4\pi(2k + l)$ as $\varepsilon \rightarrow 0$.

The existence of a global minimum for the function φ_m in $\Omega^k \times (\partial\Omega)^l$ follows from properties of the Green's function, see the proof of Lemma 7.1. In reality, associated to each *topologically nontrivial* for φ_m , a bubbling solution at a corresponding critical point exists, see §8.

It is important to remark the analogy existing between our results and those known for the Liouville-type equation

$$(9) \quad \begin{cases} \Delta u + \varepsilon^2 e^u = 0 & \text{in } \Omega \\ u = 0 & \text{on } \partial\Omega. \end{cases}$$

Asymptotic behavior of solutions of (9) for which $\varepsilon^2 \int_{\Omega} e^u$ remains uniformly bounded is well understood after the works [5, 28, 29]: $\varepsilon^2 e^u$ approaches a superposition of Dirac deltas in the interior of Ω . Construction of solutions with this behavior has been achieved in [1, 11, 16]. Related constructions for problems involving nonlinear, exponential boundary conditions for which boundary concentration have been performed in [12] and [13]. A special feature of the problem treated in this paper is the presence of **mixed** boundary-interior bubbling solutions. A similar phenomenon had only been observed in [19], for a different Neumann singularly perturbed problem. To capture such solutions, we use the so-called “localized energy method”- a combination of Lyapunov-Schmidt reduction method and variational techniques. Namely, we first use Lyapunov-Schmidt reduction method to convert the problem into a finite dimensional one, for a suitable asymptotic *reduced energy*, related with φ_m . Such a scheme has been used in many works, see for instance [2, 14, 10, 18, 19, 20, 11, 16, 35] and references therein. Our approach shares elements with that in [11], however, a different, more delicate functional setting has to be introduced. In what remains of this paper we shall prove Theorem 1.2.

2 Ansatz for the solution

Given $\xi_j \in \bar{\Omega}$, $\mu_j > 0$ we define

$$u_j(x) = \log \frac{8\mu_j^2}{(\varepsilon^2\mu_j^2 + |x - \xi_j|^2)^2}.$$

The choice of ξ_j and μ_j will be made later on.

The ansatz is

$$(10) \quad U(x) = \sum_{j=1}^m (u_j(x) + H_j^\varepsilon(x))$$

where H_j^ε is a correction term defined as the solution of

$$(11) \quad \begin{cases} -\Delta H_j^\varepsilon + H_j^\varepsilon = -u_j & \text{in } \Omega \\ \frac{\partial H_j^\varepsilon}{\partial \nu} = -\frac{\partial u_j}{\partial \nu} & \text{on } \partial\Omega. \end{cases}$$

Lemma 2.1 *For any $0 < \alpha < 1$*

$$(12) \quad H_j^\varepsilon(x) = c_j H(x, \xi_j) - \log 8\mu_j^2 + O(\varepsilon^\alpha)$$

uniformly in $\bar{\Omega}$, where H is the regular part of Green's function defined (5).

We will give the proof of this lemma at the end of the section.

It will be convenient to work with the scaling of u given by

$$v(y) = u(\varepsilon y) + 4 \log \varepsilon.$$

If u is a solution of (3) then v satisfies

$$(13) \quad \begin{cases} -\Delta v + \varepsilon^2(v - 4 \log \varepsilon) = e^v & \text{in } \Omega_\varepsilon \\ \frac{\partial v}{\partial \nu} = 0 & \text{on } \partial\Omega_\varepsilon, \end{cases}$$

where $\Omega_\varepsilon = \Omega/\varepsilon$. With this scaling u_j becomes

$$v_j(y) = \log \frac{8\mu_j}{(\mu_j^2 + |y - \xi_j'|^2)^2}$$

where $\xi_j' = \xi_j/\varepsilon$ and where we will write ν for the exterior normal unit vector to $\partial\Omega$ and $\partial\Omega_\varepsilon$.

Note that $u_j + H_j^\varepsilon$ satisfies

$$(14) \quad \begin{cases} -\Delta(u_j + H_j) + \varepsilon^2(u_j + H_j) = e^{v_j} & \text{in } \Omega_\varepsilon \\ \frac{\partial(u_j + H_j)}{\partial \nu} = 0 & \text{on } \partial\Omega_\varepsilon. \end{cases}$$

We will seek a solution v of (13) of the form

$$v = V + \phi,$$

where

$$(15) \quad V(y) = U(\varepsilon y) + 4 \log \varepsilon$$

and U is defined by (10). Problem (13) can be stated as to find ϕ a solution to

$$(16) \quad \begin{cases} -\Delta\phi + \varepsilon^2\phi = e^V\phi + N(\phi) + R & \text{in } \Omega_\varepsilon \\ \frac{\partial\phi}{\partial \nu} = 0 & \text{on } \partial\Omega_\varepsilon, \end{cases}$$

where the “nonlinear term” is

$$(17) \quad N(\phi) = e^V(e^\phi - 1 - \phi)$$

and the “error term” is given by

$$(18) \quad R = \Delta V - \varepsilon^2(V - 4 \log \varepsilon) + e^V.$$

At this point it is convenient to make a choice of the parameters μ_j , the objective being to make the error term small. We claim that if

$$(19) \quad \log 8\mu_j^2 = c_j H(\xi_j, \xi_j) + \sum_{i \neq j} c_i G(\xi_i, \xi_j),$$

then we achieve the following behavior for R : for any $0 < \alpha < 1$ there exists C independent of ε such that

$$(20) \quad |R(y)| \leq C\varepsilon^\alpha \sum_{j=1}^m \frac{1}{1 + |y - \xi_j'|^3} \quad \forall y \in \Omega_\varepsilon$$

and for $W = e^V$

$$(21) \quad W(y) = \sum_{j=1}^m \frac{8\mu_j^2}{(\mu_j^2 + |y - \xi_j|^2)^2} (1 + \theta_\varepsilon(y)), \quad \forall y \in \Omega_\varepsilon$$

with θ_ε satisfying the following estimate

$$|\theta_\varepsilon(y)| \leq C\varepsilon^\alpha + C\varepsilon \sum_{j=1}^m |y - \xi'_j| \quad \forall y \in \Omega_\varepsilon.$$

Proof of (21).

$$\begin{aligned} W(y) &= \varepsilon^4 \exp \left(\sum_{i=1}^m u_j(\varepsilon y) + H_i^\varepsilon(\varepsilon y) \right) \\ &= \exp \left(\sum_{i=1}^m \left(\log \frac{8\mu_i^2}{(\mu_i^2 + |y - \xi'_i|^2)^2} + H_i^\varepsilon(\varepsilon y) \right) \right). \end{aligned}$$

Let us fix a small constant $\delta > 0$ and consider this expression for $|y - \xi'_j| < \frac{\delta}{\varepsilon}$

$$\begin{aligned} W(y) &= \frac{8\mu_j^2}{(\mu_j^2 + |y - \xi'_j|^2)^2} \exp \left(H_j^\varepsilon(\varepsilon y) + \sum_{i \neq j}^m \left[\log \frac{8\mu_i^2}{(\varepsilon^2 \mu_i^2 + \varepsilon^2 |y - \xi'_i|^2)^2} \right. \right. \\ &\quad \left. \left. + H_i^\varepsilon(\varepsilon y) \right] \right). \end{aligned}$$

Using (12) and the fact that H is $C^1(\partial\Omega^2)$ we have

$$\begin{aligned} H_i^\varepsilon(\varepsilon y) &= c_i H(\varepsilon y, \xi_i) - \log(8\mu_i^2) + O(\varepsilon^\alpha) \quad \forall y \in \Omega_\varepsilon \\ &= c_i H(\xi_j, \xi_i) - \log(8\mu_i^2) + O(\varepsilon^\alpha) + O(\varepsilon |y - \xi'_j|) \quad \forall y \in \Omega_\varepsilon. \end{aligned}$$

Hence for $|y - \xi'_j| < \frac{\delta}{\varepsilon}$

$$\begin{aligned} H_j^\varepsilon(\varepsilon y) &+ \sum_{i \neq j}^m \left(\log \frac{8\mu_i^2}{(\varepsilon^2 \mu_i^2 + \varepsilon^2 |y - \xi'_i|^2)^2} + H_i^\varepsilon(\varepsilon y) \right) \\ &= c_j H(\xi_j, \xi_j) - \log(8\mu_j^2) + \sum_{i \neq j}^m \left(\log \frac{8\mu_i^2}{|\xi_j - \xi_i|^4} + c_i H(\xi_j, \xi_i) - \log(8\mu_i^2) \right) \\ &\quad + O(\varepsilon^\alpha) + O(\varepsilon |y - \xi'_j|). \\ &= c_j H(\xi_j, \xi_j) - \log(8\mu_j^2) + \sum_{i \neq j}^m c_i G(\xi_j, \xi_i) + O(\varepsilon^\alpha) + O(\varepsilon |y - \xi'_j|) \\ &= O(\varepsilon^\alpha) + O(\varepsilon |y - \xi'_j|) \end{aligned}$$

by the choice of μ_j , c.f. (19). Therefore

$$(22) \quad W(y) = \frac{8\mu_j^2}{(\mu_j^2 + |y - \xi'_j|^2)^2} (1 + O(\varepsilon^\alpha) + O(\varepsilon |y - \xi'_j|)) \quad \forall |y - \xi'_j| < \frac{\delta}{\varepsilon}.$$

If $|y - \xi'_j| > \frac{\delta}{\varepsilon}$ for all $j = 1, \dots, m$ we have $W = O(\varepsilon^4)$, and this together with (22) implies (21). \square

Proof of (20). We defined $R = \Delta V - \varepsilon^2(V - 4\log \varepsilon) + e^V$ with V given by (15).

By our definition and (14),

$$R = \varepsilon^4 e^{\sum_{j=1}^m (u_j + H_j)} - \sum_{j=1}^m e^{v_j}$$

For $|y - \xi'_j| < \frac{\delta}{\varepsilon}$, we have according to (22)

$$\begin{aligned} R &= e^{v_j} (1 + O(\varepsilon^\alpha) + O(\varepsilon|y - \xi'_j|)) - e^{v_j} + O(\varepsilon^4) \\ &= O(e^{v_j}(\varepsilon^\alpha + \varepsilon|y - \xi'_j|)) + O(\varepsilon^4) \end{aligned}$$

which proves (20). \square

Proof of lemma 2.1. The boundary condition satisfied by H_j^ε is

$$(23) \quad \frac{\partial H_j^\varepsilon}{\partial \nu} = -\frac{\partial u_j}{\partial \nu} = 4 \frac{(x - \xi_j) \cdot \nu(x)}{\varepsilon^2 \mu_j^2 + |x - \xi_j|^2}$$

Thus for $\xi_j \in \Omega, j = 1, \dots, k$, we have

$$(24) \quad \frac{\partial H_j^\varepsilon}{\partial \nu} = 4 \frac{(x - \xi_j) \cdot \nu(x)}{|x - \xi_j|^2} + O(\varepsilon^2) \quad \text{on } \partial\Omega.$$

For $\xi_j \in \partial\Omega, j = k+1, \dots, k+l$, we have

$$(25) \quad \lim_{\varepsilon \rightarrow 0} \frac{\partial H_j^\varepsilon}{\partial \nu}(x) = 4 \frac{(x - \xi_j) \cdot \nu(x)}{|x - \xi_j|^2} \quad \forall x \neq \xi_j.$$

The regular part of Green's function $H(x, \xi_j)$ satisfies

$$\begin{cases} -\Delta_x H(x, \xi_j) + H(x, \xi_j) = -\frac{4}{c_j} \log \frac{1}{|x - \xi_j|} & x \in \Omega \\ \frac{\partial H}{\partial \nu_x}(x, \xi_j) = \frac{4}{c_j} \frac{(x - \xi_j) \cdot \nu(x)}{|x - \xi_j|^2} & x \in \partial\Omega \end{cases}$$

For the difference $z_\varepsilon(x) = H_j^\varepsilon(x) + \log 8\mu_j^2 - c_j H(x, \xi_j)$ we have

$$\begin{cases} -\Delta z_\varepsilon + z_\varepsilon = -\log \frac{1}{(\varepsilon^2 \mu_j^2 + |x - \xi_j|^2)^2} + \log \frac{1}{|x - \xi_j|^4} & \text{in } \Omega \\ \frac{\partial z_\varepsilon}{\partial \nu} = \frac{\partial H_j^\varepsilon}{\partial \nu} - 4 \frac{(x - y) \cdot \nu(x)}{|x - y|^2} & \text{on } \partial\Omega. \end{cases}$$

We claim that for any $p > 1$ there exists $C > 0$ such that

$$(26) \quad \left\| \frac{\partial H_j^\varepsilon}{\partial \nu} - 4 \frac{(x - \xi_j) \cdot \nu(x)}{|x - \xi_j|^2} \right\|_{L^p(\partial\Omega)} \leq C\varepsilon^{1/p}.$$

For this it will be convenient to observe first that for $\xi_j \in \partial\Omega$

$$(27) \quad |(x - \xi_j) \cdot \nu(x)| \leq C|x - \xi_j|^2 \quad \forall x \in \partial\Omega,$$

which can be proved, for example, assuming that $\xi_j = 0$ and that near the origin $\partial\Omega$ is the graph of a function $G : (-a, a) \rightarrow \mathbb{R}$ with $G(0) = G'(0) = 0$. Now

$$(28) \quad \frac{\partial H_j^\varepsilon}{\partial \nu} - 4 \frac{(x - \xi_j) \cdot \nu(x)}{|x - \xi_j|^2} = \varepsilon^2 \mu_j^2 \frac{(x - \xi_j) \cdot \nu(x)}{|x - \xi_j|^2 (\varepsilon^2 \mu_j^2 + |x - \xi_j|^2)}.$$

By (27)

$$(29) \quad \left| \frac{\partial H_j^\varepsilon}{\partial \nu} - 4 \frac{(x - \xi_j) \cdot \nu(x)}{|x - \xi_j|^2} \right| \leq C \frac{\varepsilon^2}{\varepsilon^2 \mu_j^2 + |x - \xi_j|^2}.$$

Fix $\rho > 0$ small. Then

$$(30) \quad \left| \frac{\partial H_j^\varepsilon}{\partial \nu} - 4 \frac{(x - \xi_j) \cdot \nu(x)}{|x - \xi_j|^2} \right| \leq C\varepsilon^2 \quad \forall |x - \xi_j| \geq \rho, \quad x \in \partial\Omega.$$

Now let $p > 1$. Changing variables $x - \xi_j = \varepsilon y$ we have

$$\begin{aligned} \int_{B_\rho(\xi_j) \cap \partial\Omega} \left| \frac{\varepsilon^2}{(\varepsilon^2 \mu_j^2 + |x - \xi_j|^2)} \right|^p dx &= C\varepsilon \int_{B_{\rho/\varepsilon}(0) \cap \Omega_\varepsilon} \left| \frac{1}{\mu_j^2 + |y|^2} \right|^p dy \\ &\leq C\varepsilon \int_0^{\rho/\varepsilon} \frac{1}{(1+s^2)^p} ds \\ &\leq C\varepsilon. \end{aligned}$$

Combining this with (29) and (30) we conclude that (26) holds.

For $p > 1$ let us estimate now

$$\begin{aligned} \left\| \log \frac{1}{|x - \xi_j|^2} - \log \frac{1}{\varepsilon^2 \mu_j^2 + |x - \xi_j|^2} \right\|_{L^p(\Omega)}^p &= \int_{B_{10\varepsilon\mu_j}(\xi_j) \cap \Omega} \dots \\ &\quad + \int_{\Omega \setminus B_{10\varepsilon\mu_j}(\xi_j)} \dots \\ &= I_1 + I_2. \end{aligned}$$

For I_1 observe that

$$\int_{B_{10\varepsilon\mu_j}(\xi_j) \cap \Omega} \left| \log \frac{1}{|x - \xi_j|^2} \right|^p dx \leq C \int_0^{C\varepsilon} |\log r|^p r dr \leq C\varepsilon^2 (\log \frac{1}{\varepsilon})^p.$$

The same bound is true for the integral of $|\log \frac{1}{\varepsilon^2 \mu_j^2 + |x - \xi_j|^2}|^p$ in $B_{10\varepsilon\mu_j}(\xi_j) \cap \Omega$. Hence

$$|I_1| \leq C\varepsilon^2 \left(\log \frac{1}{\varepsilon}\right)^p.$$

Let us estimate I_2 as follows:

$$\left| \log \frac{1}{|x - \xi_j|^2} - \log \frac{1}{\varepsilon^2 \mu_j^2 + |x - \xi_j|^2} \right| \leq \frac{C\varepsilon}{|x - \xi_j|}.$$

Take $1 < p < 2$ and integrate

$$|I_2| \leq C\varepsilon^p \int_{10\mu\varepsilon}^D r^{1-p} dr \leq C\varepsilon^p,$$

where D is the diameter of Ω . In conclusion, for any $1 < p < 2$ we have

$$\left\| \log \frac{1}{|x - \xi_j|^2} - \log \frac{1}{\varepsilon^2 \mu_j^2 + |x - \xi_j|^2} \right\|_{L^p(\Omega)} \leq C\varepsilon.$$

By L^p theory

$$\|z_\varepsilon\|_{W^{1+s,p}(\Omega)} \leq C \left(\left\| \frac{\partial z_\varepsilon}{\partial \nu} \right\|_{L^p(\partial\Omega)} + \|\Delta z_\varepsilon\|_{L^p(\Omega)} \right) \leq C\varepsilon^{1/p}$$

for any $0 < s < \frac{1}{p}$. By the Morrey embedding we obtain

$$\|z_\varepsilon\|_{C^\gamma(\bar{\Omega})} \leq C\varepsilon^{1/p}$$

for any $0 < \gamma < \frac{1}{2} + \frac{1}{p}$. This proves the result (with $\alpha = \frac{1}{p}$). \square

Remark. The convergence (25) is not uniform in general because $\frac{\partial H_\varepsilon}{\partial \nu}(\xi_j) = 0$ while the function $x \mapsto 2\frac{(x-\xi_j) \cdot \nu(x)}{|x-\xi_j|^2}$ can be extended continuously to ξ_j with a value equal to the curvature of $\partial\Omega$ at ξ_j .

3 Solvability of a linear equation

The main result of this section is the solvability of the following linear problem: given h find $\phi, c_{11}, \dots, c_{mJ_m}$ such that

$$(31) \quad \begin{cases} -\Delta\phi + \varepsilon^2\phi = W\phi + h + \sum_{j=1}^m \sum_{i=1, J_j} c_{ij} \chi_j Z_{ij} & \text{in } \Omega_\varepsilon \\ \frac{\partial\phi}{\partial\nu} = 0 & \text{on } \partial\Omega_\varepsilon \\ \int_{\Omega_\varepsilon} \chi_j Z_{ij} \phi = 0 & \forall j = 1, \dots, m, i = 1, J_j \end{cases}$$

where $m = k + l$, W is a function that satisfies (21), $h \in L^\infty(\Omega_\varepsilon)$ and Z_{ij} , χ_j are defined as follows, $J_j = 2$ if $j = 1, \dots, k$ and $J_j = 1$ if $j = k + 1, \dots, k + l$.

Let z_{ij} be

$$z_{0j} = \frac{1}{\mu_j} - 2\frac{\mu_j}{\mu_j^2 + y^2}, \quad z_{ij} = \frac{y_i}{\mu_j^2 + y^2}.$$

It is well-known that any solution to

$$(32) \quad \Delta\phi + e^{v_j}\phi = 0, |\phi| \leq C(1 + |y|)^\sigma$$

is a linear combination of z_{ij} , $i = 0, 1, 2$. (See Lemma 2.1 of [6].)

Next we choose a large but fixed number R_0 and nonnegative smooth function $\chi : \mathbb{R} \rightarrow \mathbb{R}$ so that $\chi(r) = 1$ for $r \leq R_0$ and $\chi(r) = 0$ for $r \geq R_0 + 1$, $0 \leq \chi \leq 1$.

For $j = 1, \dots, k$ (corresponding to **interior** bubble case), we define

$$(33) \quad \chi_j(y) = \chi(|y - \xi'_j|), \quad Z_{ij}(y) = z_{ij}(y), \quad i = 0, 1, 2, \quad j = 1, \dots, k.$$

For $j = k + 1, \dots, k + l$ (corresponding to **boundary** bubble case), we have to strengthen the boundary first. More precisely, at the boundary point $\xi_j \in \partial\Omega$, we assume that $\xi_j = 0$ and the unit outward normal at ξ_j is $-\mathbf{e}_2 = (0, -1)$. Let $G(x_1)$ be the defining function for the boundary $\partial\Omega$ in a neighborhood $B_\rho(\xi_j)$ of ξ_j , that is, $\Omega \cap B_\rho(\xi_j) = \{(x_1, x_2) | x_2 > G(x_1), (x_1, x_2) \in B_\rho(\xi_j)\}$. Then, let $F_j : B_\rho(\xi_j) \cap N \rightarrow \mathbb{R}^2$ be defined by

$$(34) \quad F_j = (F_{j,1}, F_{j,2}), \quad \text{where} \quad F_{j,1} = x_1 + \frac{x_2 - G(x_1)}{1 + |G'(x_1)|^2} G'(x_1), \quad F_{j,2} = x_2 - G(x_1)$$

Then we set

$$(35) \quad F_j^\varepsilon(y) = \frac{1}{\varepsilon} F_j(\varepsilon y).$$

Note that F_j preserves the Neumann boundary condition. Define

$$(36) \quad \chi_j(y) = \chi(|F_j^\varepsilon(y)|), \quad Z_{ij}(y) = z_{ij}(F_j^\varepsilon(y)) \quad i = 0, 1 \quad j = k + 1, \dots, k + l.$$

It is important to note that

$$(37) \quad \Delta Z_{0j} + e^{v_j} Z_{0j} = O(\varepsilon(1 + |y - \xi'_j|)^{-3})$$

since

$$\nabla z_{0j} = O\left(\frac{1}{(1 + |y - \xi'_j|)^3}\right).$$

All functions above depend on ε but we omit this dependence in the notation. Furthermore, Z_{ij} now all satisfies the Neumann boundary condition (since F_j preserves the Neumann boundary condition).

Equation (31) will be solved for $h \in L^\infty(\Omega_\varepsilon)$ but we will be able to estimate the size of the solution in terms of the following norm

$$(38) \quad \|h\|_\infty = \sup_{y \in \Omega_\varepsilon} |h(y)|, \quad \|h\|_* = \sup_{y \in \Omega_\varepsilon} \frac{|h(y)|}{\varepsilon^2 + \sum_{j=1}^m (1 + |y - \xi'_j|)^{-2-\sigma}},$$

where we fix $0 < \sigma < 1$ although the precise choice will be made later on.

Proposition 3.1 *Let $d > 0$ and m a positive integer. Then there exist $\varepsilon_0 > 0$, C such that for any $0 < \varepsilon < \varepsilon_0$, any family of points $(\xi_1, \dots, \xi_m) \in \mathcal{M}_\delta$ and any $h \in L^\infty(\Omega_\varepsilon)$ there is a unique solution $\phi \in L^\infty(\Omega_\varepsilon)$, $c_{ij} \in \mathbb{R}$ to (31). Moreover*

$$\|\phi\|_{L^\infty(\Omega_\varepsilon)} \leq C \log \frac{1}{\varepsilon} \|h\|_*.$$

We begin by stating an a-priori estimate for solutions of (31) satisfying orthogonality conditions with respect to $Z_{ij}, i = 0, J_j, j = 1, \dots, m$.

Lemma 3.2 *There are $R_0 > 0$ and $\varepsilon_0 > 0$ so that for $0 < \varepsilon < \varepsilon_0$ and any solution ϕ of (31) with the orthogonality conditions*

$$(39) \quad \int_{\Omega_\varepsilon} Z_{ij} \chi_j \phi = 0 \quad \forall i = 0, \dots, J_j \quad \forall j = 1, \dots, m$$

we have

$$\|\phi\|_{L^\infty(\Omega_\varepsilon)} \leq C \|h\|_*$$

where C is independent of ε .

For the proof of this lemma we need to construct a suitable barrier.

Lemma 3.3 *For $\varepsilon > 0$ small enough there exist $R_1 > 0$, and*

$$\psi : \Omega_\varepsilon \setminus \cup_{j=1}^m B_{R_1}(\xi'_j) \rightarrow \mathbb{R}$$

smooth and positive so that

$$\begin{aligned} -\Delta \psi + \varepsilon^2 \psi - W \psi &\geq \sum_{j=1}^m \frac{1}{|y - \xi'_j|^{2+\sigma}} + \varepsilon^2 && \text{in } \Omega_\varepsilon \setminus \bigcup_{j=1}^m B_{R_1}(\xi'_j) \\ \frac{\partial \psi}{\partial \nu} &\geq 0 && \text{on } \partial \Omega_\varepsilon \setminus \bigcup_{j=1}^m B_{R_1}(\xi'_j) \\ \psi &> 0 && \text{in } \Omega_\varepsilon \setminus \bigcup_{j=1}^m B_{R_1}(\xi'_j) \\ \psi &\geq 1 && \text{on } \Omega_\varepsilon \cap \left(\bigcup_{j=1}^m \partial B_{R_1}(\xi'_j) \right) \end{aligned}$$

The constants $R_1 > 0$, $c > 0$ can be chosen independently of ε and ψ is bounded uniformly

$$0 < \psi \leq C \quad \text{in } \Omega_\varepsilon \setminus \cup_{j=1}^m B_{R_1}(\xi'_j).$$

Proof of lemma 3.2. We take $R_0 = 2R_1$, R_1 being the constant of lemma 3.3. Thanks to the barrier ψ of that lemma we deduce that the following maximum principle holds in $\Omega_\varepsilon \setminus \cup_{j=1}^m B_{R_1}(\xi'_j)$: if $\phi \in H^1(\Omega_\varepsilon \setminus \cup_{j=1}^m B_{R_1}(\xi'_j))$ satisfies

$$\left\{ \begin{array}{l} -\Delta\phi + \varepsilon^2\phi \geq W\phi \quad \text{in } \Omega_\varepsilon \setminus \bigcup_{j=1}^m B_{R_1}(\xi'_j) \\ \frac{\partial\phi}{\partial\nu} = 0 \quad \text{on } \partial\Omega_\varepsilon \setminus \bigcup_{j=1}^m B_{R_1}(\xi'_j) \\ \phi \geq 0 \quad \text{on } \Omega_\varepsilon \cap \left(\bigcup_{j=1}^m \partial B_{R_1}(\xi'_j) \right) \end{array} \right.$$

then $\phi \geq 0$ in $\Omega_\varepsilon \setminus \cup_{j=1}^m B_{R_1}(\xi'_j)$.

Let h be bounded and ϕ a solution to (31) satisfying (39). Following [11] we first claim that $\|\phi\|_{L^\infty(\Omega_\varepsilon)}$ can be controlled in terms of $\|h\|_*$ and the following inner norm of ϕ :

$$\|\phi\|_i = \sup_{\Omega_\varepsilon \cap (\cup_{j=1}^m B_{R_1}(\xi'_j))} |\phi|.$$

Indeed, set

$$\tilde{\phi} = C_1\psi(\|\phi\|_i + \|f\|_*),$$

with C_1 a constant independent of ε . By the above maximum principle we have $\phi \leq \tilde{\phi}$ and $-\phi \leq \tilde{\phi}$ in $\Omega_\varepsilon \setminus \cup_{j=1}^m B_{R_1}(\xi'_j)$. Since ψ is uniformly bounded we deduce

$$(40) \quad \|\phi\|_{L^\infty(\Omega_\varepsilon)} \leq C(\|\phi\|_i + \|f\|_*),$$

for some constant C independent of ϕ and ε .

We prove the lemma by contradiction. Assume that there exist a sequence $\varepsilon_n \rightarrow 0$, points $(\xi_1^n, \dots, \xi_m^n) \in \mathcal{M}_\delta$ and functions ϕ_n, f_n and h_n with $\|\phi_n\|_{L^\infty(\Omega_{\varepsilon_n})} = 1$ and $\|h_n\|_* \rightarrow 0$ so that for each n ϕ_n solves (31) and satisfies (39). By (40) we see that $\|\phi_n\|_i$ stays away from zero. For one of the indices, say j , we can assume that $\sup_{B_{R_1}(\xi'_j)} |\phi_n| \geq c > 0$ for all n . Consider $\hat{\phi}_n(z) = \phi_n(z - \xi'_j)$ and let us translate and rotate Ω_{ε_n} so that Ω_{ε_n} approaches the upper half plane \mathbb{R}_+^2 and $\xi'_j = 0$. Then by elliptic estimates $\hat{\phi}_n$ converges uniformly on compact sets to a nontrivial solution of

$$\Delta\phi + e^{v_j}\phi = 0, |\phi| \leq C.$$

Thus $\hat{\phi}$ is a linear combination of $z_{ij}, i = 0, \dots, J_j$. On the other hand we can take the limit in the orthogonality relations (39), observing that limits of the functions Z_{ij} are just rotations and translations of z_{ij} , and we find $\int_{\mathbb{R}_+^2} \chi_{\hat{\phi}} z_{ij} = 0$ for $i = 0, J_j$. This contradicts the fact that $\hat{\phi} \not\equiv 0$. \square

Proof of lemma 3.3.

We take

$$(41) \quad \psi_{1j}(r) = 1 - \frac{1}{r^\sigma} \text{ where } r = |y - \xi'_j|.$$

Then

$$-\Delta\psi_{1j} = \sigma^2 \frac{1}{r^{2+\sigma}}$$

If $\xi'_j \in \Omega_\varepsilon$, then we have

$$\frac{\partial\psi_{1j}}{\partial\nu_\varepsilon} = O(\varepsilon^{1+\sigma}) \text{ on } \partial\Omega_\varepsilon.$$

If $\xi'_j \in \Omega_\varepsilon$ and $|y - \xi'_j| > R$, we have

$$\frac{\partial\psi_{1j}}{\partial\nu_\varepsilon} = \sigma \frac{\langle y - \xi'_j, \nu_\varepsilon \rangle}{r^{2+\sigma}}$$

As before, we write $\partial\Omega_\varepsilon$ near ξ'_j as the graph $\{(y_1, y_2) : y_2 = \frac{1}{\varepsilon}G(\varepsilon y_1)\}$ with $G(0) = 0$ and $G'(0) = 0$. Then

$$\begin{aligned} \frac{\partial\psi_{1j}}{\partial\nu} &= \frac{\sigma}{r^{2+\sigma}} \frac{1}{\sqrt{G'(\varepsilon y_1)^2 + 1}} \left(-y_1 G'(\varepsilon y_1), \frac{1}{\varepsilon} G(\varepsilon y_1) \right) \\ &= \frac{\sigma}{r^{2+\sigma}} \frac{1}{\sqrt{O(\delta^2) + 1}} O(\varepsilon r^2) \quad \forall R_1 < r < \delta/\varepsilon \\ &= O\left(\frac{\varepsilon}{r^\sigma}\right) \quad \forall R_1 < r < \delta/\varepsilon. \end{aligned}$$

Combining together, we see that

$$(42) \quad \frac{\partial\psi_{1j}}{\partial\nu_\varepsilon} = o(\varepsilon) \text{ on } \partial\Omega_\varepsilon.$$

Now let ψ_0 be the unique solution of

$$\Delta\psi_0 - \varepsilon^2\psi_0 + \varepsilon^2 = 0 \text{ in } \Omega_\varepsilon, \quad \frac{\partial\psi_0}{\partial\nu_\varepsilon} = \varepsilon \text{ on } \partial\Omega_\varepsilon.$$

Set

$$(43) \quad \psi = \sum_{j=1}^m \psi_{1j} + C\psi_0$$

Then for $|y - \xi'_j| > R, j = 1, \dots, k+l$ where R is large

$$(44) \quad -\Delta\psi + \varepsilon^2\psi - W\psi \geq C\varepsilon^2 + \sigma^2 \sum_{j=1}^m \frac{1}{|y - \xi'_j|^{2+\sigma}} - CW \geq \frac{\sigma^2}{2} \sum_{j=1}^m \frac{1}{|y - \xi'_j|^{2+\sigma}} + \varepsilon^2$$

since

$$W \leq \sum_{j=1}^m \frac{1}{1 + |y - \xi'_j|^4}$$

On $\partial\Omega_\varepsilon$,

$$\frac{\partial\psi}{\partial\nu_\varepsilon} \geq \frac{\varepsilon}{2}$$

It is easy to see that $\frac{4}{\sigma^2}\psi$ satisfies all the properties of the lemma. \square

We will establish next an a priori estimate for solutions to problem (31) that satisfy orthogonality conditions with respect to $Z_{ij}, i = 1, J_j$ only.

Lemma 3.4 *For ε sufficiently small, if ϕ solves*

$$(45) \quad -\Delta\phi + \varepsilon^2\phi + W\phi = h \quad \text{in } \Omega_\varepsilon, \quad \frac{\partial\phi}{\partial\nu} = 0 \quad \text{on } \partial\Omega_\varepsilon$$

and satisfies

$$(46) \quad \int_{\Omega_\varepsilon} Z_{ij}\chi_j \phi = 0 \quad \forall j = 1, \dots, m, i = 1, J_j$$

then

$$(47) \quad \|\phi\|_{L^\infty(\Omega_\varepsilon)} \leq C \log \frac{1}{\varepsilon} \|h\|_*$$

where C is independent of ε .

Proof. Let ϕ satisfy (45) and (46). We will modify ϕ to satisfy all orthogonality relations in (39) and for this purpose we consider modifications with compact support of the functions Z_{0j} . Let $R > R_0 + 1$ be large and fixed.

Let

$$(48) \quad a_{0j} = \frac{1}{\mu_j \left(\frac{4}{c_j} \log \frac{1}{\varepsilon R} + H(\xi_j, \xi_j) \right)}.$$

Set

$$(49) \quad \widehat{Z}_{0j}(y) = Z_{0j}(y) - \frac{1}{\mu_j} + a_{0j}G(\xi_j, \varepsilon y).$$

Note that by our definition, $\widehat{Z}_{0,j}$ satisfies the Neumann boundary condition.

Let η be radial smooth cut-off function on \mathbb{R}^2 so that

$$0 \leq \eta \leq 1, \quad |\nabla\eta| \leq C \quad \text{in } \mathbb{R}^2, \quad \eta \equiv 1 \quad \text{in } B_R(0) \quad \text{and} \quad \eta \equiv 0 \quad \text{in } \mathbb{R}^2 \setminus B_{R+1}(0).$$

Set

$$(50) \quad \eta_j(y) = \eta(|y - \xi_j'|) \quad \text{for } j = 1, \dots, k, \quad \eta_j(y) = \eta(F_j^\varepsilon(y)) \quad \text{for } j = k+1, \dots, k+l.$$

Now define

$$(51) \quad \widetilde{Z}_{0j} = \eta_j Z_{0j} + (1 - \eta_j) \widehat{Z}_{0j}.$$

Given ϕ satisfying (45) and (46) let

$$\tilde{\phi} = \phi + \sum_{j=1}^m d_j \tilde{Z}_{0j}, \quad \text{where } d_j = -\frac{\int_{\Omega_\varepsilon} Z_{0j} \chi_j \phi}{\int_{\Omega_\varepsilon} Z_{0j}^2 \chi_j}.$$

Estimate (47) is a direct consequence of:

Claim.

$$(52) \quad |d_j| \leq C \log \frac{1}{\varepsilon} \|h\|_* \quad \forall j = 1, \dots, m.$$

We start proving this by observing, using the notation $L = -\Delta + \varepsilon^2 - W$, that

$$(53) \quad L(\tilde{\phi}) = h + \sum_{j=1}^m d_j L(\tilde{Z}_{0j}) \quad \text{in } \Omega_\varepsilon,$$

and

$$(54) \quad \frac{\partial \tilde{\phi}}{\partial \nu} = 0 \quad \text{on } \partial\Omega_\varepsilon.$$

Thus by lemma 3.2 we have

$$(55) \quad \|\tilde{\phi}\|_{L^\infty(\Omega_\varepsilon)} \leq C \sum_{j=1}^m |d_j| \|L(\tilde{Z}_{0j})\|_* + C \|h\|_*$$

Multiplying equation (53) by \tilde{Z}_{0k} , integrating by parts and using (54) we find

$$(56) \quad \sum_{j=1}^m d_j \int_{\Omega_\varepsilon} L(\tilde{Z}_{0j}) \tilde{Z}_{0k} \leq C \|h\|_* [1 + \sum_{j=1}^m \|L(\tilde{Z}_{0j})\|_*] + C \sum_{j=1}^m |d_j| \|L(\tilde{Z}_{0j})\|_*^2$$

We now measure the size of $\|L(\tilde{Z}_{0j})\|_*$. To this end, we have for $|y - \xi'_j| > R$, according to (37)

$$(57) \quad L(\hat{Z}_{0j}) = -e^{v_j} Z_{0j} + W \hat{Z}_{0j} + O(\varepsilon(1 + |y - \xi'_j|)^{-3}) = e^{v_j} (a_{0j} G(\xi_j, \varepsilon y) - \frac{1}{\mu_j}) + O(\varepsilon(1 + |y - \xi'_j|)^{-3})$$

Thus

$$(58) \quad \|(1 - \eta_j) L(\hat{Z}_{0j})\|_* \leq \frac{C}{\log \frac{1}{\varepsilon}}$$

where the number C depends in principle of the chosen large constant R .

So

$$(59) \quad \begin{aligned} L(\tilde{Z}_{0j}) &= \eta_j L(Z_{0j}) + (1 - \eta_j) L(\hat{Z}_{0j}) + 2\nabla \eta_j \nabla (Z_{0j} - \hat{Z}_{0j}) + \Delta \eta_j (Z_{0j} - \hat{Z}_{0j}) \\ &= O(\varepsilon^{2+\alpha}) + (1 - \eta_j) e^{v_j} (a_{0j} G(\xi_j, \varepsilon y) - \frac{1}{\mu_j}) + 2\nabla \eta_j \nabla (Z_{0j} - \hat{Z}_{0j}) + \Delta \eta_j (Z_{0j} - \hat{Z}_{0j}). \end{aligned}$$

Note that for $r = |y - \xi'_j| \in (R, R + 1)$, we have

$$\begin{aligned}\widehat{Z}_{0j} - Z_{0j} &= a_{0j}G(\xi_j, \varepsilon y) - \frac{1}{\mu_j} \\ &= a_{0j} \left(\frac{4}{c_j} \log \frac{1}{\varepsilon |\xi'_j - y|} + H(\xi_j, \varepsilon y) \right) - \frac{1}{\mu_j}\end{aligned}$$

Henc we derive that for $r \in (R, R + 1)$,

$$(60) \quad \widehat{Z}_{0j} - Z_{0j} = \frac{C}{\log \frac{1}{\varepsilon}} \log \frac{1}{r} + O\left(\frac{\varepsilon^\alpha}{\log \frac{1}{\varepsilon}}\right), \quad \nabla(\widehat{Z}_{0j} - Z_{0j}) = -\frac{C}{\log \frac{1}{\varepsilon}} \frac{1}{r} + O\left(\frac{\varepsilon^\alpha}{\log \frac{1}{\varepsilon}}\right).$$

From (59) and (60), we conclude that

$$(61) \quad \|L(\widetilde{Z}_{0j})\|_* \leq \frac{C}{\log \frac{1}{\varepsilon}}.$$

Now we estimate the left hand side integral of (56). From (59), we see that for $j \neq k$,

$$\int_{\Omega_\varepsilon} L(\widetilde{Z}_{0j})\widetilde{Z}_{0k} = O(\varepsilon^\alpha) + \int_{\Omega_\varepsilon} O\left(\frac{1}{\log \frac{1}{\varepsilon}}(|\eta'_j| + |\Delta\eta_j|)\right)\widetilde{Z}_{0,k} = O\left(\left(\frac{1}{\log \frac{1}{\varepsilon}}\right)^2\right)$$

For $j = k$, we decompose

$$\int_{\Omega_\varepsilon} L(\widetilde{Z}_{0k})\widetilde{Z}_{0k} = I + II + O(\varepsilon)$$

where

$$\begin{aligned}II &= \int_{\Omega_\varepsilon} O(\varepsilon^{2+\alpha}) + (1 - \eta_k)e^{v_j} \left(a_{0k}G(\xi_k, \varepsilon y) - \frac{1}{\mu_k} \right) \widetilde{Z}_{0k} \\ &= O(\varepsilon^\alpha) + O\left(\frac{1}{R} \frac{1}{\log \frac{1}{\varepsilon}}\right)\end{aligned}$$

and

$$I = \int_{\Omega_\varepsilon} (2\nabla\eta_k \nabla(Z_{0k} - \widehat{Z}_{0k}) + \Delta\eta_k(Z_{0k} - \widehat{Z}_{0k}))\widetilde{Z}_{0k}$$

Thus integrating by parts we find

$$I = \int \nabla\eta \nabla(Z_{0k} - \widehat{Z}_{0k})\widehat{Z}_{0k} - \int \nabla\eta_j(Z_{0k} - \widehat{Z}_{0k})\nabla\widehat{Z}_{0k} + O(\varepsilon)$$

Now, we observe that in the considered region, $r \in (R, R + 1)$ with $r = |y - \xi'_k|$, $|\widehat{Z}_{0k} - Z_{0k}| \leq \frac{C}{\log \frac{1}{\varepsilon}}$ while $|\nabla Z'_{0k}| \leq \frac{1}{R^3} + \frac{1}{R \log \frac{1}{\varepsilon}}$. Thus

$$\left| \int \nabla\eta_j(Z_{0k} - \widehat{Z}_{0k})\nabla\widehat{Z}_{0k} \right| \leq \frac{D}{R^3} \frac{1}{\log \frac{1}{\varepsilon}}$$

where D may be chosen independent of R . Now

$$\begin{aligned} \int \nabla \eta_k \nabla (Z_{0k} - \widehat{Z}_{0k}) \widehat{Z}_{0k} &= \int_R^{R+1} \eta' \left(a_{0k} \frac{1}{r} + O(\varepsilon) \right) \widehat{Z}_{0k} r dr \\ &= a_{0k} \int_R^{R+1} \eta' \left(1 + O(\varepsilon) + O\left(\frac{1}{R}\right) \right) \\ &= -\frac{E}{\log \frac{1}{\varepsilon}} \left[1 + O\left(\frac{1}{R}\right) \right] \end{aligned}$$

where E is a positive constant independent of ε . Thus we conclude, choosing R large enough, that $I \sim -\frac{E}{\log \frac{1}{\varepsilon}}$. Combining this and the estimate for II we find

$$(62) \quad \int_{\Omega_\varepsilon} L(\widetilde{Z}_{0k}) \widetilde{Z}_{0k} = -\frac{E}{\log \frac{1}{\varepsilon}} \left[1 + O\left(\frac{1}{R}\right) \right], \quad \int_{\Omega_\varepsilon} L(\widetilde{Z}_{0j}) \widetilde{Z}_{0k} = O\left(\frac{1}{R} \frac{1}{\log \frac{1}{\varepsilon}}\right) \quad \text{for } j \neq k$$

This, combined with (56), proves the lemma. \square

Proof of proposition 3.1.

First we prove that for any ϕ, d_1, \dots, d_m solution to (31) the bound

$$(63) \quad \|\phi\|_{L^\infty(\Omega_\varepsilon)} \leq C \log \frac{1}{\varepsilon} \|h\|_*$$

holds.

The previous lemma yields

$$(64) \quad \|\phi\|_{L^\infty(\Omega_\varepsilon)} \leq C \log \frac{1}{\varepsilon} (\|h\|_* + \sum_{j=1}^m \sum_{i=1}^{J_j} |c_{ij}|)$$

So it suffices to estimate the values of the constants c_{ij} .

To this end, we multiple (31) by Z_{ij} and integrate to find

$$(65) \quad \int_{\Omega_\varepsilon} L(\phi)(Z_{ij}) = \int_{\Omega_\varepsilon} h Z_{ij} + c_{ij} \int_{\Omega_\varepsilon} \chi_j |Z_{ij}|^2$$

Note that for $i \neq 0$

$$Z_{ij} = O\left(\frac{1}{1 + |y - \xi_j|}\right)$$

So

$$(66) \quad \int_{\Omega_\varepsilon} h Z_{ij} = O(\|h\|_*)$$

and

$$(67) \quad \int_{\Omega_\varepsilon} L(\phi) Z_{ij} = \int_{\Omega_\varepsilon} L(Z_{ij}) \phi = O(\varepsilon \|\phi\|_\infty)$$

Substituting (66) and (67) into (65), we obtain (64).

Now consider the Hilbert space

$$H = \left\{ \phi \in H^1(\Omega_\varepsilon) : \int_{\partial\Omega_\varepsilon} \chi_j Z_{ij} \phi = 0 \quad \forall j = 1, \dots, m, i = 1, J_j \right\}$$

with the norm $\|\phi\|_{H^1}^2 = \int_{\Omega_\varepsilon} |\nabla\phi|^2 + \varepsilon^2\phi^2$. Equation (31) is equivalent to find $\phi \in H$ such that

$$\int_{\Omega_\varepsilon} (\nabla\phi\nabla\psi + \varepsilon^2\phi\psi) - \int_{\Omega_\varepsilon} W\phi\psi = \int_{\partial\Omega_\varepsilon} h\psi \quad \forall\psi \in H.$$

By Fredholm's alternative this is equivalent to the uniqueness of solutions to this problem, which is guaranteed by (63). \square

The result of Proposition 3.1 implies that the unique solution $\phi = T(h)$ of (31) defines a continuous linear map from the Banach space C_* of all functions h in L^∞ for which $\|h\|_* < \infty$, into L^∞ .

It is important for later purposes to understand the differentiability of the operator T with respect to the variables $\xi'_{k,l}$. Fix $h \in C_*$ and let $\phi = T(h)$. We want to compute derivatives of ϕ with respect to, say, $\xi'_{k,l}$. Similar to that of [11], we obtain the following estimate

$$(68) \quad \|\partial_{\xi'_{k,l}} T(h)\|_\infty \leq C \left(\log \frac{1}{\varepsilon}\right)^2 \|h\|_*, \text{ for all } k = 1, \dots, m, l = 1, J_k.$$

4 The nonlinear problem

Consider the nonlinear equation

$$(69) \quad \begin{cases} -\Delta\phi + \varepsilon^2\phi - W\phi = R + N(\phi) + \sum_{ij} c_{ij}\chi_j Z_{ij} & \text{in } \Omega_\varepsilon \\ \frac{\partial\phi}{\partial\nu} = 0 & \text{on } \partial\Omega_\varepsilon \\ \int_{\Omega_\varepsilon} \chi_j Z_{ij} \phi = 0 & \forall j = 1, \dots, m, i = 1, J_j \end{cases}$$

where W is as in (21) and N, R are defined in (17) and (18) respectively.

Lemma 4.1 *Let $m > 0, d > 0$. Then there exist $\varepsilon_0 > 0, C > 0$ such that for $0 < \varepsilon < \varepsilon_0$ and any $(\xi_1, \dots, \xi_m) \in \mathcal{M}_\delta$ the problem (69) admits a unique solution ϕ, c_1, \dots, c_m such that*

$$(70) \quad \|\phi\|_{L^\infty(\Omega_\varepsilon)} \leq C\varepsilon^\alpha |\log \varepsilon|$$

where α is any number in the interval $(0, 1)$. Furthermore, the function $\xi' \rightarrow \phi(\xi') \in C(\bar{\Omega}_\varepsilon)$ is C^1 and

$$(71) \quad \|D_{\xi'} \phi\|_{L^\infty(\Omega_\varepsilon)} \leq C \varepsilon^\alpha |\log \varepsilon|^2.$$

Proof.

The proof of this lemma can be done along the lines of those of Lemma 4.1 of [11]. We omit the details. \square

5 Variational reduction

In view of lemma 4.1, given $\xi = (\xi_1, \dots, \xi_m) \in \mathcal{M}_\delta$, we define $\phi(\xi)$ and $c_{ij}(\xi)$ to be the unique solution to (69) satisfying the bound (70).

Given $\xi = (\xi_1, \dots, \xi_m) \in \Omega^k \times (\partial\Omega)^l$ we write

$$U(\xi) = \sum_{j=1}^m (u_j(x) + H_j^\varepsilon(x))$$

the ansatz defined in (10). Set

$$(72) \quad F_\varepsilon(\xi) = J_\varepsilon(U(\xi) + \tilde{\phi}(\xi)),$$

where J_ε is the functional defined by

$$(73) \quad J_\varepsilon(v) = \frac{1}{2} \int_{\Omega} (|\nabla v|^2 + v^2) - \varepsilon^2 \int_{\Omega} e^v$$

and

$$(74) \quad \tilde{\phi}(\xi)(x) = \phi(\xi)\left(\frac{x}{\varepsilon}\right), \quad x \in \Omega.$$

Lemma 5.1 *If $\xi = (\xi_1, \dots, \xi_m) \in \mathcal{M}_\delta$ is a critical point of F_ε then $u = U(\xi) + \tilde{\phi}(\xi)$ is a critical point of J_ε , that is, a solution to (4).*

Proof. Let

$$I_\varepsilon(v) = \frac{1}{2} \int_{\Omega_\varepsilon} |\nabla v|^2 + \varepsilon^2 v^2 - \varepsilon^4 \int_{\Omega_\varepsilon} e^v.$$

Then $F_\varepsilon(\xi) = J_\varepsilon(U(\xi) + \tilde{\phi}(\xi)) = I_\varepsilon(V(\xi') + \phi(\xi'))$ where $\xi' = \xi/\varepsilon$. Therefore

$$\frac{\partial F_\varepsilon}{\partial \xi_{k,l}} = \frac{1}{\varepsilon} \frac{\partial I_\varepsilon(V(\xi') + \phi(\xi'))}{\partial \xi'_{k,l}} = \frac{1}{\varepsilon} D I_\varepsilon(V(\xi') + \phi(\xi')) \left[\frac{\partial V(\xi')}{\partial \xi'_{k,l}} + \frac{\partial \phi(\xi')}{\partial \xi'_{k,l}} \right].$$

Since $v = V(\xi') + \phi(\xi')$ solves (69)

$$\frac{\partial F_\varepsilon}{\partial \xi_{k,l}} = \frac{1}{\varepsilon} \sum_{i=1, j=1, \dots, m} c_{ij} \int_{\partial\Omega_\varepsilon} \chi_j Z_{ij} \left[\frac{\partial V(\xi')}{\partial \xi'_{k,l}} + \frac{\partial \phi(\xi')}{\partial \xi'_{k,l}} \right].$$

Let us assume that $DF(\xi) = 0$. From the previous equation we conclude that

$$\sum_{j=1, \dots, m, i=1, J_j} c_{ij} \int_{\Omega_\varepsilon} \chi_j Z_{ij} \left[\frac{\partial V(\xi')}{\partial \xi'_{k,l}} + \frac{\partial \phi(\xi')}{\partial \xi'_{k,l}} \right] = 0 \quad \forall k = 1, \dots, m, l = 1, J_k.$$

Since $\left\| \frac{\partial \phi(\xi')}{\partial \xi'_{k,l}} \right\|_{L^\infty(\Omega_\varepsilon)} \leq C\varepsilon^\alpha |\log \varepsilon|^2$ and $\frac{\partial V(\xi')}{\partial \xi'_{k,l}} = \pm Z_{kl} + o(1)$ where $o(1)$ is in the L^∞ norm, it follows that

$$\sum_{j=1, \dots, m, J=1, J_j} c_{ij} \int_{\partial \Omega_\varepsilon} \chi_j Z_{ij} (\pm Z_{kl} + o(1)) \quad \forall k = 1, \dots, m,$$

which is a strictly diagonal dominant system. This implies that $c_{ij} = 0 \forall i = 1, \dots, m, j = 1, J_j$. \square

In order to solve for critical points of the function F , a key step is its expected closeness to the function $J_\varepsilon(U)$, which we will analyze in the next section.

Lemma 5.2 *The following expansion holds*

$$F_\varepsilon(\xi) = J_\varepsilon(U) + \theta_\varepsilon(\xi),$$

where

$$|\theta_\varepsilon| + |\nabla \theta_\varepsilon| \rightarrow 0,$$

uniformly on \mathcal{M}_δ .

Proof. Let $\tilde{\theta}(\xi') = I_\varepsilon(V + \phi) - I_\varepsilon(V)$. In order to get the proof of this lemma, we need to show that

$$|\tilde{\theta}| + \varepsilon^{-1} |\nabla_{\xi'} \tilde{\theta}| = o(1).$$

Taking into account $DI_\varepsilon(V + \phi)[\phi] = 0$, a Taylor expansion and an integration by parts give

$$\begin{aligned} I_\varepsilon(V + \phi) - I_\varepsilon(V) &= \int_0^1 D^2 I_\varepsilon(V + t\phi)[\phi]^2 (1-t) dt \\ (75) \quad &= \int_0^1 \left(\int_{\Omega_\varepsilon} [N(\phi) + R] \phi + \int_{\Omega_\varepsilon} e^V [1 - e^{t\phi}] \phi^2 \right) (1-t) dt, \end{aligned}$$

so we get

$$I_\varepsilon(V + \phi) - I_\varepsilon(V) = \tilde{\theta}_\varepsilon = O(\varepsilon^{2\alpha} |\log \varepsilon|^3).$$

taking into account that $\|\phi\|_{L^\infty(\Omega_\varepsilon)} \leq C\varepsilon^\alpha |\log \varepsilon|$. Let us differentiate with respect to $\xi'_{k,l}$

$$\partial_{\xi'_{k,l}} [I_\varepsilon(V + \phi) - I_\varepsilon(V)] = \int_0^1 \left(\int_{\partial \Omega_\varepsilon} \partial_{\xi'_{k,l}} [(N(\phi) + R) \phi] \right)$$

$$+ \int_{\Omega_\varepsilon} \partial_{\xi'_k, l} [e^V [1 - e^{t\phi}] \phi^2] (1-t) dt.$$

Using the fact that $\|\partial_{\xi'_k} \phi\|_* \leq C \varepsilon^\alpha |\log \varepsilon|^2$ and the estimates of the previous sections we get

$$\partial_{\xi'_k, l} [I_\varepsilon(V + \phi) - I_\varepsilon(V)] = \partial_{\xi'_k, l} \tilde{\theta}_\varepsilon = O(\varepsilon^{2\alpha} |\log \varepsilon|^4).$$

The continuity in ξ of all these expressions is inherited from that of ϕ and its derivatives in ξ in the L^∞ norm. The proof is complete. \square

6 Expansion of the energy

Lemma 6.1 *Let μ_j be given by (19). Then for any $0 < \alpha < 1$*

$$J_\varepsilon(U) = (8\pi k + 4\pi l)(\beta - 1 + \log 8) + 2(8\pi k + 4\pi l) \log \frac{1}{\varepsilon} - \frac{1}{2} \sum_{j=1}^m c_j \left[c_j H(\xi_j, \xi_j) + \sum_{i, i \neq j} c_i G(\xi_i, \xi_j) \right] + O(\varepsilon^\alpha).$$

where

$$\beta = \int_0^\infty \frac{1}{(1+x^2)^2} \log \frac{1}{(1+x^2)^2} x dx.$$

Proof. Define

$$U_j(x) = u_j(x) + H_j^\varepsilon(x)$$

so we may rewrite (10) in equivalent form $U = \sum_{j=1}^m U_j$. Then

$$\begin{aligned} J_\varepsilon(U) &= \frac{1}{2} \int_\Omega \left| \sum_{j=1}^m \nabla U_j \right|^2 + \frac{1}{2} \int_\Omega \left(\sum_{j=1}^m U_j \right)^2 - \varepsilon^2 \int_\Omega \exp\left(\sum_{j=1}^m U_j \right) \\ &= \sum_{j=1}^m \int_\Omega \frac{1}{2} (|\nabla U_j|^2 + U_j^2) + \frac{1}{2} \sum_{i \neq j} \int_\Omega (\nabla U_i \nabla U_j + U_i U_j) - \varepsilon^2 \int_\Omega \exp\left(\sum_{j=1}^m U_j \right) \\ &= I_A + I_B + I_C. \end{aligned}$$

Let us analyze the behavior of I_A . Note that U_j satisfies

$$\Delta U_j - U_j + \varepsilon^2 e^{u_j} = 0 \text{ in } \Omega, \quad \frac{\partial U_j}{\partial \nu} = 0 \text{ on } \partial\Omega$$

which gives

$$(76) \quad \int_\Omega (|\nabla U_j|^2 + U_j^2) = \varepsilon^2 \int_\Omega e^{u_j} (u_j + H_j^\varepsilon).$$

Let us find the asymptotic behavior of the expression:

$$\int_{\Omega} (|\nabla U_j|^2 + U_j^2) = \varepsilon^2 \int_{\Omega} \frac{8\mu_j^2}{(\varepsilon^2\mu_j^2 + |x - \xi_j|^2)^2} \left(\log \frac{8\mu_j^2}{(\varepsilon^2\mu_j^2 + |x - \xi_j|^2)^2} + c_j H(x, \xi_j) + O(\varepsilon^\alpha) \right).$$

Changing variables $\varepsilon\mu_j y = x - \xi_j$

$$\int_{\Omega} |\nabla U_j|^2 + U_j^2 = \int_{\Omega_{\varepsilon\mu_j}} \frac{8}{(1 + |y|^2)^2} \left(\log \frac{1}{(1 + |y|^2)^2} + c_j H(\xi_j + \varepsilon\mu_j y, \xi_j) - 4 \log(\varepsilon\mu_j) \right) + O(\varepsilon^\alpha).$$

But

$$\int_{\Omega_{\varepsilon\mu_j}} \frac{8}{(1 + |y|^2)^2} = 2c_j + O(\varepsilon), \quad \int_{\Omega_{\varepsilon\mu_j}} \frac{8}{(1 + |y|^2)^2} \log \frac{1}{(1 + |y|^2)^2} = c_j\beta + O(\varepsilon^\alpha)$$

and for $0 < \alpha < 1$

$$\begin{aligned} \int_{\Omega_{\varepsilon\mu_j}} \frac{8}{(1 + |y|^2)^2} (H(\varepsilon\mu_j y, \xi_j) - H(\xi_j, \xi_j)) &= \int_{\Omega_{\varepsilon\mu_j}} \frac{1}{(1 + |y|^2)^2} O(\varepsilon^\alpha |y|^\alpha) \\ &= O(\varepsilon^\alpha). \end{aligned}$$

Therefore

$$(77) \quad \int_{\Omega} |\nabla U_j|^2 + U_j^2 = 2c_j\beta + c_j^2 H(\xi_j, \xi_j) - 4c_j \log(\varepsilon\mu_j) + O(\varepsilon^\alpha).$$

Thus

$$(78) \quad I_A = \sum_j c_j\beta - 2 \sum_{j=1}^m c_j \log(\varepsilon\mu_j) + \sum_{j=1}^m \frac{1}{2} c_j^2 H(\xi_j, \xi_j) + O(\varepsilon^\alpha).$$

We consider now

$$\begin{aligned} I_B &= \frac{1}{2} \sum_{i \neq j}^m \int_{\Omega} (\nabla U_i \nabla U_j + U_i U_j) \\ &= \frac{\varepsilon^2}{2} \sum_{i \neq j} \int_{\Omega} e^{u_i} (u_j + H_j) \end{aligned}$$

A similar argument as for I_A shows that

$$(79) \quad I_B = \frac{1}{2} \sum_{i \neq j}^m c_i c_j G(\xi_i, \xi_j) + O(\varepsilon^\alpha).$$

Regarding the expression I_C we have

$$I_C = -\varepsilon^2 \int_{\Omega} e^{\sum_{k=1}^m U_k} = -\varepsilon^2 \sum_{j=1}^m \int_{\Omega \cap B_{\delta}(\xi_j)} e^{\sum_{k=1}^m (u_k + H_k^{\varepsilon})} + O(\varepsilon^2).$$

Using the definition of u_j and (12) for each term we have

$$\varepsilon^2 \int_{\Omega \cap B_{\delta}(\xi_j)} e^{\sum_{k=1}^m (u_k + H_k^{\varepsilon})} = \varepsilon^2 \int_{\partial \Omega \cap B_{\delta}(\xi_j)} e^{u_j} e^{c_j H(x, \xi_j) - \log(8\mu_j^2) + O(\varepsilon^{\alpha})} E_j(x),$$

where

$$E_j(x) = \exp \left(\sum_{i \neq j} \log \frac{1}{(\varepsilon^2 \mu_i^2 + |x - \xi_i|^2)^2} + c_i H(x, \xi_i) + O(\varepsilon^{\alpha}) \right).$$

Changing variables $\varepsilon \mu_j y = x - \xi_j$ we have

$$e^{c_j H(\xi_j + \varepsilon \mu_j y, \xi_j) - \log(8\mu_j^2) + O(\varepsilon^{\alpha})} = e^{c_j H(\xi_j, \xi_j) - \log(8\mu_j^2)} + O(\varepsilon^{\alpha} |y|^{\alpha}).$$

and

$$E_j(\xi_j + \varepsilon \mu_j y, \xi_j) = \exp \left(\sum_{i, i \neq j} c_i G(\xi_j, \xi_i) \right) + O(\varepsilon^{\alpha} |y|^{\alpha}).$$

Therefore, by the definition of μ_j in (19)

$$\begin{aligned} \varepsilon^2 \int_{\Omega \cap B_{\delta}(\xi_j)} e^{\sum_{k=1}^m (u_k + H_k^{\varepsilon})} &= \varepsilon^2 \int_{\Omega \cap B_{\delta}(\xi_j)} e^{u_j + O(\varepsilon^{\alpha})} \\ &= c_j + O(\varepsilon^{\alpha}). \end{aligned}$$

Thus

$$(80) \quad I_C = - \sum_j c_j + O(\varepsilon^{\alpha}).$$

Thanks to (78), (79) and (80) we have

$$\begin{aligned} J_{\varepsilon}(U) &= \sum_{j=1}^m c_j (\beta - 1 + \log 8) + 2 \sum_{j=1}^m c_j \log \frac{1}{\varepsilon} + \sum_{j=1}^m c_j \left[-\log(8\mu_j^2) + \frac{1}{2} c_j H(\xi_j, \xi_j) \right. \\ &\quad \left. + \frac{1}{2} \sum_{i \neq j} c_i c_j G(\xi_i, \xi_j) \right] + O(\varepsilon^{\alpha}). \end{aligned}$$

Employing again (19) we have

$$\begin{aligned} J_{\varepsilon}(U) &= \sum_{j=1}^m c_j (\beta - 1 + \log 8) + 2 \sum_{j=1}^m c_j \log \frac{1}{\varepsilon} - \frac{1}{2} \sum_{j=1}^m c_j \left[\sum_{j=1}^m c_j H(\xi_j, \xi_j) + \sum_{i, i \neq j} c_i G(\xi_i, \xi_j) \right] \\ &\quad + O(\varepsilon^{\alpha}). \end{aligned}$$

□

7 Proof of theorem 1.2

Let

$$(81) \quad \varphi_m(\xi) = \sum_{j=1}^m c_j \left[c_j H(\xi_j, \xi_j) + \sum_{i, i \neq j} c_i G(\xi_i, \xi_j) \right].$$

We have

Lemma 7.1 *We have*

$$(82) \quad \min_{\xi \in \partial \mathcal{M}_\delta} \rightarrow +\infty \text{ as } \delta \rightarrow 0.$$

Proof: Let $\xi = (\xi_1, \dots, \xi_m) \in \partial \mathcal{M}_\delta$. There are two possibilities: either there exists $j_0 \leq k$ such that $d(\xi_{j_0}, \partial \Omega) = \delta$, or there exists $i_0 \neq j_0$, $|\xi_{i_0} - \xi_{j_0}| = \delta$.

In the first case, we claim that for all $\xi \in \Omega$

$$(83) \quad H(\xi, \xi) \geq C \log \frac{1}{Cd(\xi, \partial \Omega)}.$$

In fact, if ξ is close to the boundary, let ξ_0 be the nearest point of $\partial \Omega$ to ξ . It is easily checked that

$$(84) \quad H(x, \xi) = \frac{1}{2\pi} \log \frac{1}{|x - \xi^*|} + O(1) \quad \text{as } d(\xi, \partial \Omega) \rightarrow 0$$

uniformly in Ω , where ξ^* is the reflection of ξ across the boundary, that is the symmetric point to ξ with respect to ξ_0

(83) follows from (84), using the fact that $G(x, y) > 0$.

In the second case, we may assume that there exists a fixed constant C such that $d(\xi_i, \partial \Omega) \geq C$, $i = 1, \dots, k$, as otherwise it follows into the first case. But then it is easy to see that

$$(85) \quad G(\xi_i, \xi_j) \geq C \log \frac{1}{|\xi_i - \xi_j|}$$

From (83) and (85), the proof of this lemma is complete. \square

Proof of Theorem 1.2: For $\delta > 0$ sufficiently small, we define a configuration space as:

$$(86) \quad \mathcal{M}_\delta := \left\{ \xi = (\xi_1, \dots, \xi_m) \in (\Omega)^k \times (\partial \Omega)^L \mid \min_{i=1, \dots, k} d(\xi_i, \partial \Omega) \geq \delta, \min_{i \neq j} |\xi_i - \xi_j| \geq \delta \right\}.$$

According to Lemma 6.2, the function $U(\xi) + \tilde{\phi}(\xi)$, where U and $\tilde{\phi}$ are defined respectively by (10) and (74), is a solution of Problem (3) if we adjust ξ so that it is a critical point of $F_\varepsilon(\xi) = J_\varepsilon(U(\xi) + \tilde{\phi}(\xi))$ defined by (72). This is obviously equivalent to finding a critical point of

$$\tilde{F}_\varepsilon(\xi) = -2 \left(F_\varepsilon(\xi) - (8\pi k + 4\pi l)(\beta - 1 + \log 8) - 2(8\pi k + 4\pi l) \log \frac{1}{\varepsilon} \right)$$

On the other hand, from Lemmas 5.2 and 6.1, we have that for $\xi \in \mathcal{M}_\delta$,

$$(87) \quad \tilde{F}_\varepsilon(\xi) = \varphi_m(\xi) + \varepsilon^\alpha \Theta_\varepsilon(\xi)$$

where φ_m is give by (81), Θ_ε and $\nabla_\xi \Theta_\varepsilon$ are uniformly bounded in the considered region as $\varepsilon \rightarrow 0$.

From the above Lemma, the function φ_m is C^1 , bounded from below in \mathcal{M}_δ and such that

$$\varphi_m(\xi_1, \dots, \xi_m) \rightarrow +\infty \text{ as } \delta \rightarrow 0$$

Hence, for δ is arbitrarily small, φ_m has an absolute minimum M in \mathcal{M}_δ . This implies that $\tilde{F}_\varepsilon(\xi)$ also has an absolute minimum $(\xi_1^\varepsilon, \dots, \xi_m^\varepsilon) \in \mathcal{M}_\delta$ such that

$$\varphi_m(\xi_1^\varepsilon, \dots, \xi_m^\varepsilon) \rightarrow \min_{\xi \in \mathcal{M}_\delta} \varphi_m(\xi) \text{ as } \varepsilon \rightarrow 0,$$

Hence Lemma 5.1 guarantees the existence of a solution u_ε for (4). Furthermore, from the ansatz (10), we get that, as $\varepsilon \rightarrow 0$, u_ε remains uniformly bounded on $\Omega \setminus \cup_{j=1}^m B_\delta(\xi_j^\varepsilon)$, and

$$\sup_{B_\delta(\xi_j^\varepsilon)} u_\varepsilon \rightarrow +\infty,$$

for any $\delta > 0$. □

Remark 7.2 By using Ljusternik-Schnirelmann theory, one can get a second, distinct solution satisfying Theorem 1.2. The proof is similar to [12].

Remark 7.3 As mentioned in the introduction, one can get a stronger result than Theorem 1.2 under the assumption that the function φ_m has, in addition to the ones described in the proof of Theorem 1.2, some other critical points in $\hat{\Omega}_m$ with the property of being *topologically non trivial*, for instance (possibly degenerate) local minima or maxima, or saddle points.

Let us define what we mean with *topologically non trivial* critical point for φ_m . Let Σ be an open set compactly contained in \mathcal{M}_δ with smooth boundary. We recall that φ_m *links in Σ at critical level \mathcal{C} relative to B and B_0* if B and B_0 are closed subsets of $\bar{\Sigma}$ with B connected and $B_0 \subset B$ such that the following conditions hold: Let us set Γ to be the class of all maps $\Phi \in C(B, \Sigma)$ with the property that there exists a function $\Psi \in C([0, 1] \times B, \Sigma)$ such that

$$\Psi(0, \cdot) = \text{Id}_B, \quad \Psi(1, \cdot) = \Phi, \quad \Psi(t, \cdot)|_{B_0} = \text{Id}_{B_0} \text{ for all } t \in [0, 1].$$

We assume

$$(88) \quad \sup_{y \in B_0} \varphi_m(y) < \mathcal{C} \equiv \inf_{\Phi \in \Gamma} \sup_{y \in B} \varphi_m(\Phi(y)),$$

and for all $y \in \partial\Sigma$ such that $\varphi_m(y) = \mathcal{C}$, there exists a vector τ_y tangent to $\partial\Sigma$ at y such that

$$(89) \quad \nabla \varphi_m(y) \cdot \tau_y \neq 0.$$

Under these conditions a critical point $\bar{y} \in \Sigma$ of φ_m with $\varphi_m(\bar{y}) = \mathcal{C}$ exists. Not only this: any function C^1 close to φ_m inherits such critical point.

Acknowledgment

The first author has been partly supported by grants Fondecyt 1030840 and FONDAP, Chile. The research of the second author is supported by an Earmarked Grant from RGC of Hong Kong (RGC402503).

References

- [1] S. Baraket, F. Pacard, Construction of singular limits for a semilinear elliptic equation in dimension 2. *Calc. Var. Partial Differential Equations* **6** (1998), no. 1, 1–38.
- [2] P. Bates, E.N. Dancer and J. Shi, Multi-spike stationary solutions of the Cahn-Hilliard equation in higher-dimension and instability, *Adv. Diff. Eqns* **4** (1999), 1–69.
- [3] P. Biler, Local and global solvability of some parabolic system modeling chemotaxis, *Adv. Math. Sci. Appl.* **8**(1998), 715–743.
- [4] M.P. Brenner, P. Constantin, L.P. Kadanoff, A. Schenkel, S.C. Venkataramani, Diffusion, attraction and collapse, *Nonlinearity* **12** (4) (1999) 1071–1098.
- [5] H. Brezis, F. Merle, Uniform estimates and blow-up behavior for solutions of $-\Delta u = V(x)e^u$ in two dimensions. *Comm. Partial Differential Equations* **16** (1991), no. 8–9, 1223–1253.
- [6] C.-C. Chen and C.-S. Lin, Sharp estimates for solutions of multi-bubbles in compact Riemann surfaces, *Comm. Pure Appl. Math.* **55** (2002), no. 6, 728–771.
- [7] S. Childress, Chemotactic Collapse in Two Dimensions, Lecture Notes in Biomathematics, Vol. 55, Springer, Berlin, 1984, pp. 61–68.
- [8] S. Childress and J. K. Percus, Nonlinear aspects of chemotaxis, *Math. Biosci.* **56** (1981), 217–237.
- [9] L. Corrias, B. Perthame, H. Zaag, Global solutions of some chemotaxis and angiogenesis systems in high space dimensions. *Milan J. Math.* **72** (2004), 1–28.
- [10] E.N. Dancer and S. Yan, Multipole solutions for a singular perturbed Neumann problem, *Pacific J. Math.* **189**(1999), 241–262.
- [11] M. Del Pino, M. Kowalczyk and M. Musso, Singular limits in Liouville-type equations. *Calc. Var. Partial Differential Equations*, to appear.

- [12] J. Dávila, M. del Pino, M. Musso, Concentrating solutions in a two-dimensional elliptic problem with exponential Neumann data, *To appear J. Funct. Anal.*
- [13] Juan Dávila, M. del Pino, M. Musso and J. Wei Singular limits of a two-dimensional boundary value problem arising in corrosion modelling, *Preprint*.
- [14] M. Del Pino, P. Felmer, M. Musso, Two-bubble solutions in the supercritical Bahri-Coron's problem, *Calc. Var. and Part. Diff. Equ.* **16** (2003) 113-145.
- [15] J. Dolbeault, B. Perthame, Optimal critical mass in the two-dimensional Keller-Segel model in \mathbb{R}^2 . *C. R. Math. Acad. Sci. Paris* 339 (2004), no. 9, 611–616.
- [16] P. Esposito, M. Grossi, A. Pistoia, On the existence of blowing-up solutions for a mean field equation. *Ann. Inst. H. Poincaré Anal. Non Linéaire*, to appear.
- [17] I. Guerra, M. Peletier, Self-similar blow-up for a diffusion-attraction problem. *Nonlinearity* 17 (2004), no. 6, 2137–2162.
- [18] C. Gui and J. Wei, Multiple interior spike solutions for some singular perturbed Neumann problems, *J. Diff. Eqns.* 158(1999), 1-27.
- [19] C. Gui and J. Wei, On multiple mixed interior and boundary peak solutions for some singularly perturbed Neumann problems, *Can. J. Math.* 52(2000), 522-538.
- [20] C. Gui, J. Wei and M. Winter, Multiple boundary peak solutions for some singularly perturbed Neumann problems, *Ann. Inst. H. Poincaré Anal. Non Linéaire* 17 (2000), 249-289.
- [21] D. Henry, *Geometric Theory of Semilinear Parabolic Equations*, Springer, Berlin, 1981.
- [22] M. A. Herrero, J.J.L. Velazquez, Singularity patterns in a chemotaxis model, *Math. Ann.* 306 (1996), 583–623.
- [23] M. A. Herrero, J.J.L. Velazquez, Chemotactic collapse for the Keller-Segel model, *J. Math. Biol.* 35 (1996), 177–196.
- [24] M. A. Herrero, J.J.L. Velazquez, A blow-up mechanism for a chemotaxis model, *Ann. Scuola Norm. Sup. Pisa IV* 35 (1997), 633–683.
- [25] D. Horstmann, On the existence of radially symmetric blow-up solutions for the Keller–Segel model, *J. Math. Biol.* 44 (5) (2002) 463–478.

- [26] W. Jager and S. Luckhaus, On explosions of solutions to a system of partial differential equations modelling chemotaxis, *Trans. Amer. Math. Soc.* **329** (1992), 819–824
- [27] E. F. Keller and L. A. Segel, Initiation of slime mold aggregation viewed as an instability. *J. Theor. Biol.* **26** (1970), 399–415.
- [28] Y. Li, I. Shafrir, Blow-up analysis for solutions of $-\Delta u = Ve^u$ in dimension two. *Indiana Univ. Math. J.* **43** (1994), no. 4, 1255–1270.
- [29] K. Nagasaki, Y. Suzuki, Asymptotic analysis for two-dimensional elliptic eigenvalue problems with exponentially dominated nonlinearities, *Asymptotic Anal.* **3** (1990), 173–188.
- [30] T. Nagai, Blow-up of radially symmetric solutions to a chemotaxis system. *Adv. Math. Sci. Appl.* **5** (1995), no. 2, 581–601
- [31] T. Nagai, Global existence and blowup of solutions to a chemotaxis system. Proceedings of the Third World Congress of Nonlinear Analysts, Part 2 (Catania, 2000). *Nonlinear Anal.* **47** (2001), no. 2, 777–787.
- [32] V. Nanjudiah, Chemotaxis, signal relaying and aggregation morphology, *J. Theor. Biol.*, **42** (1973), pp. 63–105.
- [33] T. Nagai, T. Senba and K. Yoshida, Application of the Trudinger-Moser inequality to a parabolic system of chemotaxis, *Funckcj. Ekvac.* **40** (1997), 411–433.
- [34] T. Nagai, T. Senba, T. Suzuki, Chemotactic collapse in a parabolic system of mathematical biology, *Hiroshima Math. J.* **30** (3) (2000) 463–497.
- [35] O. Rey, J. Wei, Blow-up solutions for an elliptic Neumann problem with sub-or-supercritical nonlinearity, I: $N = 3$, *J. Funct. Anal.* **212**(2004), no.2, 472–499.
- [36] R. Schaaf, Stationary solutions of chemotaxis systems, *Trans. Amer. Math. Soc.* **292**(1985), 531–556.
- [37] T. Senba and T. Suzuki, Some structures of the solution set for a stationary system of chemotaxis. *Adv. Math. Sci. Appl.* **10**(2000), no. 1, 191–224.
- [38] T. Senba, T. Suzuki, Time global solutions to a parabolic-elliptic system modelling chemotaxis. *Asymptot. Anal.* **32** (2002), no. 1, 63–89.
- [39] R. Temam, *Infinite-Dimensional Dynamical Systems in Mechanics and Physics* (New York: Springer), 1988.
- [40] T. Senba and T. Suzuki, Weak solutions to a parabolic-elliptic system of chemotaxis, *J. Funct. Anal.* **191** (2002), no. 1, 17–51

- [41] J.J.L. Velazquez, Stability of some mechanisms of chemotactic aggregation, *SIAM J. Appl. Math.* 62 (5) (2002) 1581–1633
- [42] J.J.L. Velazquez, Point dynamics in a singular limit of the Keller-Segel model. II. Formation of the concentration regions. *SIAM J. Appl. Math.* 64 (2004), no. 4, 1224–1248.
- [43] J.J.L. Velazquez, Well-posedness of a model of point dynamics for a limit of the Keller-Segel system. *J. Differential Equations* 206 (2004), no. 2, 315–352.
- [44] G. Wang and J. Wei, Steady state solutions of a reaction-diffusion system modeling Chemotaxis. *Math. Nach.* 233-234 (2002), 221-236.