

POINT RUPTURES FOR A MEMS EQUATION WITH FRINGING FIELD

JUAN DÁVILA AND JUNCHENG WEI

ABSTRACT. We construct solutions of the equation

$$-\Delta u = \frac{\lambda(1 + |\nabla u|^2)}{(1 - u)^2}, \quad 0 < u < 1$$

in a bounded smooth domain of \mathbb{R}^2 with Dirichlet boundary condition, for $\lambda > 0$ small. These solutions approach 1 as $\lambda \rightarrow 0$ at one point, and if Ω is not simply connected we find solutions forming singularities at many points. The equation arises in the modeling of a MEMS with fringing field. A surprising connection with plasma problem is found.

1. INTRODUCTION

The following elliptic equation arises in the modeling of electrostatic Micro-Electromechanical Systems (MEMS),

$$(1.1) \quad \begin{cases} -\Delta u = \frac{\lambda}{(1 - u)^2}, & 0 < u < 1 & \text{in } \Omega \\ u = 0 & & \text{on } \partial\Omega \end{cases}$$

where Ω is a bounded domain in \mathbb{R}^2 with smooth boundary and $\lambda > 0$. Taking into account a fringing field in the modeling of the MEMS yields an extra term:

$$(1.2) \quad \begin{cases} -\Delta u = \lambda \frac{1 + \delta |\nabla u|^2}{(1 - u)^2}, & 0 < u < 1 & \text{in } \Omega \\ u = 0 & & \text{on } \partial\Omega, \end{cases}$$

where $\delta > 0$, see [13, 14, 15].

Of special interest are solutions that give rise to singularities in the equation, that is such that $u \approx 1$ in some region, which in the physical model represents a rupture in the device. We will say that a family of classical solutions u_λ of (1.1) or (1.2) develops ruptures as λ approaches a critical parameter if $\sup_\Omega u_\lambda \rightarrow 1$.

As observed by Ye and Wei [17], there is a striking difference between (1.2) with $\delta > 0$ and (1.1). On one hand, for (1.1) in the unit ball, one knows that there is a family of solutions u_λ developing a rupture at the origin for $\lambda \rightarrow \lambda_0 \neq 0$. In this case λ_0 and the limit function u_0 are explicit, see [12]. More properties on equation (1.1) can be found in [6, 7, 9, 10, 11]. On the other hand, for (1.2) with $\delta > 0$, if λ has a fixed positive lower bound, then there is an a-priori estimate $u \leq C < 1$ for any solution u of (1.2), see Theorem 4 in [17]. Then the implicit function theorem

Date: August 8, 2011.

2000 Mathematics Subject Classification. 35J60, 35B25, 35B40.

Key words and phrases. MEMS, fringing-field, singularity formation.

and the global bifurcation theorem of Rabinowitz [16], imply that there is a family (λ, u) of solutions of (1.2) with $\lambda \rightarrow 0$ and $\sup_{\Omega} u \rightarrow 1$.

The precise behavior of solutions developing ruptures as $\lambda \rightarrow 0$ and the set of possible ruptures is not known so far. We give an answer to this question by constructing families of solutions that develop one or more ruptures as $\lambda \rightarrow 0$, and obtain a precise asymptotic description. The analysis also reveals connections with the plasma problem and the Liouville equation. This seems to be the first result for the construction of multiple point ruptures.

For simplicity we will work with $\delta = 1$, namely we work with

$$(1.3) \quad \begin{cases} -\Delta u = \lambda \frac{1 + |\nabla u|^2}{(1-u)^2}, & 0 < u < 1 \quad \text{in } \Omega \\ u = 0 & \text{on } \partial\Omega, \end{cases}$$

but the results are valid for any $\delta > 0$. Let $m \geq 1$ be an integer. We say that a family of solutions u_λ of (1.3) defined for all $\lambda > 0$ small develops m isolated ruptures as $\lambda \rightarrow 0$, if there are points $\xi_{1,\lambda}, \dots, \xi_{m,\lambda} \in \Omega$, uniformly separated between them and from the boundary, such that for any $\delta > 0$

$$\limsup_{\lambda \rightarrow 0} \sup_{\Omega \setminus \bigcup_{j=1}^m B_\delta(\xi_{j,\lambda})} u < 1$$

and for any $\delta > 0$ and all $i = 1, \dots, m$:

$$\sup_{B_\delta(\xi_{i,\lambda})} u \rightarrow 1 \quad \text{as } \lambda \rightarrow 0.$$

Our main results are the following.

Theorem 1.1. *There exists $\lambda_0 > 0$ such that for $\lambda \in (0, \lambda_0)$ there is a solution u_λ of (1.3) developing one isolated rupture as $\lambda \rightarrow 0$.*

Theorem 1.2. *If Ω is not simply connected, for any integer $m \geq 1$ there exists $\lambda_m > 0$ such that for $\lambda \in (0, \lambda_m)$ there is a solution u_λ of (1.3) developing m isolated ruptures as $\lambda \rightarrow 0$.*

The location of the ruptures is determined by the Green function and its regular part. Let G denote the Green function for the Laplacian with Dirichlet boundary condition:

$$\begin{cases} -\Delta G(\cdot, y) = \delta_y & \text{in } \Omega \\ G(x, y) = 0 & \text{for all } x \in \partial\Omega \end{cases}$$

and H its regular part, given by

$$(1.4) \quad H(x, y) = G(x, y) - \frac{1}{2\pi} \log\left(\frac{1}{|x-y|}\right).$$

For an integer $m \geq 1$ and ξ_1, \dots, ξ_m different points in Ω we define

$$(1.5) \quad \varphi_m(\xi_1, \dots, \xi_m) = \sum_{i=1}^m H(\xi_i, \xi_i) + \sum_{j \neq i} G(\xi_i, \xi_j).$$

Then in Theorems 1.1 and 1.2, after passing to a subsequence, the rupture points $\xi_{1,\lambda}, \dots, \xi_{m,\lambda} \in \Omega$ of the solution u_λ , converge to a critical point of φ_m .

Also we obtain the following result.

Theorem 1.3. *For any non-degenerate critical point $\xi = (\xi_1, \dots, \xi_m)$ of φ_m there is $\lambda_0 > 0$ such that $\lambda \in (0, \lambda_0)$ there is a solution u_λ of (1.3) developing m isolated ruptures as $\lambda \rightarrow 0$ at points $\xi_{1,\lambda}, \dots, \xi_{m,\lambda} \in \Omega$ that converge to (ξ_1, \dots, ξ_m) .*

Regarding the asymptotic behavior of the solutions we construct, if u_λ is the solution of any of the three theorems above developing m ruptures $\xi_{1,\lambda}, \dots, \xi_{m,\lambda} \in \Omega$, then

$$u(x) = \frac{2\pi}{|\log \lambda|} \sum_{j=1}^m G(x, \xi_{j,\lambda})(1 + o(1))$$

as $\lambda \rightarrow 0$, for points $x \in \Omega$ away from $\xi_{1,\lambda}, \dots, \xi_{m,\lambda}$. Very close to the points $\xi_{j,\lambda}$ we have an expansion of the form

(1.6)

$$u(x) = 1 - \frac{\lambda}{|\log \lambda|} - \frac{\lambda \log |\log \lambda|}{|\log \lambda|^2} + \frac{\lambda}{|\log \lambda|^2} \log \left(\alpha_0 V_0 \left(\frac{|\log \lambda|^2}{\lambda} |x - \xi_{j,\lambda}| \right) \right) + \text{l.o.t.}$$

where l.o.t. contains smaller order terms and the function V_0 is the unique radial function satisfying

$$-\Delta V_0 = (V_0)_+ \quad \text{in } \mathbb{R}^2, \quad \max_{\mathbb{R}^2} V_0 = 1.$$

The number α_0 is given by $\alpha_0 = \left(\frac{1}{2\pi} \int_{\mathbb{R}^2} (V_0)_+ \right)^{-1}$. The expansion (1.6) is valid for points $x \in \Omega$ such that $|x - \xi_{j,\lambda}| \leq (R_0 - \delta) \frac{\lambda}{|\log \lambda|^2}$, where $R_0 > 0$ is the radius for which $V_0(R_0) = 0$ and $0 < \delta < R_0$.

The key behind these results is a change of variables introduced in Ye and Wei [17], which allows us to rewrite problem (1.3) in the form

$$(1.7) \quad \begin{cases} -\frac{\lambda^2}{|\log \lambda|^4} \Delta v = p_\lambda(v-1) & \text{in } \Omega \\ v = 0 & \text{on } \partial\Omega, \end{cases}$$

with a nonlinearity p_λ that satisfies

$$p_\lambda(x) \rightarrow x_+ \quad \text{as } \lambda \rightarrow 0$$

for all $x \in \mathbb{R}$, where $x_+ = \max(x, 0)$. See the derivation in Section 2 and the definition of p_λ in (2.13). For $\lambda > 0$ we may expect solutions of (1.3) to be related to solutions of

$$\begin{cases} -\frac{\lambda^2}{|\log \lambda|^4} \Delta v = (v-1)_+ & \text{in } \Omega \\ v = 0 & \text{on } \partial\Omega. \end{cases}$$

This problem is, after some transformations, the same as the *plasma* problem studied in [3, 2] and also presents similarities with the Liouville equation [1, 4, 5, 8]. For both problems there are existence and classification results of solutions exhibiting point concentration. In both problems the location of the concentration points is determined by an energy expansion that at main nontrivial order involves the Green function and its regular part.

The proof of all theorems is carried out with equation (1.7), or more precisely, with a rescaled version of it, equation (3.1) below. We construct an approximate solution for this equation, in the same spirit as in [2, 4, 5], which is precise enough to perform a finite dimensional reduction using a Lyapunov-Schmidt procedure. The

main difficulty in this process is that the nonlinearity p_λ converges to x_+ , which is only Lipschitz and hence, some estimates of derivatives of the solution, which involve p'_λ , become delicate as $\lambda \rightarrow 0$. This is similar to the difficulty in [2], except that for us the nonlinearity is not explicit but smooth. After the reduction, the problem becomes one of finding critical points of a function that is close in C^1 norm to φ_m on compact subsets of its domain of definition. In the case of two or more points of concentration in a non-simply connected domain, this is guaranteed by a result in [5], which has also appeared in [2].

2. CHANGE OF VARIABLES

Equation (1.3) is equivalent to

$$\begin{cases} \Delta u = \lambda \frac{1 + |\nabla u|^2}{u^2}, & 0 < u < 1 & \text{in } \Omega \\ u = 1 & & \text{on } \partial\Omega. \end{cases}$$

Let us write u in the form $u(x) = \lambda w(x/\lambda)$, $x \in \Omega$, so that w satisfies

$$\begin{cases} \Delta w = \frac{1 + |\nabla w|^2}{w^2}, & 0 < w < \frac{1}{\lambda} & \text{in } \Omega/\lambda \\ w = \frac{1}{\lambda} & & \text{on } \partial\Omega/\lambda. \end{cases}$$

Motivated by [17] we introduce the change of variables

$$v = g(w) \quad \text{where} \quad g(w) = \int_w^1 e^{1/s} ds,$$

and we compute

$$\Delta v = -\frac{1}{w^2} e^{1/w}.$$

Note that $g : (0, +\infty) \rightarrow \mathbb{R}$ is a decreasing convex function with range equal to \mathbb{R} . Therefore $g^{-1} : \mathbb{R} \rightarrow (0, +\infty)$ is well defined. Let

$$h(w) = \frac{1}{w^2} e^{1/w} \quad \text{for all } w > 0$$

and

$$(2.1) \quad f(v) = h(g^{-1}(v)).$$

Then v satisfies

$$(2.2) \quad \begin{cases} -\Delta v = f(v) & \text{in } \Omega/\lambda \\ v = -c_\lambda & \text{on } \partial\Omega/\lambda \end{cases}$$

where

$$(2.3) \quad c_\lambda = -g(1/\lambda) = \int_1^{1/\lambda} e^{1/s} ds.$$

Note that

$$c_\lambda = \frac{1}{\lambda} + |\log \lambda| + O(1) \quad \text{as } \lambda \rightarrow 0$$

and that $f : \mathbb{R} \rightarrow (0, \infty)$ is a convex increasing function. Moreover

$$(2.4) \quad f(v) = v(\log v)^4 + o(v(\log v)^4) \quad \text{as } v \rightarrow +\infty$$

$$(2.5) \quad f(v) = O\left(\frac{1}{v^2}\right) \quad \text{as } v \rightarrow -\infty.$$

We perform one more change of variables

$$v(x) = M\tilde{v}((\log M)^2 x) \quad x \in \frac{1}{\lambda}\Omega,$$

where $M > 0$ is a new large parameter to be chosen later on. We find

$$-\Delta\tilde{v} = \frac{1}{M(\log M)^2} f(M\tilde{v}) \quad \text{in } \frac{(\log M)^2}{\lambda}\Omega.$$

By (2.4), (2.5)

$$\frac{1}{M(\log M)^2} f(M\tilde{v}) \rightarrow \tilde{v}_+ \quad \text{as } M \rightarrow +\infty$$

for any $\tilde{v} \in \mathbb{R}$, where

$$x_+ = \max(x, 0) \quad \forall x \in \mathbb{R}.$$

If we choose M such that $\max\tilde{v} = 1$, then \tilde{v} solves at main order

$$\begin{aligned} -\Delta\tilde{v} &= \tilde{v}_+ \quad \text{in } \frac{(\log M)^2}{\lambda}\Omega \\ \max\tilde{v} &= 1. \end{aligned}$$

Let $R_0 > 0$ be the unique number such that the principal Dirichlet eigenvalue of $-\Delta$ in $B_{R_0}(0)$ is 1, that is, the problem

$$\begin{cases} -\Delta\varphi_1 = \varphi_1, & \varphi_1 > 0 \quad \text{in } B_{R_0}(0) \\ \varphi_1 = 0 & \text{on } \partial B_{R_0}(0) \end{cases}$$

has a solution. We normalize the principal eigenfunction so that $\varphi_1(0) = 1$ and let

$$(2.6) \quad d_0 = -\varphi_1'(R_0).$$

Define

$$(2.7) \quad V_0(x) = \begin{cases} \varphi_1(x) & \text{if } x \in B_{R_0}(0) \\ -d_0 R_0 \log(|x|/R_0) & \text{if } x \notin B_{R_0}(0). \end{cases}$$

Then V_0 is a radially symmetric function, with $\max_{\mathbb{R}^2} V_0 = 1$ and V_0 satisfies

$$-\Delta V_0 = (V_0)_+ \quad \text{in } \mathbb{R}^2.$$

Heuristically we can expect that

$$\tilde{v}(x) \approx V_0(x)$$

on compact sets, and this means

$$v(x) \approx M V_0((\log M)^2 x).$$

Evaluating this relation on the boundary of $\frac{1}{\lambda}\Omega$ we find

$$c_\lambda = M d_0 R_0 \log\left(\frac{(\log M)^2 D}{\lambda R_0}\right)$$

where D represents the diameter of the domain. Note that this gives the relation

$$(2.8) \quad M_\lambda = \frac{1}{d_0 R_0 \lambda |\log \lambda|} + o\left(\frac{1}{\lambda |\log \lambda|}\right) \quad \text{as } \lambda \rightarrow 0.$$

Since $\log(M_\lambda) = |\log \lambda| + o(|\log \lambda|)$ as $\lambda \rightarrow 0$ it is natural to choose from the beginning $M = |\log \lambda|$. Thus, we start from (2.2) and rewrite it as

$$\begin{cases} -\Delta \tilde{v} = f(\tilde{v} - c_\lambda) & \text{in } \Omega/\lambda \\ \tilde{v} = 0 & \text{on } \partial\Omega/\lambda \end{cases}$$

where c_λ is defined in (2.3). Motivated by (2.8), we write \tilde{v} as

$$(2.9) \quad \tilde{v}(x) = \frac{c_\lambda}{|\log \lambda|} v(|\log \lambda|^2 x)$$

which leads to

$$(2.10) \quad \begin{cases} -\Delta v = f_\lambda(v - |\log \lambda|) & \text{in } \Omega_\lambda = \frac{|\log \lambda|^2}{\lambda} \Omega \\ v = 0 & \text{on } \partial\Omega_\lambda, \end{cases}$$

where

$$(2.11) \quad f_\lambda(t) = \frac{1}{c_\lambda |\log \lambda|^3} f\left(\frac{c_\lambda}{|\log \lambda|} t\right).$$

Then, thanks to (2.4), (2.5), $f_\lambda(x) \rightarrow x_+$ as $\lambda \rightarrow 0$.

Remark 2.1. *If instead of the change of variables (2.9) we choose*

$$\tilde{v}(x) = c_\lambda v(|\log \lambda|^2 x)$$

then we obtain

$$(2.12) \quad \begin{cases} -\Delta v = p_\lambda(v - 1) & \text{in } \Omega_\lambda \\ v = 0 & \text{on } \partial\Omega_\lambda \end{cases}$$

where

$$(2.13) \quad p_\lambda(t) = \frac{1}{c_\lambda |\log \lambda|^4} f(c_\lambda t)$$

and f is given by (2.1). For this nonlinearity we also have $p_\lambda(x) \rightarrow x_+$ as $\lambda \rightarrow 0$. Here we see the similarity of (2.12) with the plasma problem studied in [2].

Finally, we collect here some useful estimates for the non-linearity (we omit the computations).

Lemma 2.2. *Let f be defined by (2.1) and f_λ be the function (2.11). Then the following properties hold:*

$$f(v) = v((\log v)^4 + O((\log v)^3 \log \log v)) \quad \text{as } v \rightarrow +\infty$$

$$f(v) \leq \frac{C}{1+v^2} \quad \text{for all } x \leq 0$$

$$(2.14) \quad f_\lambda(x) = x_+ + O\left(\frac{\log |\log \lambda|}{|\log \lambda|}\right) \quad \text{as } \lambda \rightarrow 0, \text{ uniformly for } x \text{ on bounded sets}$$

$$(2.15) \quad f_\lambda(t) \leq C \frac{\lambda^3}{|\log \lambda|(t^2 + \lambda^2 |\log \lambda|)} \quad \text{for } t \leq 0$$

$$(2.16) \quad f'_\lambda(t) = O(1) \quad \text{uniformly for } t \text{ in bounded sets}$$

$$(2.17) \quad f'_\lambda(t) \leq \frac{C\lambda^3}{|\log \lambda|(|t|^3 + \lambda^3 |\log \lambda|^3)} \quad \text{for all } t \leq 0$$

$$(2.18) \quad f''_\lambda(t) = O(1/\lambda) \quad \text{uniformly for } t \text{ in bounded sets}$$

$$(2.19) \quad f''_\lambda(t) \leq \frac{C}{t} \quad \text{if } t/(\lambda |\log \lambda|) \rightarrow \infty$$

$$(2.20) \quad f''_\lambda(t) \leq C \frac{\lambda^3}{|\log \lambda|(|t|^4 + \lambda^4 |\log \lambda|^4)} \quad \text{for all } t \leq 0$$

3. FIRST APPROXIMATION AND ITS ENERGY

We will work mainly with the following reformulation of problem (1.3),

$$(3.1) \quad \begin{cases} -\Delta v = f_\lambda(v - |\log \lambda|) & \text{in } \Omega_\lambda \\ v = 0 & \text{on } \partial\Omega_\lambda \end{cases}$$

where

$$f_\lambda(x) = \frac{1}{c_\lambda |\log \lambda|^3} f\left(\frac{c_\lambda}{|\log \lambda|} x\right)$$

with f defined in (2.1) and

$$\Omega_\lambda = \frac{|\log \lambda|^2}{\lambda} \Omega.$$

We will define an initial approximation of a solution to (3.1) based on the solutions $w_{\lambda,\alpha}$ of the following problem

$$(3.2) \quad \begin{cases} w''_{\lambda,\alpha}(r) + \frac{1}{r} w'_{\lambda,\alpha}(r) = -f_\lambda(w_{\lambda,\alpha}) & r > 0 \\ w_{\lambda,\alpha}(0) = \alpha, \quad w'_{\lambda,\alpha}(0) = 0, \end{cases}$$

where $\alpha > 0$.

Lemma 3.1. *We have:*

a)

$$(3.3) \quad w_{\lambda,\alpha} = \alpha V_0 + O\left(\frac{\log |\log \lambda|}{|\log \lambda|}\right)$$

as $\lambda \rightarrow 0$ in the C^1 norm over compact sets of \mathbb{R}^2 , where V_0 is the function defined in (2.7).

b) *There is a unique $R_{\lambda,\alpha} > 0$ such that*

$$w_{\lambda,\alpha}(R_{\lambda,\alpha}) = 0,$$

and it satisfies

$$(3.4) \quad R_{\lambda,\alpha} = R_0 + O\left(\frac{\log|\log\lambda|}{|\log\lambda|}\right), \quad \frac{d}{d\alpha}R_{\lambda,\alpha} = O\left(\frac{\log|\log\lambda|}{|\log\lambda|}\right)$$

as $\lambda \rightarrow 0$ where R_0 is the number defined just before (2.7) and $O\left(\frac{\log|\log\lambda|}{|\log\lambda|}\right)$ is uniform for α in compact sets of $(0, +\infty)$.

c)

$$(3.5) \quad -w'_{\lambda,\alpha}(R_{\lambda,\alpha}) \rightarrow d_0\alpha \quad \text{as } \lambda \rightarrow 0$$

uniformly for α in compact sets of $(0, +\infty)$, where d_0 is defined in (2.6).

For the proof see the end of this section.

We proceed now with the construction of an initial approximation of a solution to (3.1). Let $m \geq 1$ be a fixed integer, $\xi_1, \dots, \xi_m \in \Omega$, and $\mu_1, \dots, \mu_m > 0$. We will always assume that for some small $\delta > 0$

$$(3.6) \quad |\xi_i - \xi_j| \geq \delta \quad \text{for all } i \neq j$$

$$(3.7) \quad \text{dist}(\xi_j, \partial\Omega) \geq \delta, \quad \text{for all } 1 \leq j \leq m,$$

$$(3.8) \quad \delta \leq \mu_j \leq \delta^{-1} \quad \text{for all } 1 \leq j \leq m.$$

Let us use the notation

$$\bar{\xi}_j = \frac{|\log\lambda|^2}{\lambda} \xi_j \in \Omega_\lambda.$$

The parameters μ_j will be chosen later on.

Thanks to (3.5), we can find for $\lambda > 0$ small a unique positive number α_j such that

$$(3.9) \quad -w'_{\lambda,\alpha_j}(R_{\lambda,\alpha_j})R_{\lambda,\alpha_j} = \mu_j,$$

Let us write

$$(3.10) \quad w_j(r) = w_{\lambda,\alpha_j}(r), \quad R_j = R_{\lambda,\alpha_j}.$$

We define

$$V_j(x) = \begin{cases} w_j(|x - \bar{\xi}_j|) + |\log\lambda| & \text{if } |x - \bar{\xi}_j| \leq R_j \\ -\mu_j \log\left(\frac{|x - \bar{\xi}_j|}{R_j}\right) + |\log\lambda| & \text{if } |x - \bar{\xi}_j| > R_j. \end{cases}$$

The function V_j is radial about the point $\bar{\xi}_j$ and it is C^1 across the boundary of the ball $B_{R_j}(\bar{\xi}_j)$, thanks to (3.9). It satisfies

$$(3.11) \quad \begin{cases} -\Delta V_j = f_\lambda(V_j - |\log\lambda|) \chi_{B_{R_j}(\bar{\xi}_j)} & \text{on } \mathbb{R}^2, \\ \max_{\mathbb{R}^2} V_j = V_j(\bar{\xi}_j) = \alpha_j + |\log\lambda| \end{cases}$$

Since the function V_j does not satisfy the boundary condition on $\partial\Omega_\lambda$ we consider $V_j - H_j$ where H_j is the solution of the problem

$$(3.12) \quad \begin{cases} \Delta H_j = 0 & \text{in } \Omega_\lambda \\ H_j = V_j & \text{on } \partial\Omega_\lambda. \end{cases}$$

Then

$$H_j(x) = -\mu_j \log\left(\frac{|x - \bar{\xi}_j|}{R_j}\right) + |\log\lambda| \quad \text{for all } x \in \partial\Omega_\lambda,$$

and we get the formula

$$H_j(x) = -2\pi\mu_j H\left(\frac{\lambda}{|\log \lambda|^2}x, \xi_j\right) - 2\mu_j \log |\log \lambda| + \mu_j \log R_j + (1 - \mu_j)|\log \lambda|$$

where H is the regular part of the Green function, c.f. (1.4). We define the initial approximation as:

$$(3.13) \quad V_\lambda = \sum_{j=1}^m V_j - H_j.$$

We look for a solution v of (3.1) of the form $v = V_\lambda + \phi$ where ϕ is small compared to V_λ . Then problem (3.1) gets reformulated in terms of ϕ as follows:

$$(3.14) \quad \begin{cases} \Delta\phi + f'_\lambda(V_\lambda - |\log \lambda|)\phi + E_\lambda + N(\phi) = 0 & \text{in } \Omega_\lambda \\ \phi = 0 & \text{on } \partial\Omega_\lambda, \end{cases}$$

where

$$\begin{aligned} E_\lambda &= \Delta V_\lambda + f_\lambda(V_\lambda - |\log \lambda|) \\ N(\phi) &= f_\lambda(V_\lambda + \phi - |\log \lambda|) - f_\lambda(V_\lambda - |\log \lambda|) - f'_\lambda(V_\lambda - |\log \lambda|)\phi. \end{aligned}$$

Up to now the parameters μ_j were free in the interval (δ, δ^{-1}) ($\delta > 0$ a small constant). To ensure that E_λ is small in an appropriate norm to be introduced later it is necessary to adjust the numbers μ_j in a suitable way.

Lemma 3.2. *Assume that $\mu_1, \dots, \mu_m > 0$ satisfy the system of equations*

$$(3.15) \quad \mu_i = 1 - 2\mu_i \frac{\log |\log \lambda|}{|\log \lambda|} + \mu_i \frac{\log R_i}{|\log \lambda|} - \frac{2\pi}{|\log \lambda|} \left[\mu_i H(\xi_i, \xi_i) + \sum_{i \neq j} \mu_j G(\xi_i, \xi_j) \right]$$

for all $i = 1, \dots, m$. Then for all $R > 0$, we have

$$(3.16) \quad V_\lambda(x) = w_i(|x - \bar{\xi}_i|) + |\log \lambda| + O\left(\frac{\lambda}{|\log \lambda|^2}\right), \quad \text{for all } x \in B_R(\bar{\xi}_i)$$

as $\lambda \rightarrow 0$, where the term $O\left(\frac{\lambda}{|\log \lambda|^2}\right)$ is in C^1 norm in $B_R(\bar{\xi}_i)$

We note that the system of equations (3.15) is nonlinear since the functions R_i depend on μ_i . Nevertheless this system is solvable for small $\lambda > 0$ and we obtain the following expansion for the solution

$$(3.17) \quad \mu_i = 1 - 2\frac{\log |\log \lambda|}{|\log \lambda|} + \frac{\log(R_0) - 2\pi}{|\log \lambda|} \left[H(\xi_i, \xi_i) + \sum_{j \neq i} G(\xi_i, \xi_j) \right] + O\left(\frac{(\log |\log \lambda|)^2}{|\log \lambda|^2}\right)$$

as $\lambda \rightarrow 0$.

Because of Lemma 3.2 we will work in the sequel only with μ_i satisfying (3.15) and in particular we will always assume that

$$(3.18) \quad \mu_i = 1 + O\left(\frac{\log |\log \lambda|}{\log \lambda}\right) \quad \text{as } \lambda \rightarrow 0.$$

Also, thanks to Lemma 3.1, we will assume

$$R_i = R_0 + O\left(\frac{\log |\log \lambda|}{\log \lambda}\right) \quad \text{as } \lambda \rightarrow 0.$$

To solve (3.14) with ϕ small in a convenient sense we need to choose the points $\xi_1, \dots, \xi_m \in \Omega_\lambda$ appropriately, and for this we use a variational formulation of (3.1): v is a solution of (3.1) if and only if $v \in H_0^1(\Omega)$ is a critical point of the energy functional

$$(3.19) \quad I_\lambda(v) = \frac{1}{2} \int_{\Omega_\lambda} |\nabla v|^2 - \int_{\Omega_\lambda} F_\lambda(v - |\log \lambda|)$$

where

$$F_\lambda(v) = \int_0^v f_\lambda(s) ds.$$

We note that I_λ is a C^1 functional on $H_0^1(\Omega)$. It ϕ is a small solution of (3.14) one may expect that $I_\lambda(V_\lambda + \phi)$ is critical with respect to ξ_1, \dots, ξ_m . Therefore, in order to find the good choice of the points ξ_1, \dots, ξ_m it becomes important to compute $I_\lambda(V_\lambda)$.

Lemma 3.3. *Suppose $\xi_1, \dots, \xi_m \in \Omega$ satisfy the separation conditions (3.6), (3.7), and let μ_j satisfy (3.15). Then*

$$(3.20) \quad I_\lambda(V_\lambda) = a_1 |\log \lambda| - 2a_1 \log |\log \lambda| - a_2 \varphi_m(\xi_1, \dots, \xi_m) + o(1)$$

as $\lambda \rightarrow 0$, where φ_m is defined in (1.5),

$$a_1 = \pi m R_0, \quad a_2 = \pi R_0 (2\pi - \log R_0),$$

and $o(1)$ is uniform with respect to the C^1 norm in the region (3.6), (3.7).

Proof. We compute

$$I_\lambda(V_\lambda) = \frac{1}{2} \int_{\Omega_\lambda} |\nabla V_\lambda|^2 - \int_{\Omega_\lambda} F_\lambda(V_\lambda - |\log \lambda|).$$

Using (3.11) and (3.12) we have

$$\int_{\Omega_\lambda} |\nabla V_\lambda|^2 = \int_{\Omega_\lambda} \left| \sum_{j=1}^m \nabla(V_j - H_j) \right|^2 = \sum_{j=1}^m \int_{B_{R_j}(\bar{\xi}_j)} f_\lambda(w_j(|x - \bar{\xi}_j|)) V_\lambda.$$

Therefore By Lemma 3.2

$$(3.21) \quad \begin{aligned} & \int_{\Omega_\lambda} |\nabla V_\lambda|^2 \\ &= \sum_{j=1}^m \int_{B_{R_j}(\bar{\xi}_j)} f_\lambda(w_j(|x - \bar{\xi}_j|)) \left(w_j(|x - \bar{\xi}_j|) + |\log \lambda| + O\left(\frac{\lambda}{|\log \lambda|^2}\right) \right) dx \\ &= |\log \lambda| \sum_{j=1}^m \int_{B_{R_j}(0)} f_\lambda(w_j(x)) dx + \sum_{j=1}^m \int_{B_{R_j}(0)} f_\lambda(w_j(x)) w_j(x) dx + O\left(\frac{\lambda}{|\log \lambda|^2}\right) \end{aligned}$$

as $\lambda \rightarrow 0$. Integrating (3.11) in $B_{R_j}(\bar{\xi}_j)$ and then using (3.4) and (3.17) we obtain

$$(3.22) \quad \int_{B_{R_j}(0)} f_\lambda(w_j(x)) dx = 2\pi R_j \mu_j = 2\pi R_0 - 4\pi R_0 \frac{\log |\log \lambda|}{|\log \lambda|} \\ + 2\pi R_0 \frac{\log R_0 - 2\pi}{|\log \lambda|} \left[H(\xi_j, \xi_j) + \sum_{i \neq j} G(\xi_j, \xi_i) \right] \\ + O\left(\frac{(\log |\log \lambda|)^2}{|\log \lambda|^2}\right).$$

For the terms $\int_{B_{R_j}(0)} f_\lambda(w_j) w_j$, we note that multiplying (3.11) by w_j and integrating in B_{R_j} we obtain

$$\int_{B_{R_j}(0)} f_\lambda(w_j) w_j = \int_{B_{R_j}(0)} |\nabla w_j|^2.$$

But by (3.5)

$$w_j = \alpha_j V_0 + O\left(\frac{\log |\log \lambda|}{|\log \lambda|}\right)$$

in the C^1 norm over compact sets of \mathbb{R}^2 , where α_j satisfies (3.9), which gives

$$\mu_j = d_0 R_0 \alpha_j + O\left(\frac{\log |\log \lambda|}{|\log \lambda|}\right).$$

This gives

$$|\nabla w_j|^2 = \frac{1}{d_0^2 R_0^2} |\nabla V_0|^2 + O\left(\frac{\log |\log \lambda|}{|\log \lambda|}\right).$$

Using this, (3.4) and (3.18) we see that

$$\int_{B_{R_j}(0)} f_\lambda(w_j) w_j = \frac{1}{d_0^2 R_0^2} \int_{B_{R_0}(0)} |\nabla V_0|^2 + O\left(\frac{\log |\log \lambda|}{|\log \lambda|}\right).$$

Therefore we find

$$(3.23) \quad \frac{1}{2} \int_{\Omega_\lambda} |\nabla V_\lambda|^2 = \pi m R_0 |\log \lambda| - \pi m R_0 \log |\log \lambda| + \pi R_0 (\log R_0 - 2\pi) \varphi_m(\xi_1, \dots, \xi_m) \\ + \frac{m}{2d_0^2 R_0^2} \int_{B_{R_0}} |\nabla V_0|^2 + O\left(\frac{(\log |\log \lambda|)^2}{|\log \lambda|}\right).$$

Now we compute

$$\int_{\Omega_\lambda} F_\lambda(V_\lambda - |\log \lambda|) = \sum_{j=1}^m \int_{B_{R_j}(\bar{\xi}_j)} F_\lambda(V_\lambda - |\log \lambda|) + \int_{\tilde{\Omega}_\lambda} F_\lambda(V_\lambda - |\log \lambda|)$$

where $\tilde{\Omega}_\lambda = \Omega \setminus \cup_{j=1}^m B_{R_j}(\bar{\xi}_j)$. First we have, using (3.2)

$$(3.24) \quad \int_{B_{R_j}(\bar{\xi}_j)} F_\lambda(V_\lambda - |\log \lambda|) = \int_{B_{R_j}(0)} F_\lambda(w_j) + O\left(\frac{\lambda}{|\log \lambda|^2}\right).$$

Using Lemma 3.2, $F_\lambda(t) = \frac{1}{2} t_+^2 + O\left(\frac{\log |\log \lambda|}{|\log \lambda|}\right)$, (3.4) and (3.18), we obtain

$$(3.25) \quad \int_{B_{R_j}(\bar{\xi}_j)} F_\lambda(V_\lambda - |\log \lambda|) = \frac{1}{2d_0^2 R_0^2} \int_{B_{R_0}} V_0^2 + O\left(\frac{\log |\log \lambda|}{|\log \lambda|}\right)$$

as $\lambda \rightarrow 0$. To estimate the integral in $\tilde{\Omega}_\lambda$ we use the inequality (2.15) which implies that for any fixed $M > 0$

$$(3.26) \quad |F_\lambda(x)| \leq C \frac{\lambda}{|\log \lambda|^2} \quad \text{for all } -M \leq x \leq 0$$

and

$$(3.27) \quad |F_\lambda(x)| \leq C \frac{\lambda^3}{|\log \lambda|} \quad \text{for all } x \leq -M.$$

We write

$$\begin{aligned} \int_{\Omega_\lambda} F_\lambda(V_\lambda - |\log \lambda|) &= \sum_{j=1}^m \int_{B_{10R_j}(\bar{\xi}_j) \setminus B_{R_j}(\bar{\xi}_j)} F_\lambda(V_\lambda - |\log \lambda|) \\ &\quad + \int_{\Omega_\lambda \setminus \cup B_{10R_j}(\bar{\xi}_j)} F_\lambda(V_\lambda - |\log \lambda|) \end{aligned}$$

and estimate, using Lemma 3.2 and (3.26),

$$\begin{aligned} \int_{B_{10R_j}(\bar{\xi}_j) \setminus B_{R_j}(\bar{\xi}_j)} F_\lambda(V_\lambda - |\log \lambda|) &= \int_{B_{10R_j}(\bar{\xi}_j) \setminus B_{R_j}(\bar{\xi}_j)} F_\lambda(w_j + O(\frac{\lambda}{|\log \lambda|^2})) \\ &= \int_{B_{10R_j}(\bar{\xi}_j) \setminus B_{R_j}(\bar{\xi}_j)} F_\lambda(w_j) + O(\frac{\lambda}{|\log \lambda|^2}) \\ (3.28) \quad &= O(\frac{\lambda}{|\log \lambda|^2}). \end{aligned}$$

Far away from the points $\bar{\xi}_j$ we argue as follows. For points x in $\partial B_{10R_j}(\bar{\xi}_j)$ we have, because of Lemma 3.2, that $V_\lambda(x) - |\log \lambda| \leq -M$ where $M > 0$ is a fixed constant. But $V_\lambda - |\log \lambda|$ is harmonic in $\Omega \setminus \cup_{j=1}^m B_{10R_j}(\bar{\xi}_j)$ and equal to $-|\log \lambda|$ on $\partial\Omega_\lambda$, and therefore $V_\lambda(x) - |\log \lambda| \leq -M$ holds for all points in $\Omega \setminus \cup_{j=1}^m B_{10R_j}(\bar{\xi}_j)$. Thus we can use (3.27) and deduce

$$\int_{\Omega_\lambda \setminus \cup B_{10R_j}(\bar{\xi}_j)} F_\lambda(V_\lambda - |\log \lambda|) = O(\lambda |\log \lambda|^3).$$

Combining the estimates for each term, and noting that

$$\int_{B_{R_0}} |\nabla V_0|^2 = \int_{B_{R_0}} V_0^2,$$

we conclude that formula (3.20) is valid with $o(1)$ in the C^0 norm over the region (3.6), (3.7).

Regarding the estimate in C^1 norm, the error term in (3.23) is also $O(\frac{(\log |\log \lambda|)^2}{|\log \lambda|})$ in C^1 norm for the parameters in the region (3.6), (3.7), since we can use (3.16) for the error in (3.21) and the dependence on the ξ_j in the terms appearing in (3.21) is through the μ_i in expression (3.22). Similarly, the errors in (3.24), (3.25) and (3.28) are also C^1 with respect to ξ_i . \square

Proof of Lemma 3.1. First we remark that for $\lambda > 0$ the nonlinearity f_λ is smooth, and that the solution $w_{\lambda, \alpha}$ of (3.2) exists for all $r \geq 0$. It is also smooth with respect to $\alpha > 0$, $\lambda > 0$.

a) Estimate (3.3) follows from $f_\lambda(t) = t_+ + O(\frac{\log |\log \lambda|}{|\log \lambda|})$ as $\lambda \rightarrow 0$ uniformly on compact subsets of \mathbb{R} (c.f. (2.14)).

b) The existence of $R_{\lambda,\alpha}$ follows from the convergence in part a), and the uniqueness because $\frac{\partial}{\partial r}w_{\lambda,\alpha}(r) < 0$. The estimate for $R_{\lambda,\alpha}$ in (3.4) follows by the implicit function theorem, noting that $w_{\lambda,\alpha}(r)$ is C^1 in r , that $w_{\lambda,\alpha}$ and $\frac{\partial}{\partial r}w_{\lambda,\alpha}(r)$ have continuous extensions to $\lambda = 0$, and $\frac{\partial}{\partial r}w_{\lambda,\alpha}(R_0) < 0$ (with a uniform distance as $\lambda \rightarrow 0$). What needs to be verified is that the expansion

$$w_{\lambda,\alpha}(r) = w_{\lambda,\alpha}(R_0) + \frac{\partial}{\partial r}w_{\lambda,\alpha}(R_0)(r - R_0) + o(|r - R_0|)$$

as $r \rightarrow R_0$, has an error $o(|r - R_0|)$ which is uniform as $\lambda \rightarrow 0$. This estimate can be obtained from elliptic estimates, since $\Delta w_{\lambda,\alpha}$ remains bounded, and so is uniformly $C^{1,\mu}$ on compact sets.

We prove now the estimate $\frac{d}{d\alpha}R_{\lambda,\alpha} = O(\frac{\log|\log\lambda|}{|\log\lambda|})$ in (3.4). Differentiating the relation $w_{\lambda,\alpha}(R_{\lambda,\alpha}) = 0$ with respect to α we obtain

$$\frac{d}{d\alpha}R_{\lambda,\alpha} = -\frac{\frac{\partial}{\partial\alpha}w_{\lambda,\alpha}(R_{\lambda,\alpha})}{\frac{\partial}{\partial r}w_{\lambda,\alpha}(R_{\lambda,\alpha})}.$$

Since $\frac{\partial}{\partial r}w_{\lambda,\alpha}(R_{\lambda,\alpha})$ is bounded away from zero, we need only to estimate $\frac{\partial}{\partial\alpha}w_{\lambda,\alpha}(R_{\lambda,\alpha})$. Let $z_{\lambda,\alpha} = \frac{\partial}{\partial\alpha}w_{\lambda,\alpha}$ which is smooth and satisfies

$$(3.29) \quad \Delta z_{\lambda,\alpha} = f'_\lambda(w_{\lambda,\alpha})z_{\lambda,\alpha} \quad \text{in } \mathbb{R}^2.$$

We claim that

$$(3.30) \quad |z_{\lambda,\alpha}(R_0)| \leq C \frac{\log|\log\lambda|}{|\log\lambda|}.$$

To prove this we let $\phi(r) = z_{\lambda,\alpha}(r) - V_0(r)$ and note that

$$\Delta\phi = f'_\lambda(w_{\lambda,\alpha})z_{\lambda,\alpha} - \chi_{B_{R_0}}V_0 = \chi_{B_{R_0}}\phi + (f'_\lambda(w_{\lambda,\alpha}) - \chi_{B_{R_0}})z_{\lambda,\alpha}.$$

Multiplying this equation by V_0 and integrating in B_{R_0} gives:

$$-2\pi R_0\phi(R_0)V'(R_0) = \int_{B_{R_0}} (f'_\lambda(w_{\lambda,\alpha}) - \chi_{B_{R_0}})z_{\lambda,\alpha}V_0$$

and the integral on the right hand side can be verified to be $O(\frac{\log|\log\lambda|}{|\log\lambda|})$ as $\lambda \rightarrow 0$. Equation (3.29) shows that $z_{\lambda,\alpha}$ is $C^{1,\mu}$ on compact set, for any $\mu \in (0, 1)$. Using this property and (3.30) we deduce

$$|z_{\lambda,\alpha}(R_{\lambda,\alpha})| \leq C \frac{\log|\log\lambda|}{|\log\lambda|}.$$

c) This property follows from the convergence $w_{\lambda,\alpha} \rightarrow \alpha V_0$ as $\lambda \rightarrow 0$ in $C^{1,\mu}$ on compact sets of \mathbb{R}^2 and $R_{\lambda,\alpha} \rightarrow R_0$ as $\lambda \rightarrow 0$. \square

Proof of Lemma 3.2. Fix $i = 1, \dots, m$. For $x \in B_{R_i}(\bar{\xi}_i)$ we have

$$\begin{aligned} V_i(x) - H_i(x) &= w_i(|x - \bar{\xi}_i|) + |\log\lambda| + 2\pi\mu_i H\left(\frac{\lambda}{|\log\lambda|^2}x, \xi_i\right) \\ &\quad + 2\mu_i \log|\log\lambda| - \mu_i \log R_i - (1 - \mu_i)|\log\lambda|. \end{aligned}$$

For $x \in B_{R_i}(\bar{\xi}_i)$, if we take $j \neq i$ then

$$V_j(x) - H_j(x) = 2\pi\mu_j G\left(\frac{\lambda}{|\log\lambda|^2}x, \xi_j\right).$$

(This is valid actually for $|x - \bar{\xi}_j| > R_j$.) Therefore, for $x \in B_{R_i}(\bar{\xi}_i)$

$$\begin{aligned} V_\lambda(x) &= \sum_{j=1}^m (V_j(x) - H_j(x)) = w_i(|x - \bar{\xi}_i|) + |\log \lambda| + 2\pi\mu_i H\left(\frac{\lambda}{|\log \lambda|^2}x, \xi_i\right) \\ &\quad + 2\mu_i \log |\log \lambda| - \mu_i \log R_i - (1 - \mu_i)|\log \lambda| \\ &\quad + 2\pi\mu_j G\left(\frac{\lambda}{|\log \lambda|^2}x, \xi_j\right). \end{aligned}$$

Using the equations satisfied by μ_j , (3.15), we find, for $x \in B_{R_i}(\bar{\xi}_i)$

$$\begin{aligned} (3.31) \quad V_\lambda(x) &= w_i(|x - \bar{\xi}_i|) + |\log \lambda| + 2\pi \left[H\left(\frac{\lambda}{|\log \lambda|^2}x, \xi_i\right) - H(\xi_i, \xi_i) \right] \\ &\quad + 2\pi \sum_{j \neq i} \left[G\left(\frac{\lambda}{|\log \lambda|^2}x, \xi_j\right) - G(\xi_i, \xi_j) \right] \\ &= w_i(|x - \bar{\xi}_i|) + |\log \lambda| + O\left(\frac{\lambda}{|\log \lambda|^2}\right), \end{aligned}$$

where the $O\left(\frac{\lambda}{|\log \lambda|^2}\right)$ is in C^1 norm. \square

4. LINEAR THEORY

Here we study the invertibility of the operator

$$L\phi = \Delta\phi + W\phi$$

in Ω_λ where

$$W = f'_\lambda(V_\lambda - |\log \lambda|).$$

Throughout this section we assume that $\xi_1, \dots, \xi_m \in \Omega$ satisfy the separation conditions (3.6), (3.7) and $\bar{\xi}_j = \frac{|\log \lambda|^2}{\lambda} \xi_j \in \Omega_\lambda$. We also assume that μ_1, \dots, μ_m satisfy (3.8).

We consider the linear problem of given h in an appropriate space, finding ϕ and c_{ij} , $i = 1, \dots, m$, $j = 1, 2$, such that

$$(4.1) \quad \begin{cases} L\phi = h + \sum_{i=1}^m \sum_{j=1,2} c_{ij} Z_{ij} & \text{in } \Omega_\lambda \\ \phi = 0 & \text{on } \partial\Omega_\lambda \end{cases}$$

$$(4.2) \quad \int_{\Omega_\lambda} \phi Z_{ij} = 0 \quad \text{for all } i = 1, 2, j = 1, \dots, m,$$

where the functions Z_{ij} are defined by

$$Z_{ij}(x) = z_{ij}(x)\eta_0(x - \bar{\xi}_i)$$

with

$$(4.3) \quad z_{ij}(x) = \frac{\partial V_0}{\partial x_j}(x - \bar{\xi}_i),$$

and η_0 is a smooth radial function in \mathbb{R}^2 with support in $B_{R_0}(0)$ and identically 1 in $B_{R_0/2}(0)$, and V_0 is defined in (2.7). Note that $\frac{\partial V_0}{\partial x_j}(x - \bar{\xi}_j)$ is continuous but not C^1 . The choice of η_0 makes Z_{ij} a smooth function with compact support.

Let Y be the space of measurable functions $h : \Omega_\lambda \rightarrow \mathbb{R}$ such that

$$h \in L^\infty(\hat{\Omega}_\lambda) \text{ and } h \in L^p(B_{2R_0}(\bar{\xi}_j)) \text{ for all } j = 1, \dots, m,$$

where

$$\hat{\Omega}_\lambda = \Omega_\lambda \setminus \cup_{j=1}^m B_{2R_0}(\bar{\xi}_j)$$

and $2 < p < +\infty$ is fixed. We will consider the following norm on Y :

$$\|h\|_Y = \sup_{x \in \hat{\Omega}_\lambda} \left(\sum_{j=1}^m |x - \bar{\xi}_j|^{-2-\sigma} \right)^{-1} |h(x)| + \sum_{j=1}^m \|h\|_{L^p(B_{2R_0}(\bar{\xi}_j))},$$

where $0 < \sigma < 1$ is a small constant. Note that since $p > 2$, if $h \in Y$ then any solution $\phi \in H_0^1(\Omega_\lambda)$ of (4.1) is $C^1(\Omega_\lambda)$.

Proposition 4.1. *There is $\lambda_0 > 0$ such that for any $0 < \lambda \leq \lambda_0$ and for all $h \in Y$ there is a unique $\phi \in L^\infty(\Omega_\lambda)$ and unique $c_{ij} \in \mathbb{R}$ that solve (4.1), (4.2). Moreover*

$$(4.4) \quad \|\phi\|_{L^\infty(\Omega_\lambda)} + \sum_{i=1}^m \sum_{j=1,2} |c_{ij}| \leq C \|h\|_Y.$$

In addition, the maps $\xi_1, \dots, \xi_m \mapsto \phi, c_{ij}$ are differentiable and

$$(4.5) \quad \|\partial_{\bar{\xi}_k} \phi\|_{L^\infty(\Omega_\lambda)} + |\partial_{\bar{\xi}_k} c_{ij}| \leq C \lambda^{1/p-1} \|h\|_Y.$$

The proof of this result relies on the non-degeneracy of V_0 (defined in (2.7)), which satisfies

$$-\Delta V_0 = (V_0)_+ \text{ in } \mathbb{R}^2,$$

and is radially symmetric with $\max_{\mathbb{R}^2} V_0 = 1$.

Proposition 4.2. *Let $\phi \in L^\infty(\mathbb{R}^2)$ be a solution of*

$$-\Delta \phi = \chi_{[V_0 > 0]} \phi \text{ in } \mathbb{R}^2,$$

where $\chi_{[V_0 > 0]}$ is the characteristic function of the set $[V_0 > 0] = B_{R_0}(0)$. Then ϕ is a linear combination of

$$\frac{\partial V_0}{\partial x_1}, \frac{\partial V_0}{\partial x_2}.$$

For the proof see [2, Proposition 3.1].

To prove Proposition 4.1 we start with an apriori estimate.

Lemma 4.3. *There is $C > 0$ such that for all $\lambda > 0$ small, and for any $h \in Y$, $\phi \in L^\infty(\Omega_\lambda)$, and $c_{ij} \in \mathbb{R}$ that verify (4.1), (4.2) we have*

$$\|\phi\|_{L^\infty(\Omega_\lambda)} + \sum_{i=1}^m \sum_{j=1,2} |c_{ij}| \leq C \|h\|_Y.$$

Proof. We first prove that

$$(4.6) \quad |c_{ij}| \leq C \|h\|_Y + o(1) \|\phi\|_{L^\infty(\Omega_\lambda)}$$

as $\lambda \rightarrow 0$. For this let η be a radial function in $C^\infty(\mathbb{R}^2)$ with support in $B_2(0)$ and $\eta \equiv 1$ in $B_1(0)$. Let $\eta_\lambda(x) = \eta(\lambda^{1/2}(x - \bar{\xi}_i))$ and

$$\tilde{Z}_{ij} = z_{ij} \eta_\lambda$$

where z_{ij} is defined in (4.3). Multiplying (4.1) by \tilde{Z}_{ij} we find

$$\int_{\Omega_\lambda} \phi(\Delta \tilde{Z}_{ij} + W \tilde{Z}_{ij}) = \int_{\Omega_\lambda} h \tilde{Z}_{ij} + c_{ij} \int_{\Omega_\lambda} Z_{ij} \tilde{Z}_{ij}.$$

This gives

$$|c_{ij}| \leq C \|h\|_Y + \|\phi\|_{L^\infty(\Omega_\lambda)} \int_{\Omega_\lambda} |\Delta \tilde{Z}_{ij} + W \tilde{Z}_{ij}|.$$

We compute

$$\Delta \tilde{Z}_{ij} + W \tilde{Z}_{ij} = \eta_\lambda (\Delta z_{ij} + W z_{ij}) + 2\nabla \eta_\lambda \nabla z_{ij} + \Delta \eta_\lambda z_{ij}.$$

Using that $|z_{ij}(x)| \leq C|x - \bar{\xi}_i|^{-1}$ and $|\nabla z_{ij}(x)| \leq C|x - \bar{\xi}_i|^{-2}$ for $|x - \bar{\xi}_i| \geq 2R_0$, we see that

$$\int_{\Omega_\lambda} |2\nabla \eta_\lambda \nabla z_{ij} + \Delta \eta_\lambda z_{ij}| = O(\lambda^{1/2})$$

as $\lambda \rightarrow 0$. The other term can be estimated as follows:

$$\int_{\Omega_\lambda} |\eta_\lambda (\Delta z_{ij} + W z_{ij})| \leq \|\eta_\lambda\|_{L^\infty} \|z_{ij}\|_{L^\infty} \int_{\Omega_\lambda} |\chi_{B_{R_0}}(\bar{\xi}_i) - f'_\lambda(V_\lambda - |\log \lambda|)| \rightarrow 0$$

as $\lambda \rightarrow 0$. Therefore

$$\int_{\Omega_\lambda} |\Delta \tilde{Z}_{ij} + W \tilde{Z}_{ij}| = o(1)$$

as $\lambda \rightarrow 0$ and this proves (4.6).

Now we claim that if $R > 0$ is large enough, then

$$(4.7) \quad \|\phi\|_{L^\infty(\Omega_\lambda)} \leq C(\|\phi\|_i + \|h\|_Y + \sum_{i=1}^m \sum_{j=1,2} |c_{ij}|)$$

where

$$\|\phi\|_i = \sup_{\cup_{i=1}^m B_R(\bar{\xi}_i)} |\phi|.$$

For this we use a barrier argument. Let

$$\psi(x) = \sum_{j=1}^m (1 - |x - \bar{\xi}_j|^{-\sigma}).$$

Fix $R > 2R_0$. By (2.17), $W(x) \leq C\lambda^3/|\log \lambda|$ for $x \in \Omega_\lambda \setminus \cup_{i=1}^m B_R(\bar{\xi}_i)$. Then

$$\Delta \psi + W(x)\psi < 0 \quad \text{in } \Omega_\lambda \setminus \cup_{i=1}^m B_R(\bar{\xi}_i),$$

provided $\lambda > 0$ is small. This shows that the operator $\Delta + W$ satisfies the maximum principle in this region. Applying the maximum principle to $(\|h\|_Y + \sum |c_{ij}| + \|\phi\|_i)\psi \pm \phi$ we arrive at (4.7).

Using (4.6) and (4.7) we obtain

$$\|\phi\|_{L^\infty(\Omega_\lambda)} \leq C(\|\phi\|_i + \|h\|_Y).$$

Therefore to prove the lemma it suffices to show

$$(4.8) \quad \|\phi\|_i \leq C\|h\|_Y.$$

We prove this estimate by contradiction. Assume that there are sequences $\lambda_n \rightarrow 0$, (ϕ_n) in $L^\infty(\Omega_{\lambda_n})$, $(c_{ij}^{(n)})$ in \mathbb{R} and (h_n) in Y that satisfy (4.1), (4.2) and

$$\|\phi_n\|_i > n\|h_n\|_Y.$$

By linearity we can assume that $\|\phi_n\|_i = 1$. Then (4.6) implies that $\|h_n\|_Y \rightarrow 0$ and $c_{ij}^{(n)} \rightarrow 0$ as $n \rightarrow +\infty$. Then for a fixed $i \in \{1, \dots, m\}$ and a subsequence (denoted the same as the original sequence)

$$\sup_{B_R(\bar{\xi}_i)} |\phi_n| \geq c$$

for some $c > 0$. By translating we can assume that $\bar{\xi}_i = 0$. Using the equation, we get that up to another subsequence, $\phi_n \rightarrow \phi$ uniformly on compact sets of \mathbb{R}^2 and that $\phi \not\equiv 0$ is a bounded solution of

$$\Delta \phi + \chi_{B_{R_0}(0)} \phi = 0 \quad \text{in } \mathbb{R}^2.$$

By Proposition 4.2, $\phi = a_1 z_{i1} + a_2 z_{i2}$ for some $a_1, a_2 \in \mathbb{R}$. But ϕ also satisfies

$$\int_{\mathbb{R}^2} \phi Z_{ij} = 0 \quad j = 1, 2,$$

which shows that $a_1 = a_2 = 0$ so $\phi \equiv 0$, a contradiction. This proves (4.8) and finishes the proof of the lemma. \square

Proof of Proposition 4.1. Consider the Hilbert space

$$H = \left\{ \phi \in H_0^1(\Omega_\lambda) : \int_{\Omega_\lambda} Z_{ij} \phi = 0 \quad i = 1, \dots, m, j = 1, 2 \right\}$$

with inner product $\langle \phi_1, \phi_2 \rangle = \int_{\Omega_\lambda} \nabla \phi_1 \nabla \phi_2$. For $h \in Y$, the variational problem: find $\phi \in H$ such that

$$\langle \phi, \psi \rangle = \int_{\Omega_\lambda} (-W\phi + h)\phi \quad \text{for all } \phi \in H$$

is a weak formulation of (4.1), (4.2). Using the Riesz representation theorem, this variational problem is equivalent to solve

$$(4.9) \quad \phi + K(\phi) = \tilde{h}$$

where $\tilde{h} \in H$ and $K : H \rightarrow H$ is a compact operator. When $h = 0$ then $\tilde{h} = 0$ and by Lemma 4.3 $\phi = 0$. By the Fredholm alternative there is a solution $\phi \in H$ of (4.9) giving a weak solution of (4.1), (4.2). By standard regularity theory $\phi \in C(\bar{\Omega}_\lambda)$ and we get the estimate (4.4) from Lemma 4.3.

Now we proceed with the differentiability properties of ϕ , c_{ij} with respect to ξ_1, \dots, ξ_m . For this we proceed formally, assuming the differentiability, and obtain estimate (4.5). This argument can then later be justified by applying it to finite differences instead of derivatives. We recall that ϕ is the unique solution of

$$L\phi = h + \sum_{i=1}^m \sum_{j=1,2} c_{ij} Z_{ij} \quad \text{in } \Omega_\lambda$$

satisfying $\phi = 0$ on $\partial\Omega_\lambda$ and the orthogonality conditions (4.2), and that the c_{ij} are uniquely determined. Assuming that ϕ , c_{ij} are differentiable and setting $\psi = \partial_{\bar{\xi}_k} \phi$ we find

$$L\psi = -\partial_{\bar{\xi}_k} W\phi + h + \sum d_{ij} Z_{ij} + \sum c_{ij} \partial_{\bar{\xi}_k} Z_{ij}$$

where $d_{ij} = \partial_{\bar{\xi}_k} c_{ij}$. Differentiating the orthogonality condition (4.2) we get

$$\int_{\Omega_\lambda} \psi Z_{ij} + \phi \partial_{\bar{\xi}_k} Z_{ij} = 0.$$

Setting

$$\tilde{\psi} = \psi + b_{ij} Z_{ij} \quad \text{where} \quad b_{ij} = \frac{\int_{\Omega_\lambda} \phi \partial_{\bar{\xi}_k} Z_{ij}}{\int_{\Omega_\lambda} Z_{ij}^2}$$

we get that $\tilde{\psi}$ satisfies (4.2) and

$$L\tilde{\psi} = -\partial_{\bar{\xi}_k} W\phi + h + \sum d_{ij} Z_{ij} + \sum c_{ij} \partial_{\bar{\xi}_k} Z_{ij} + \sum b_{ij} Z_{ij} \quad \text{in } \Omega_\lambda.$$

Hence, applying the apriori estimate of Lemma 4.3 we deduce

$$\|\psi\|_{L^\infty(\Omega_\lambda)} + |d_{ij}| \leq C(\|\partial_{\bar{\xi}_k} W\phi\|_Y + \|h\|_Y + \sum |c_{ij}| \|\partial_{\bar{\xi}_k} Z_{ij}\|_Y + \sum |b_{ij}| \|Z_{ij}\|_Y).$$

We claim that each term is bounded by $C\lambda^{1/p-1}\|h\|_Y$. Let us verify this explicitly for $\|\partial_{\bar{\xi}_k} W\phi\|_Y$, because the others are direct. Since $\|\phi\|_{L^\infty} \leq C\|h\|_Y$, it suffices to verify that

$$(4.10) \quad \|\partial_{\bar{\xi}_k} W\|_Y \leq C\lambda^{1/p-1}.$$

But $\partial_{\bar{\xi}_k} W = f'_\lambda(V_\lambda - |\log \lambda|) \partial_{\bar{\xi}_k} V_\lambda$. One can verify directly that $\|\partial_{\bar{\xi}_k} V_\lambda\|_{L^\infty(\Omega_\lambda)} \leq C$. We then conclude the validity of (4.10) by using the next lemma. \square

Lemma 4.4. *We have*

$$(4.11) \quad \|f''_\lambda(V_\lambda - |\log \lambda|)\|_Y \leq C\lambda^{1/p-1}$$

for some $C > 0$ and all $\lambda > 0$ small.

Proof. Let us write $\hat{\Omega}_\lambda = \Omega_\lambda \setminus \cup_{j=1}^m B_{2R_0}(\bar{\xi}_j)$. Using (2.20) and that $V_\lambda - |\log \lambda| \leq -a$ for some $a > 0$ on $\hat{\Omega}_\lambda$ we get

$$\sup_{x \in \hat{\Omega}_\lambda} \left(\sum_{j=1}^m |x - \bar{\xi}_j|^{-2-\sigma} \right)^{-1} |f''_\lambda(V_\lambda - |\log \lambda|)| \leq C\lambda^{1-2\sigma}.$$

To estimate $\|f''_\lambda(V_\lambda - |\log \lambda|)\|_{L^p(B_{2R_0}(\bar{\xi}_j))}$ we split the integral

$$\int_{B_{2R_0}(\bar{\xi}_j)} |f''_\lambda(V_\lambda - |\log \lambda|)|^p = \int_{D_1} \dots + \int_{D_2} \dots$$

where $D_1 = B_{R_j+L\lambda}(\bar{\xi}_j) \setminus B_{R_j-L\lambda}(\bar{\xi}_j)$, $D_2 = B_{2R_0}(\bar{\xi}_j) \setminus D_1$ and $L > 0$ is some large fixed number. In D_1 , by (2.18) we estimate $|f''_\lambda| \leq C/\lambda$ and we get

$$\int_{D_1} |f''_\lambda(V_\lambda - |\log \lambda|)|^p \leq C\lambda^{1-p}.$$

To estimate the integral over D_2 we recall that $V_\lambda(x) - |\log \lambda| = w_j(|x - \bar{\xi}_j|) + O(\frac{\lambda}{|\log \lambda|^p})$ for $x \in B_{2R_0}(\bar{\xi}_j)$ (c.f. (3.16)) in C^1 norm. Since $w_j(R_j) = 0$, $V_\lambda - |\log \lambda|$ has a zero set that is at distance $O(\lambda)$ from $\partial B_{R_j}(\bar{\xi}_j)$, and $V_\lambda(x) - |\log \lambda|$ separates linearly from this curve. So thanks to (2.19), $|f''_\lambda(V_\lambda(x) - |\log \lambda|)| \leq C \text{dist}(x, \partial B_{R_j}(\bar{\xi}_j))^{-1}$ on A_2 . Therefore

$$\int_{A_2} |f''_\lambda(V_\lambda - |\log \lambda|)|^p \leq C \int_{L\lambda}^C y^{-p} dy \leq C\lambda^{1-p}.$$

It follows that

$$\|f''_\lambda(V_\lambda - |\log \lambda|)\|_{L^p(B_{2R_0}(\bar{\xi}_j))} \leq C\lambda^{1/p-1},$$

and taking $\sigma > 0$ small we get the stated estimate (4.11). \square

5. A NONLINEAR PROJECTED PROBLEM

In this section we solve the nonlinear problem

$$(5.1) \quad \begin{cases} L\phi + N(\phi) + E_\lambda = \sum_{i=1}^m \sum_{j=1,2} c_{ij} Z_{ij} & \text{in } \Omega_\lambda \\ \phi = 0 & \text{on } \partial\Omega_\lambda \end{cases}$$

with ϕ satisfying

$$(5.2) \quad \int_{\Omega_\lambda} \phi Z_{ij} = 0 \quad \text{for all } i = 1, 2, j = 1, \dots, m.$$

We always assume that $\xi_1, \dots, \xi_m \in \Omega$ satisfy the separation conditions (3.6), (3.7), and $\bar{\xi}_j = \frac{|\log \lambda|^2}{\lambda} \xi_j$.

Proposition 5.1. *Assume μ_1, \dots, μ_m satisfy (3.15). Then there is $\lambda_0 > 0$ such that for $\lambda \in (0, \lambda_0)$ (5.1) has a unique solution $\phi = \phi(\xi_1, \dots, \xi_m)$ and $c_{ij} = c_{ij}(\xi_1, \dots, \xi_m)$ such that*

$$(5.3) \quad \|\phi\|_{L^\infty} + \sum_{i=1}^m \sum_{j=2}^2 |c_{ij}| \leq C\lambda^{1-2\sigma}.$$

Proof. For the argument it is better to work with the space X defined as the set of continuous functions ϕ on $\bar{\Omega}_\lambda$ such that ϕ restricted to $B_{2R_0}(\bar{\xi}_j)$ belongs to $W^{1,\infty}(B_{2R_0}(\bar{\xi}_j))$ for all $j = 1, \dots, m$, equipped with the norm

$$\|\phi\|_X = \|\phi\|_{L^\infty(\Omega_\lambda)} + \sum_{j=1}^m \|\phi\|_{W^{1,\infty}(B_{2R_0}(\bar{\xi}_j))}.$$

We rewrite problem (5.1), (5.2) as the fixed point problem

$$\phi = F(\phi)$$

where $F = -T(N(\phi) + E_\lambda)$ and T is the linear operator defined in Proposition 4.1, which by estimate (4.4) satisfies

$$\|T(h)\|_\infty \leq C\|h\|_Y \quad \text{for all } h \in Y.$$

By elliptic estimates we also deduce

$$\|T(h)\|_X \leq C\|h\|_Y$$

with a constant C independent of λ , and here it is important that $p > 2$.

Let us estimate $\|E_\lambda\|_Y$. Using (2.15) we have

$$\sup_{x \in \hat{\Omega}_\lambda} \left(\sum_{j=1}^m |x - \bar{\xi}_j|^{-2-\sigma} \right)^{-1} |E_\lambda(x)| \leq C\lambda^{1-\sigma} |\log \lambda|^{3+2\sigma} \leq C\lambda^{1-2\sigma}$$

for $\lambda > 0$ small, where $\hat{\Omega}_\lambda = \Omega_\lambda \setminus \cup_{j=1}^m B_{2R_0}(\bar{\xi}_j)$. Also, using Lemma 3.2 we see that

$$|E_\lambda| \leq C \frac{\lambda}{|\log \lambda|^2} \quad \text{in } B_{2R_0}(\bar{\xi}_j).$$

So we deduce

$$(5.4) \quad \|E_\lambda\|_Y \leq C\lambda^{1-2\sigma}.$$

For fixed $\gamma > 0$, let $\mathcal{B} = \{\phi \in X : \|\phi\|_X \leq \gamma\lambda^{1-2\sigma}\}$. We claim that there exist constants $C, a > 0$ such that for $\phi_1, \phi_2 \in \mathcal{B}$

$$(5.5) \quad \|N(\phi_1) - N(\phi_2)\|_Y \leq C\lambda^a \|\phi_1 - \phi_2\|_X,$$

and we can prove this with $a = \frac{1-2\sigma}{p}$.

Indeed,

$$N(\phi_1) - N(\phi_2) = (\phi_1 - \phi_2) \int_0^1 N'(\phi_2 + s(\phi_1 - \phi_2)) ds.$$

Therefore, to obtain (5.5) it is enough to show that for all $s \in [0, 1]$

$$\|N'(\phi_2 + s(\phi_1 - \phi_2))\|_Y \leq C\lambda^a.$$

Part of this norm is $\|N'(\phi_2 + s(\phi_1 - \phi_2))\|_{L^p(B_{2R_0}(\bar{\xi}_j))}$. Write $\phi = \phi_2 + s(\phi_1 - \phi_2)$ and note that $\|\phi\|_X \leq \gamma\lambda^{1-2\sigma}$. Then

$$\int_{B_{2R_0}(\bar{\xi}_j)} |N'(\phi)|^p = \int_{B_{2R_0}(\bar{\xi}_j)} |f'_\lambda(V_\lambda + \phi - |\log \lambda|) - f'_\lambda(V_\lambda - |\log \lambda|)|^p = I_1 + I_2,$$

where

$$I_k = \int_{A_k} |f'_\lambda(V_\lambda + \phi - |\log \lambda|) - f'_\lambda(V_\lambda - |\log \lambda|)|^p, \quad k = 1, 2$$

$$A_1 = B_{R_j + \lambda^m}(\bar{\xi}_j) \setminus B_{R_j - \lambda^m}(\bar{\xi}_j), \quad A_2 = A \setminus A_1,$$

the numbers R_j are given in (3.10) and $m > 0$ is a parameter to be chosen. Since f'_λ is uniformly bounded in the range of its arguments (see (2.16)), we find

$$(5.6) \quad |I_1| \leq C|A_1| \leq C\lambda^m.$$

On A_2 we obtain

$$\begin{aligned} |I_2| &\leq \int_{A_2} |f'_\lambda(V_\lambda + \phi - |\log \lambda|) - f'_\lambda(V_\lambda - |\log \lambda|)|^p \\ &\leq \int_{A_2} \int_0^1 |f''_\lambda(V_\lambda + \tau\phi - |\log \lambda|)|^p |\phi|^p d\tau dx. \end{aligned}$$

We recall that $V_\lambda(x) - |\log \lambda| = w_j(|x - \bar{\xi}_j|) + O(\frac{\lambda}{|\log \lambda|^\sigma})$ for $x \in B_{2R_0}(\bar{\xi}_j)$ (c.f. (3.16)) in C^1 norm, and note that the gradient of $\tau\phi$ is small in uniform norm in $B_{2R_0}(\bar{\xi}_j)$, because $\|\phi\|_X \leq \gamma\lambda^{1-2\sigma}$. This and $w_j(R_j) = 0$ yield

$$|V_\lambda + \tau\phi - |\log \lambda|| \geq c\lambda^m \quad \text{in } A_2$$

for some constant $c > 0$. Therefore, from estimate (2.19)

$$|f''_\lambda(V_\lambda + \tau\phi - |\log \lambda|)| \leq C \text{dist}(x, \partial B_{R_j}(\bar{\xi}_j))^{-1}$$

on A_2 and we get

$$|I_2| \leq C\lambda^{m(1-p)} \|\phi\|_\infty^p \leq C\lambda^{m(1-p) + (1-2\sigma)p}$$

Combining this last estimate with (5.6) we obtain

$$\|N'(\phi_2 + s(\phi_1 - \phi_2))\|_{L^p(B_{2R_0}(\bar{\xi}_j))} \leq C(\lambda^{m/p} + \lambda^{m(1/p-1) + 1 - 2\sigma}).$$

We choose $m = 1 - 2\sigma$ and we obtain

$$(5.7) \quad \|N'(\phi)\|_{L^p(B_{2R_0}(\bar{\xi}_j))} \leq C\lambda^{\frac{1-2\sigma}{p}}.$$

Next we estimate the weighted L^∞ norm of $N'(\phi)$ away from the points $\bar{\xi}_j$. For this we recall that if $|x - \bar{\xi}_j| \geq 2R_0$ for all j , then $V_\lambda - |\log \lambda| \leq -M$ for some fixed $M > 0$ and the same holds for $V_\lambda + s\phi - |\log \lambda|$. Therefore (2.17) yields

$$(5.8) \quad \sup_{x \in \hat{\Omega}_\lambda} \left(\sum_{j=1}^m |x - \bar{\xi}_j|^{-2-\sigma} \right)^{-1} |N'(\phi)| \leq C\lambda^{1-2\sigma}.$$

The combination of (5.7), (5.8) proves (5.5).

Using (5.4) and (5.5), we see that choosing $\gamma > 0$ fixed and large in the definition of \mathcal{B} , for $\lambda > 0$ sufficiently small, F is a contraction in \mathcal{B} , and by the contraction mapping principle F has a unique fixed point in \mathcal{B} . \square

Proposition 5.2. *For $\lambda > 0$ sufficiently small, the maps $\xi_1, \dots, \xi_m \rightarrow \phi, c_{ij}$ constructed in Proposition 5.1 are differentiable in the region defined by (3.6), (3.7) and for any $k = 1, \dots, m$, and $\lambda > 0$ small*

$$(5.9) \quad \|\partial_{\bar{\xi}_k} \phi\|_{L^\infty} + \|\partial_{\bar{\xi}_k} c_{ij}\|_{L^\infty} \leq C\lambda^{1/p-2\sigma}.$$

For the proof we need the following lemma.

Lemma 5.3. *We have*

$$\|\partial_{\bar{\xi}_k} E\|_Y \leq C\lambda^{\min(1-2\sigma, 1/p)}.$$

Proof. We compute $\partial_{\bar{\xi}_k} E_\lambda = \Delta \partial_{\bar{\xi}_k} V_\lambda + f'_\lambda(V_\lambda - |\log \lambda|) \partial_{\bar{\xi}_k} V_\lambda$. It can be verified that $|\partial_{\bar{\xi}_k} V_\lambda| \leq C$ in Ω_λ for some fixed constant.

Let us compute $\partial_{\bar{\xi}_k} E_\lambda(x)$ for $x \in \Omega_\lambda \setminus \cup_{j=1}^m B_{R_j}(\bar{\xi}_j)$. In this region V_λ is harmonic and therefore $\partial_{\bar{\xi}_k} E_\lambda = f'_\lambda(V_\lambda - |\log \lambda|) \partial_{\bar{\xi}_k} V_\lambda$. Therefore, using (2.17) and that $\|\partial_{\bar{\xi}_k} V_\lambda\|_{L^\infty(\Omega_\lambda)} \leq C$ we get

$$(5.10) \quad \sup_{x \in \hat{\Omega}_\lambda} \left(\sum_{j=1}^m |x - \bar{\xi}_j|^{-2-\sigma} \right)^{-1} |E_\lambda| \leq C\lambda^{1-\sigma} |\log \lambda|^{3+2\sigma} \leq C\lambda^{1-2\sigma}$$

for $\lambda > 0$ small, where $\hat{\Omega}_\lambda = \Omega_\lambda \setminus \cup_{j=1}^m B_{2R_0}(\bar{\xi}_j)$.

Let us estimate $\|\partial_{\bar{\xi}_k} E_\lambda\|_{L^p(B_{2R_0}(\bar{\xi}_i))}$. We split $\|\partial_{\bar{\xi}_k} E_\lambda\|_{L^p(B_{2R_0}(\bar{\xi}_i))}^p = I_1 + I_2$ where

$$I_1 = \int_{B_{2R_0}(\bar{\xi}_i) \setminus B_{R_i+L\lambda}(\bar{\xi}_i)} |\partial_{\bar{\xi}_k} E_\lambda|^p \quad I_2 = \int_{B_{R_i+L\lambda}(\bar{\xi}_i) \setminus B_{R_i}(\bar{\xi}_i)} |\partial_{\bar{\xi}_k} E_\lambda|^p$$

where $L > 0$ is some large constant. Using inequality (2.17) we can estimate

$$(5.11) \quad |f'_\lambda(V_\lambda - |\log \lambda|)| \leq \frac{C\lambda^3}{\text{dist}(x, \partial B_{R_i}(\bar{\xi}_i))^3}$$

for $x \in B_{2R_0}(\bar{\xi}_i) \setminus B_{R_i+L\lambda}(\bar{\xi}_i)$. Therefore

$$I_1 \leq C\lambda^{3p} \int_{L\lambda}^{2R_0} \frac{1}{y^{3p}} dy \leq C\lambda.$$

For I_2 we have, using the uniform bound for f'_λ and $\partial_{\bar{\xi}_k} V_\lambda$, $|I_2| \leq C\lambda$. Hence we find

$$(5.12) \quad \int_{B_{2R_0}(\bar{\xi}_i) \setminus B_{R_i}(\bar{\xi}_i)} |\partial_{\bar{\xi}_k} E_\lambda|^p \leq C\lambda.$$

To estimate inside $B_{R_i}(\bar{\xi}_i)$ we use (3.31) to write

$$E_\lambda = f_\lambda(w_i(|x - \bar{\xi}_i|) + g(x)) - f_\lambda(w_i(|x - \bar{\xi}_i|))$$

for $x \in B_{R_i}(\bar{\xi}_i)$, where

$$g(x) = 2\pi \left[H\left(\frac{\lambda}{|\log \lambda|^2}x, \xi_i\right) - H(\xi_i, \xi_i) \right] + 2\pi \sum_{j \neq i} \left[G\left(\frac{\lambda}{|\log \lambda|^2}x, \xi_j\right) - G(\xi_i, \xi_j) \right]$$

We recall that w_i depends on μ_i because the initial condition α_i in the ODE (3.2) is determined by μ_i from the relation (3.9). We make the dependence of the solution w of (3.2) explicit by writing $w = w(r, \alpha)$. We also note that μ_i depend on $\bar{\xi}_j$ through the formula (3.15). Then

$$\partial_{\bar{\xi}_k} E_\lambda = E_1 + E_2$$

where

$$\begin{aligned} E_1 &= [f'_\lambda(w_i(|x - \bar{\xi}_i|) + g(x)) - f'_\lambda(w_i(|x - \bar{\xi}_i|))] \partial_{\bar{\xi}_k} w_i(|x - \bar{\xi}_i|, \alpha_i) \\ E_2 &= f'_\lambda(w_i(|x - \bar{\xi}_i|) + g(x)) \partial_{\bar{\xi}_k} g \end{aligned}$$

Then we compute

$$\partial_{\bar{\xi}_k} w_i(|x - \bar{\xi}_i|, \alpha_i) = -\delta_{ik} \frac{\partial w_i}{\partial r} \frac{x - \bar{\xi}_i}{|x - \bar{\xi}_i|} + \frac{\partial w_i}{\partial \alpha} \sum_j \frac{\partial \alpha_i}{\partial \mu_j} \frac{\partial \mu_j}{\partial \bar{\xi}_k}$$

and we get

$$(5.13) \quad |\partial_{\bar{\xi}_k} w_i(|x - \bar{\xi}_i|, \alpha_i)| \leq C \quad \text{for } x \in B_{R_i}(\bar{\xi}_i).$$

To estimate E_1 we write $A_1 = B_{R_i}(\bar{\xi}_i) \setminus B_{R_i - L\lambda}(\bar{\xi}_i)$, $A_2 = B_{R_i - L\lambda}(\bar{\xi}_i)$ and $L > 0$ is a large constant. Then, using the uniform bounds (5.13) and (2.16) we get $\int_{A_1} |E_1|^p \leq C\lambda$. Since $g(x) = O(\lambda/|\log \lambda|^2)$, we estimate using (2.19)

$$\begin{aligned} \int_{A_2} |E_1|^p &\leq C \int_{A_2} |f'_\lambda(w_i(|x - \bar{\xi}_i|) + g(x)) - f'_\lambda(w_i(|x - \bar{\xi}_i|))|^p \\ &\leq C \int_{A_2} \int_0^1 |f''_\lambda(w_i(|x - \bar{\xi}_i|) + \tau g(x))|^p |g|^p d\tau dx \leq C\lambda. \end{aligned}$$

Also $\int_{B_{R_i}(\bar{\xi}_i)} |E_2|^p \leq C\lambda^p$ and we conclude that

$$(5.14) \quad \int_{B_{R_i}(\bar{\xi}_i)} |\partial_{\bar{\xi}_k} E_\lambda|^p \leq C\lambda.$$

Combining (5.10), (5.12) and (5.14) we obtain the result of the lemma. \square

Proof of Proposition 5.2. The proof that ϕ , c_{ij} are differentiable with respect to ξ_1, \dots, ξ_m can be done with the contraction mapping principle, using that the linear operator defined in Proposition 4.1 is differentiable with respect to ξ_1, \dots, ξ_m . We proceed to prove estimate (5.9). Differentiating the equation in (5.1) with respect to $\bar{\xi}_k$ we obtain, for $\psi = \partial_{\bar{\xi}_k} \phi$,

$$\Delta \psi + W\psi + (\partial_{\bar{\xi}_k} W)\phi + \partial_{\bar{\xi}_k} E + \partial_{\bar{\xi}_k} N(\phi) + \sum (\partial_{\bar{\xi}_k} c_{ij}) Z_{ij} + \sum c_{ij} \partial_{\bar{\xi}_k} Z_{ij} = 0$$

in Ω_λ . Let $\tilde{\psi} = \psi - \sum d_{ij} Z_{ij}$ where $d_{ij} = -\int_{\Omega_\lambda} \phi \partial_{\bar{\xi}_k} Z_{ij} / \int_{\Omega_\lambda} Z_{ij}^2$. In this way, $\tilde{\psi}$ satisfies the orthogonality conditions (4.2). Applying Proposition 4.1 we can estimate

$$\begin{aligned} \|\tilde{\psi}\|_{L^\infty} + \sum |\partial_{\bar{\xi}_k} c_{ij}| &\leq C \left(\|(\partial_{\bar{\xi}_k} W)\phi\|_Y + \|\partial_{\bar{\xi}_k} E\|_Y + \|\partial_{\bar{\xi}_k} N(\phi)\|_Y \right. \\ &\quad \left. + \sum |c_{ij}| \|\partial_{\bar{\xi}_k} Z_{ij}\|_Y + \sum |d_{ij}| \|\Delta Z_{ij} + W Z_{ij}\|_Y \right) \end{aligned}$$

Estimate (5.3) gives that $|c_{ij}| = O(\lambda^{1-2\sigma})$ and $|d_{ij}| = O(\lambda^{1-2\sigma})$ as $\lambda \rightarrow 0$.

Lemma 5.3 yields

$$\|\partial_{\bar{\xi}_k} E\|_Y \leq C \lambda^{\min(1-2\sigma, 1/p)}.$$

We compute

$$(\partial_{\bar{\xi}_k} W)\phi + \partial_{\bar{\xi}_k} N(\phi) = \left[f'_\lambda(V_\lambda + \phi - |\log \lambda|) - f'_\lambda(V_\lambda - |\log \lambda|) \right] (\partial_{\bar{\xi}_k} V_\lambda + \psi).$$

The same computations that lead to (5.7) and (5.8) show that

$$\|(f'_\lambda(V_\lambda + \phi - |\log \lambda|) - f'_\lambda(V_\lambda - |\log \lambda|))\|_Y \leq C \lambda^{\frac{1-2\sigma}{p}}.$$

Therefore

$$\|(\partial_{\bar{\xi}_k} W)\phi + \partial_{\bar{\xi}_k} N(\phi)\|_Y \leq C(\lambda^{1/p-2\sigma} + \lambda^{\frac{1-2\sigma}{p}}) \|\psi\|_{L^\infty}.$$

Collecting the estimates above, we obtain

$$\|\psi\|_{L^\infty} \leq C(\lambda^{1/p-2\sigma} + C \lambda^{\frac{1-2\sigma}{p}}) \|\psi\|_{L^\infty}.$$

For $\lambda > 0$ small we deduce (5.9). \square

6. PROOF OF THE MAIN RESULTS

We work with points $\xi_j \in \Omega$ satisfying (3.6), (3.7), that is, in the set

$$\tilde{\Omega}_m = \{(\xi_1, \dots, \xi_m) \in \Omega^m : |\xi_i - \xi_j| \geq \delta \forall i \neq j, \text{dist}(\xi_j, \partial\Omega) \geq \delta \forall 1 \leq j \leq m\},$$

where $\delta > 0$ is small. Recall that $\bar{\xi}_j = \frac{|\log \lambda|^2}{\lambda} \xi_j \in \Omega_\lambda$. We also work with μ_j given by (3.15).

For $\xi = (\xi_1, \dots, \xi_m) \in \tilde{\Omega}_m$, let $\phi(\xi)$ denote the solution of (3.14) that satisfies $\|\phi\|_\infty \leq C \lambda^{1-2\sigma}$ constructed in Proposition 5.1 and let $c_{ij}(\xi)$ denote the constants appearing in equation (5.1).

Writing the initial approximation $V_\lambda = V_\lambda(\xi)$ (defined in (3.13)), we set

$$J_\lambda(\xi) = I_\lambda(V_\lambda(\xi) + \phi(\xi))$$

where I_λ is the functional given in (3.19).

Lemma 6.1. *If $\xi = (\xi_1, \dots, \xi_j) \in \tilde{\Omega}_m$ is a critical point of J_λ then $c_{ij}(\xi) = 0$ for all $i = 1, \dots, m$ and all $j = 1, 2$, so that $V_\lambda(\xi) + \phi(\xi)$ is a solution of (3.1).*

The proof of this lemma is very similar to Lemma 4.1 in [2] or Lemma 5.1 in [5].

Lemma 6.2. *We have the expansion*

$$I_\lambda(V_\lambda + \phi_\lambda) = I_\lambda(V_\lambda) + \Theta_\lambda(\xi)$$

as $\lambda \rightarrow 0$, where $\Theta_\lambda(\xi) = o(1)$ in the C^1 norm in $\tilde{\Omega}_m$.

Proof. Using that

$$DI_\lambda(V_\lambda + \phi_\lambda)[\phi_\lambda] = - \sum c_{ij} \int_{\Omega_\lambda} Z_{ij} \phi_\lambda = 0$$

we compute

$$\begin{aligned} I_\lambda(V_\lambda + \phi_\lambda) - I_\lambda(V_\lambda) &= - \int_0^1 s \int_{\Omega_\lambda} D^2 I_\lambda(V_\lambda + s\phi_\lambda)[\phi_\lambda, \phi_\lambda] dx ds \\ &= - \int_0^1 s \int_{\Omega_\lambda} (f_\lambda(V_\lambda + \phi_\lambda - |\log \lambda|) - f_\lambda(V_\lambda - |\log \lambda|)) \phi_\lambda \\ &\quad + \int_0^1 s \int_{\Omega_\lambda} (f'_\lambda(V_\lambda + s\phi_\lambda - |\log \lambda|)\phi_\lambda - E_\lambda) \phi_\lambda \end{aligned}$$

We compute with detail the estimate for the derivative of $I_\lambda(V_\lambda + \phi_\lambda) - I_\lambda(V_\lambda)$ with respect to $\bar{\xi}_k$. The estimate for the C^0 norm is similar. Differentiating with respect to $\bar{\xi}_k$,

$$\partial_{\bar{\xi}_k} (I_\lambda(V_\lambda + \phi_\lambda) - I_\lambda(V_\lambda)) = \int_0^1 s(I_1 + I_2 + \dots) ds$$

where

$$\begin{aligned} I_1 &= - \int_{\Omega_\lambda} (f'_\lambda(V_\lambda + \phi_\lambda - |\log \lambda|) - f'_\lambda(V_\lambda - |\log \lambda|)) (\partial_{\bar{\xi}_k} V_\lambda) \phi_\lambda \\ I_2 &= - \int_{\Omega_\lambda} (f'_\lambda(V_\lambda + \phi_\lambda - |\log \lambda|) - 2f'_\lambda(V_\lambda + s\phi_\lambda)) (\partial_{\bar{\xi}_k} \phi_\lambda) \phi_\lambda \\ I_3 &= - \int_{\Omega_\lambda} f''_\lambda(V_\lambda + s\phi_\lambda - |\log \lambda|) (\partial_{\bar{\xi}_k} (V_\lambda + \phi_\lambda)) \phi_\lambda^2 \\ I_4 &= - \int_{\Omega_\lambda} (\partial_{\bar{\xi}_k} E_\lambda) \phi_\lambda + E_\lambda (\partial_{\bar{\xi}_k} \phi_\lambda) \\ I_5 &= - \int_{\Omega_\lambda} (f_\lambda(V_\lambda + \phi_\lambda - |\log \lambda|) - f_\lambda(V_\lambda - |\log \lambda|)) \partial_{\bar{\xi}_k} \phi_\lambda. \end{aligned}$$

Each term I_i will be estimated below, but before, it will be convenient to have

$$(6.1) \quad \int_{\Omega_\lambda} |f'_\lambda(V_\lambda + s\phi_\lambda - |\log \lambda|)| \leq C |\log \lambda|,$$

by (2.16), (2.17), and

$$(6.2) \quad \int_{\Omega_\lambda} |f''_\lambda(V_\lambda + s\phi_\lambda - |\log \lambda|)| \leq C |\log \lambda|,$$

which is proved by a computation similar to Lemma 4.4.

We start with

$$\begin{aligned} |I_1| &\leq \int_{\Omega_\lambda} |f'_\lambda(V_\lambda + \phi_\lambda - |\log \lambda|) - f'_\lambda(V_\lambda - |\log \lambda|)| |\partial_{\bar{\xi}_k} V_\lambda| |\phi_\lambda| \\ &\leq \|\partial_{\bar{\xi}_k} V_\lambda\|_{L^\infty} \|\phi_\lambda\|_{L^\infty} \int_{\Omega_\lambda} |f'_\lambda(V_\lambda + \phi_\lambda - |\log \lambda|) - f'_\lambda(V_\lambda - |\log \lambda|)| \\ &\leq C \|\phi_\lambda\|_{L^\infty}^2 \int_0^1 \int_{\Omega_\lambda} |f''_\lambda(V_\lambda + \tau\phi - |\log \lambda|)| d\tau \\ &\leq C \lambda^{2-4\sigma} |\log \lambda|, \end{aligned}$$

where we have used (5.3) and (6.2). Next we estimate

$$\begin{aligned} |I_2| &\leq \int_0^1 s \int_{\Omega_\lambda} (|f'_\lambda(V_\lambda + \phi_\lambda - |\log \lambda|)| + |f'_\lambda(V_\lambda + s\phi_\lambda - |\log \lambda|)|) |\partial_{\bar{\xi}_k} \phi_\lambda| |\phi_\lambda| \\ &\leq C \|\partial_{\bar{\xi}_k} \phi_\lambda\|_{L^\infty} \|\phi_\lambda\|_{L^\infty} \leq C \lambda^{1+1/p-4\sigma} \end{aligned}$$

by (5.3), (5.9) and (6.1). Similarly

$$|I_3| \leq \int_0^1 \int_{\Omega_\lambda} |f''_\lambda(V_\lambda + s\phi_\lambda - |\log \lambda|) \partial_{\bar{\xi}_k} (V_\lambda + \phi) \phi_\lambda^2| \leq C \lambda^{2-4\sigma} |\log \lambda|.$$

and

$$|I_4| \leq \|\phi_\lambda\|_{L^\infty} \int_{\Omega_\lambda} |\partial_{\bar{\xi}_k} E_\lambda| + \|\partial_{\bar{\xi}_k} \phi_\lambda\|_{L^\infty} \int_{\Omega_\lambda} |E_\lambda| \leq C \lambda^{1+1/p-2\sigma} |\log \lambda|^3,$$

since outside big ball around the $\bar{\xi}_i$, $|f_\lambda|, |f'_\lambda| \leq C \lambda^3 / |\log \lambda|$ and the area of Ω_λ is proportional to $|\log \lambda|^4 / \lambda^2$. In summary

$$|\partial_{\bar{\xi}_k} (I_\lambda(V_\lambda + \phi_\lambda) - I_\lambda(V_\lambda))| \leq C \lambda^{1+1/p-4\sigma}.$$

We choose now $p > 2$ close to 1 and $\sigma > 0$ small. Since $\partial_{\bar{\xi}_k} = \frac{|\log \lambda|^2}{\lambda} \partial_{\xi_k}$ we deduce

$$|\partial_{\xi_k} I_\lambda(V_\lambda + \phi_\lambda) - \partial_{\xi_k} I_\lambda(V_\lambda)| \leq C \lambda^a$$

for some $a > 0$. □

Proof of Theorems 1.1, 1.2, 1.3. According to Lemma 6.1, the function $V_\lambda(\xi) + \phi(\xi)$ is a solution of problem (3.1) if we adjust $\xi = (\xi_1, \dots, \xi_m)$ so that it is a critical point of $J_\lambda(\xi) = I_\lambda(V_\lambda(\xi) + \phi(\xi))$. This is equivalent to finding a critical point of

$$\tilde{J}_\lambda(\xi) = -\frac{1}{\pi R_0(2\pi - \log R_0)} [I_\lambda(V_\lambda) - \pi m R_0 |\log \lambda| + 2\pi m R_0 \log |\log \lambda|].$$

Thanks to Lemmas 3.3 and 6.2 for $\xi \in \tilde{\Omega}_m$ we have

$$(6.3) \quad \tilde{J}_\lambda(\xi) = \varphi_m(\xi) + \Theta_\lambda(\xi).$$

where $\Theta_\lambda = o(1)$ as $\lambda \rightarrow 0$ in the C^1 norm of $\tilde{\Omega}_m$. To prove Theorem 1.1, we note that the function φ_m is C^1 , bounded from above in $\tilde{\Omega}_m$, and satisfies

$$\varphi_m(\xi_1, \dots, \xi_m) \rightarrow -\infty \text{ as } |\xi_i - \xi_j| \rightarrow 0 \text{ for some } i \neq j.$$

Hence, choosing $\delta > 0$ small, φ_m has a strict absolute maximum in $\tilde{\Omega}_m$. By (6.3), \tilde{J}_λ has also an absolute maximum in $\tilde{\Omega}_m$ for $\lambda > 0$ small, and this yields the existence of a solution to problem (3.1).

Theorem 1.3 is a direct consequence of the 6.3, since a non-degenerate critical point of φ_m gives rise to critical to a critical point of a small C^1 perturbation of this function.

To prove Theorem 1.2 we invoke the work [5], where it is proved that if Ω is not simply connected, then φ_m and any sufficiently C^1 close map have critical points in $\tilde{\Omega}_m$, for $\delta > 0$ small enough. This result also appears in [2]. □

Acknowledgment. J.D. was supported by Fondecyt 1090167, CAPDE-Anillo ACT-125 and Fondo Basal CMM. J.W was supported by GRF grant "On Elliptic Equations with Negative Exponent" from RGC of Hong Kong. We thank Professor M.J. Ward for useful discussions.

REFERENCES

- [1] S. Baraket, F. Pacard, *Construction of singular limits for a semilinear elliptic equation in dimension 2*. Calc. Var. Partial Differential Equations **6** (1998), no. 1, 1–38.
- [2] D. Cao, S. Peng, S. Yan, *Multiplicity of solutions for the plasma problem in two dimensions*. Advances in Mathematics **225** (2010), 2741–2785.
- [3] L. A. Caffarelli, A. Friedman, *Asymptotic estimates for the plasma problem*. Duke Math. J. **47** (1980), 705–742.
- [4] C. Chen, C. Lin, *Topological degree for a mean field equation on Riemann surfaces*. Comm. Pure Appl. Math. **56** (2003), no. 12, 1667–1727.
- [5] M. del Pino, M. Kowalczyk, M. Musso, *Singular limits in Liouville-type equations*. Calc. Var. Partial Differential Equations **24** (2005), no. 1, 47–81.
- [6] P. Esposito, N. Ghoussoub, Y. Guo, *Compactness along the branch of semistable and unstable solutions for an elliptic problem with a singular nonlinearity*. Comm. Pure Appl. Math. **60** (2007), 1731–1768.
- [7] P. Esposito, N. Ghoussoub, Y. Guo, *Mathematical analysis of partial differential equations modeling electrostatic MEMS*. Courant Lecture Notes in Mathematics, 20. Courant Institute of Mathematical Sciences, New York; American Mathematical Society, Providence, RI, 2010.
- [8] P. Esposito, M. Grossi, A. Pistoia, *On the existence of blowing-up solutions for a mean field equation*. Ann. Inst. H. Poincaré Anal. Non Linéaire **22** (2005), no. 2, 227–257.
- [9] N. Ghoussoub, Y. Guo, *On the partial differential equations of electrostatic MEMS devices: stationary case*. SIAM J. Math. Anal. **38** (2006/07), 1423–1449.
- [10] Z. Guo, J. Wei, *Asymptotic behavior of touch-down solutions and global bifurcations for an elliptic problem with a singular nonlinearity*. Commun. Pure Appl. Anal. **7** (2008), no. 4, 765–786.
- [11] Z. Guo, J. Wei, *Infinitely many turning points for an elliptic problem with a singular nonlinearity*. J. Lond. Math. Soc. (2) **78** (2008), 21–35.
- [12] D.D. Joseph, T.S. Lundgren, *Quasilinear Dirichlet problems driven by positive sources*. Arch. Rational Mech. Anal. **49** (1972/73), 241–269.
- [13] A. E. Lindsay, M. J. Ward, *Asymptotics of some nonlinear eigenvalue problems for a MEMS capacitor. I. Fold point asymptotics*. Methods Appl. Anal. **15** (2008), 297–325.
- [14] J. A. Pelesko, T. A. Driscoll, *The Effect of the Small Aspect Ratio Approximation on Canonical Electrostatic MEMS Models*. J. Engrg. Math., **53**, No. 3-4, (2005), 239–252.
- [15] J.A. Pelesko, D.H. Bernstein, *D Modeling MEMS and NEMS*. Chapman & Hall/CRC, Boca Raton, FL, 2003.
- [16] P.H. Rabinowitz, *Some global results for nonlinear eigenvalue problems*. J. Functional Analysis **7** (1971), 487–513.
- [17] J. Wei, D. Ye, *On MEMS equation with fringing field*. Proc. Amer. Math. Soc. **138** (2010), 1693–1699.

J. DÁVILA - DEPARTAMENTO DE INGENIERÍA MATEMÁTICA AND CMM, UNIVERSIDAD DE CHILE, CASILLA 170 CORREO 3, SANTIAGO, CHILE.

E-mail address: `jdavila@dim.uchile.cl`

J. WEI - DEPARTMENT OF MATHEMATICS, CHINESE UNIVERSITY OF HONG KONG, SHATIN, HONG KONG.

E-mail address: `wei@math.cuhk.edu.hk`