

**MAT6082 Topics in Analysis**2<sup>nd</sup> term, 2015-16**Teacher:** Professor Ka-Sing Lau**Schedule:** Wednesday, 2.30-5.00 pm**Venue:** LSB 222**Topics:** Introduction to Stochastic Calculus

In the past thirty years, there has been an increasing demand of stochastic calculus in mathematics as well as various disciplines such as mathematical finance, pde, physics and biology. The course is a rigorous introduction to this topic. The material include conditional expectation, Markov property, martingales, stochastic processes, Brownian motions, Ito's calculus, and stochastic differential equations.

**Prerequisites**

Students are expected to have good background in real analysis, probability theory and some basic knowledge of stochastic processes.

**References:**

1. A Course in Probability Theory, K.L. Chung, (1974).
2. Measure and Probability, P. Billingsley, (1986).
3. Introduction to Stochastic Integration, H.H. Kuo, (2006).
4. Intro. to Stochastic Calculus with Application, F. Klebaner, (2001).
5. Brownian Motion and Stoch. Cal., I. Karatzas and S. Shreve, (1998).
6. Stoch. Cal. for Finance II– Continuous time model, S. Shreve, (2004).

Everyone knows calculus deals with deterministic objects. On the other hand *stochastic calculus* deals with random phenomena. The theory was introduced by Kiyosi Ito in the 40's, and therefore stochastic calculus is also called *Ito calculus*. Besides its interest in mathematics, it has been used extensively in statistical mechanics in physics, the filter and control theory in engineering. Nowadays it is very popular in the option price and hedging in finance. For example the well-known Black-Scholes model is

$$dS(t) = rS(t)dt + \sigma S(t)dB(t)$$

where  $S(t)$  is the stock price,  $\sigma$  is the volatility, and  $r$  is the interest rate, and  $B(t)$  is the Brownian motion. The most important notion for us is the Brownian motion. As is known the botanist R. Brown (1828) discovered certain zigzag random movement of pollens suspended in liquid. A. Einstein (1915) argued that the movement is due to bombardment of particle by the molecules of the fluid. He set up some basic equations of Brownian motion and use them to study diffusion. It was N. Wiener (1923) who made a rigorous study of the Brownian motion using the then new theory of Lebesgue measure. Because of that a Brownian motion is also frequently called a Wiener process.

Just like calculus is based on the *fundamental theorem of calculus*, the Ito calculus is based on the *Ito Formula*: Let  $f$  be a twice differentiable function on  $\mathbb{R}$ , then

$$f(B(t)) - f(B(0)) = \int_0^t f'(B(t))dB(t) + \frac{1}{2} \int_0^t f''(B(t))dt$$

where  $B(0) = 0$  to denote the motion starts at 0. There are formula for integration, for example, we have

$$\int_0^T B(t)dB(t) = \frac{1}{2}B(t)^2 - \frac{1}{2}T; \quad \int_0^T tdB(t) = TB(T) - \int_0^T B(t)dt.$$

In this course, the prerequisite is real analysis and basic probability theory. In real analysis, one needs to know  $\sigma$ -fields, measurable functions, measures

and integration theory, various convergence theorems, Fubini theorem and the Radon-Nikodym theorem. We will go through some of the probability theory on conditional expectation, optional r.v. (stopping time), Markov property, martingales ([1], [2]). Then we will go onto study the Brownian motion ([2], [3], [5]), the stochastic integration and the Ito calculus ([3], [4], [5]).



# Chapter 1

## Basic Probability Theory

### 1.1 Preliminaries

Let  $\Omega$  be a set and let  $\mathcal{F}$  be a family of subsets of  $\Omega$ ,  $\mathcal{F}$  is called a *field* if it satisfies

- (i)  $\emptyset, \Omega \in \mathcal{F}$ ;
- (ii) for any  $A \in \mathcal{F}$ ,  $A^c \in \mathcal{F}$ ;
- (iii) for any  $A, B \in \mathcal{F}$ ,  $A \cup B \in \mathcal{F}$  (hence  $A \cap B \in \mathcal{F}$ ).

It is called a  $\sigma$ -*field* if (iii) is replaced by

- (iii)' for any  $\{A_n\}_{n=1}^{\infty} \subset \mathcal{F}$ ,  $\cup_{n=1}^{\infty} A_n \in \mathcal{F}$  (hence  $\cap_{n=1}^{\infty} A_n \in \mathcal{F}$ ).

If  $\Omega = \mathbb{R}$  and  $\mathcal{F}$  is the smallest  $\sigma$ -field generated by the open sets, then we call it the Borel field and denote by  $\mathcal{B}$ .

A *probability space* is a triple  $(\Omega, \mathcal{F}, P)$  such that  $\mathcal{F}$  is a  $\sigma$ -field in  $\Omega$ , and  $P : \mathcal{F} \rightarrow [0, 1]$  satisfies

(i)  $P(\Omega) = 1$

(ii) countable additivity : if  $\{A_n\}_{n=1}^{\infty} \subseteq \mathcal{F}$  is a disjoint family, then

$$P\left(\bigcup_{n=1}^{\infty} A_n\right) = \sum_{n=1}^{\infty} P(A_n).$$

We call  $\Omega$  a sample space,  $A \in \mathcal{F}$  an event (or measurable set) and  $P$  a probability measure on  $\Omega$ ; an element  $\omega \in \Omega$  is called an outcome.

**Theorem 1.1.1.** (*Caratheodory Extension Theorem*) Let  $\mathcal{F}_0$  be a field of subsets in  $\Omega$  and let  $\mathcal{F}$  be the  $\sigma$ -field generated by  $\mathcal{F}_0$ . Let  $P : \mathcal{F}_0 \rightarrow [0, 1]$  satisfies (i) and (ii) (on  $\mathcal{F}_0$ ). Then  $P$  can be extended uniquely to  $\mathcal{F}$ , and  $(\Omega, \mathcal{F}, P)$  is a probability space.

The proof of the theorem is to use the outer measure argument.

**Example 1.** Let  $\Omega = [0, 1]$ , let  $\mathcal{F}_0$  be the family of set consisting of finite disjoint unions of half open intervals  $(a, b]$  and  $[0, b]$ , Let  $P([a, b]) = |b - a|$ . Then  $\mathcal{F}$  is the Borel field and  $P$  is the Lebesgue measure on  $[0, 1]$ .

**Example 2.** Let  $\{(\Omega_n, \mathcal{F}_n, P_n)\}_n$  be a sequence of probability spaces. Let  $\Omega = \prod_{n=1}^{\infty} \Omega_n$  be the product space and let  $\mathcal{F}_0$  be the family of subsets of the form  $E = \prod_{n=1}^{\infty} E_n$ , where  $E_n \in \mathcal{F}_n$ ,  $E_n = \Omega_n$  except for finitely many  $n$ . Define

$$P(E) = \prod_{n=1}^{\infty} P(E_n)$$

Let  $\mathcal{F}$  be the  $\sigma$ -field generated  $\mathcal{F}_0$ , then  $(\Omega, \mathcal{F}, P)$  is the standard infinite product measure space.

**Example 3.** (Kolmogorov Extension Theorem) Let  $P_n$  be probability measures on  $(\prod_{k=1}^n \Omega_k, \mathcal{F}_n)$  satisfying the following consistency condition: for  $m \leq n$

$$P_n \circ \pi_{nm}^{-1} = P_m$$

where  $\pi_{nm}(x_1 \cdots x_n) = (x_1 \cdots x_m)$ . On  $\Omega = \prod_{k=1}^{\infty} \Omega_k$ , we let  $\mathcal{F}_0$  be the field of sets  $F = E \times \prod_{k=n+1}^{\infty} \Omega_k$ ,  $E \in \mathcal{F}_n$  and let

$$P(F) = P_n(E).$$

Then this defines a probability spaces  $(\Omega, \mathcal{F}, P)$ , where  $\mathcal{F}$  is the  $\sigma$ -field generated by  $\mathcal{F}_0$ .

**Remark:** The probability space in Example 2 is the underlying space for a sequence of independent random variables. Example 3 is for more general sequence of random variables (with the consistency condition).

A random variable (r.v.)  $X$  on  $(\Omega, \mathcal{F})$  is an (extended) real valued function  $X : (\Omega, \mathcal{F}) \rightarrow \mathbb{R}$  such that for any Borel subset  $B$  of  $\mathbb{R}$ ,

$$X^{-1}(B) = \{\omega : X(\omega) \in B\} \in \mathcal{F}.$$

(i.e.  $X$  is  $\mathcal{F}$ -measurable). We denote this by  $X \in \mathcal{F}$ . It is well known that

- For  $X \in \mathcal{F}$ ,  $X$  is either a simple function (i.e.,  $\sum_{k=1}^n a_k \chi_{A_k}(\omega)$  where  $A_k \in \mathcal{F}$ ), or is the pointwise limit of a sequence of simple functions.
- Let  $X \in \mathcal{F}$  and  $g$  is a Borel measurable function, then  $g(X) \in \mathcal{F}$ .
- If  $\{X_n\} \subseteq \mathcal{F}$  and  $\lim_{n \rightarrow \infty} X_n = X$ , then  $X \in \mathcal{F}$ .
- Let  $\mathcal{F}_X$  be the  $\sigma$ -field generated by  $X$ , i.e., the sub- $\sigma$ -field  $\{X^{-1}(B) : B \in \mathcal{B}\}$ . Then for any  $Y \in \mathcal{F}_X$ ,  $Y = \varphi(X)$  for some extended-valued Borel function  $\varphi$  on  $\mathbb{R}$ .

*Sketch of proof* ([1, p.299]): First prove this for simple r.v.  $Y$  so that  $Y = \phi(X)$  for some simple function  $\phi$ . For a bounded r.v.  $Y \geq 0$ , we can find a sequence of increasing simple functions  $\{Y_n\}$  such that  $Y_n = \phi_n(X)$  and

$Y_n \nearrow Y$ . Let  $\phi(x) = \overline{\lim}_n \phi_n(x)$ , hence  $Y = \phi(X)$ . Then prove  $Y$  for the general case.

A r.v.  $X : (\Omega, \mathcal{F}) \rightarrow \mathbb{R}$  induces a *distribution (function)* on  $\mathbb{R}$ :

$$F(x) = F_X(x) = P(X \leq x).$$

It is a non-decreasing, right continuous function with  $\lim_{n \rightarrow -\infty} F(x) = 0$ ,  $\lim_{n \rightarrow \infty} F(x) = 1$ . The distribution defines a measure  $\mu$

$$\mu((a, b]) = F(b) - F(a)$$

(use the Caratheodory Extension Theorem here). More directly, we can define  $\mu$  by

$$\mu(B) = P(X^{-1}(B)) , \quad B \in \mathcal{B}.$$

The jump of  $F$  at  $x$  is  $F(x) - F(x-) = P(X = x)$ . A r.v.  $X$  is called a *discrete* if  $F$  is a jump function;  $X$  is called a *continuous* r.v. if  $F$  is continuous, i.e.,  $P(X = x) = 0$  for each  $x \in \mathbb{R}$ , and  $X$  is said to have a density function  $f(x)$  if  $F$  is absolutely continuous with the Lebesgue measure and  $f(x) = F'(x)$  a.e., equivalently  $F(x) = \int_{-\infty}^x f(y)dy$ .

For two random variables  $X, Y$  on  $(\Omega, \mathcal{F})$ , the random vector  $(X, Y) : (\Omega, \mathcal{F}) \rightarrow \mathbb{R}^2$  induces a distribution  $F$  on  $\mathbb{R}^2$

$$F(x, y) = P(X \leq x, Y \leq y)$$

and  $F$  is called the joint distribution of  $(X, Y)$ , the corresponding measure  $\mu$  is given by

$$\mu((a, b] \times (c, d]) = F(b, d) - F(a, d) - F(b, c) + F(a, c),$$

Similarly we can define the joint distribution  $F(x_1 \cdots x_n)$  and the corresponding measure.



For a sequence of r.v.,  $\{X_n\}_{n=1}^\infty$ , there are various notions of convergence.

(a)  $X_n \rightarrow X$  a.e. (or a.s.) if  $\lim_{n \rightarrow \infty} X_n(\omega) = X(\omega)$  (pointwise) for  $\omega \in \Omega \setminus E$  where  $P(E) = 0$ .

(b)  $X_n \rightarrow X$  in probability if for any  $\epsilon > 0$ ,  $\lim_{n \rightarrow \infty} P(|X_n - X| \geq \epsilon) = 0$ .

(c)  $X_n \rightarrow X$  in distribution if  $F_n(x) \rightarrow F(x)$  at every point  $x$  of continuity. It is equivalent to  $\mu_n \rightarrow \mu$  vaguely i.e.,  $\mu_n(f) \rightarrow \mu(f)$  for all  $f \in C_0(\mathbb{R})$ , the space of continuous functions vanish at  $\infty$  (detail in [1]).

The following relationships are basic ([1] or Royden):  $(a) \Rightarrow (b) \Rightarrow (c)$ ;  $(b) \Rightarrow (a)$  on some subsequence. On the other hand we cannot expect  $(c)$  to imply  $(b)$  as the distribution does not determine  $X$ . For example consider the interval  $[0, 1]$  with the Lebesgue measure, the r.v.'s  $X_1 = \chi_{[0, \frac{1}{2}]}$ ,  $X_2 = \chi_{[\frac{1}{2}, 1]}$ ,  $X_3 = \chi_{[0, \frac{1}{4}]} + \chi_{[\frac{3}{4}, 1]}$  all have the same distribution.

The expectation of a random variable is defined as

$$E(X) = \int_{\Omega} X(\omega) dP(\omega) = \int_{-\infty}^{\infty} x dF(x) (= \int_{-\infty}^{\infty} x d\mu(x))$$

and for a Borel measurable  $h$ , we have

$$E(h(X)) = \int_{\Omega} h(X(\omega)) dP(\omega) = \int_{-\infty}^{\infty} h(x) dF(x).$$

The most basic convergence theorems are:

(a) Fatou lemma:

$$X_n \geq 0, \quad \text{then } E(\underline{\lim}_{n \rightarrow \infty} X_n) \leq \underline{\lim}_{n \rightarrow \infty} E(X_n).$$

(b) Monotone convergence theorem:

$$X_n \geq 0, \quad X_n \nearrow X, \quad \text{then } \lim_{n \rightarrow \infty} E(X_n) = E(X).$$

(c) Dominated convergence theorem:

$$|X_n| \leq Y, E(Y) < \infty \text{ and } X_n \rightarrow X \text{ a.e.}, \text{ then } \lim_{n \rightarrow \infty} E(X_n) = E(X).$$

We say that  $X_n \rightarrow X$  in  $L^p, p > 0$  if  $E(|X|^p) < \infty$  and  $E(|X_n - X|^p) \rightarrow 0$  as  $n \rightarrow \infty$ . It is known that  $L^p$  convergence implies convergence in probability. The converse also holds if we assume further  $E(|X_n|^p) \rightarrow E(|X|^p) < \infty$  ([1], p.97).

Two events  $A, B \in \mathcal{F}$  are said to be *independent* if

$$P(A \cap B) = P(A)P(B).$$

Similarly we say that the events  $A_1, \dots, A_n \in \mathcal{F}$  are independent if for any subsets  $A_{j_1}, \dots, A_{j_k}$ ,

$$P\left(\bigcap_{i=1}^k A_{j_i}\right) = \prod_{i=1}^k P(A_{j_i}).$$

Two sub- $\sigma$ -fields  $\mathcal{F}_1$  and  $\mathcal{F}_2$  are said to be independent if any choice of sets of each of these  $\sigma$ -fields are independent. Two r.v.'s  $X, Y$  are independent if the  $\sigma$ -fields  $\mathcal{F}_X$  and  $\mathcal{F}_Y$  they generated are independent. Equivalently we have

$$P(X \leq x, Y \leq y) = P(X \leq x) P(Y \leq y),$$

(i.e., the joint distribution equals the product of their marginal distributions).

We say that  $X_1 \dots X_n$  are independent if for any  $X_{i_1} \dots X_{i_k}$ , their joint distribution is a product of their marginal distributions.

**Proposition 1.1.2.** *Let  $X, Y$  be independent, then  $f(X)$  and  $g(Y)$  are independent for any Borel measurable functions  $f$  and  $g$ .*

**Exercises**

1. Can you identify the interval  $[0, 1]$  with the Lebesgue measure to the probability space for tossing a fair coin repeatedly?
2. Prove Proposition 1.1.2.
3. Suppose that  $\sup_n |X_n| \leq Y$  and  $E(Y) < \infty$ . Show that

$$E(\overline{\lim}_{n \rightarrow \infty} X_n) \geq \overline{\lim}_{n \rightarrow \infty} E(X_n)$$

4. If  $p > 0$  and  $E(|X|^p) < \infty$ , then  $\lim_{n \rightarrow \infty} x^p P(|X| > x) = 0$ . Conversely, if  $\lim_{n \rightarrow \infty} x^p P(|X| > x) = 0$ , then  $E(|X|^{p-\epsilon}) < \infty$  for  $0 < \epsilon < p$ .
5. For any d.f.  $F$  and any  $a \geq 0$ , we have

$$\int_{-\infty}^{\infty} (F(x+a) - F(x)) dx = a$$

6. Let  $X$  be a positive r.v. with a distribution  $F$ , then

$$\int_0^{\infty} (1 - F(x)) dx = \int_0^{\infty} x dF(x).$$

and

$$E(X) = \int_0^{\infty} P(X > x) dx = \int_0^{\infty} P(X \geq x) dx$$

7. Let  $\{X_n\}$  be a sequence of identically distributed r.v. with finite mean, then

$$\lim_{n \rightarrow \infty} \frac{1}{n} E(\max_{1 \leq j \leq n} |X_j|) = 0.$$

(Hint: use Ex.6 to express the mean of the maximum)

8. If  $X_1, X_2$  are independent r.v.'s each takes values  $+1$  and  $-1$  with probability  $\frac{1}{2}$ , then the three r.v.'s  $\{X_1, X_2, X_1 X_2\}$  are pairwise independent but not independent.
9. A r.v. is independent of itself if and only if it is constant with probability one. Can  $X$  and  $f(X)$  be independent when  $f \in \mathcal{B}$ ?

- 10 .** Let  $\{X_j\}_{j=1}^n$  be independent with distributions  $\{F_j\}_{j=1}^n$ . Find the distribution for  $\max_j X_j$  and  $\min_j X_j$ .
- 11.** If  $X$  and  $Y$  are independent and  $E(|X + Y|^p) < \infty$  for some  $p > 0$ , then  $E(|X|^p) < \infty$  and  $E(|Y|^p) < \infty$ .
- 12.** If  $X$  and  $Y$  are independent,  $E(|X|^p) < \infty$  for some  $p \geq 1$ , and  $E(Y) = 0$ , then  $E(|X + Y|^p) \geq E(|X|^p)$ .

## 1.2 Conditional Expectation

Let  $\Lambda \in \mathcal{F}$  with  $P(\Lambda) > 0$ , we define

$$P(E|\Lambda) = \frac{P(\Lambda \cap E)}{P(\Lambda)} \quad \text{where } P(\Lambda) > 0.$$

It follows that for a discrete random vector  $(X, Y)$ ,

$$P(Y = y|X = x) = \begin{cases} \frac{P(Y = y, X = x)}{P(X = x)}, & \text{if } P(X = x) > 0, \\ 0, & \text{otherwise.} \end{cases}$$

Moreover if  $(X, Y)$  is a continuous random variable with joint density  $f(x, y)$ , the conditional density of  $Y$  given  $X = x$  is

$$f(y|x) = \begin{cases} \frac{f(x, y)}{f_X(x)}, & \text{if } f_X(x) > 0, \\ 0, & \text{otherwise.} \end{cases}$$

where  $f_X(x) = \int_{-\infty}^{\infty} f(x, y)dy$  is the marginal density. The conditional expectation of  $Y$  given  $X = x$  is

$$E(Y|X = x) = \int_{-\infty}^{\infty} yf(y|x)dy.$$

Note that

$$g(x) := E(Y|X = x) \quad \text{is a function on } x,$$

and hence

$$g(X(\cdot)) := E(Y|X(\cdot)) \quad \text{is a r.v. on } \Omega. \quad (1.2.1)$$

In the following we have a more general consideration for the conditional expectation (and also the conditional probability):  $E(Y|\mathcal{G})$  where  $\mathcal{G}$  is a sub- $\sigma$ -field of  $\mathcal{F}$ .

First let us look at a special case where  $\mathcal{G}$  is generated by a measurable partition  $\{\Lambda_n\}_n$  of  $\Omega$  (each member in  $\mathcal{G}$  is a union of  $\{\Lambda_n\}_n$ ). Let  $Y$  be an

integrable r.v., then

$$E(Y|\Lambda_n) = \int_{\Omega} Y(\omega) dP_{\Lambda_n}(\omega) = \frac{1}{P(\Lambda_n)} \int_{\Lambda_n} Y(\omega) dP(\omega). \quad (1.2.2)$$

(Here  $P_{\Lambda_n}(\cdot) = \frac{P(\cdot \cap \Lambda_n)}{P(\Lambda_n)}$  is a probability measure for  $P(\Lambda_n) > 0$ ). Consider the random variable (as in (1.2.1))

$$Z(\cdot) = E(Y|\mathcal{G})(\cdot) := \sum_n E(Y|\Lambda_n) \chi_{\Lambda_n}(\cdot) \in \mathcal{G}.$$

It is easy to see that if  $\omega \in \Lambda_n$ , then  $Z(\omega) = E(Y|\Lambda_n)$ , and moreover

$$\int_{\Omega} E(Y|\mathcal{G}) dP = \sum_n \int_{\Lambda_n} E(Y|\mathcal{G}) dP = \sum_n E(Y|\Lambda_n) P(\Lambda_n) = \int_{\Omega} Y dP.$$

We can also replace  $\Omega$  by  $\Lambda \in \mathcal{G}$  and obtain

$$\int_{\Lambda} E(Y|\mathcal{G}) dP = \int_{\Lambda} Y dP \quad \forall \Lambda \in \mathcal{G}.$$

Recall that for  $\mu, \nu$  two  $\sigma$ -finite measures on  $(\Omega, \mathcal{F})$  and  $\mu \geq 0, \nu$  is called *absolutely continuous* with respect to  $\mu$  ( $\nu \ll \mu$ ) if for any  $\Lambda \in \mathcal{F}$  and  $\mu(\Lambda) = 0$ , then  $\nu(\Lambda) = 0$ . The Radon-Nikodym theorem says that there exists  $g = \frac{d\nu}{d\mu}$  such that

$$\nu(\Lambda) = \int_{\Lambda} g d\mu \quad \forall \Lambda \in \mathcal{F}.$$

**Theorem 1.2.1.** *If  $E(|Y|) < \infty$  and  $\mathcal{G}$  is a sub- $\sigma$ -field of  $\mathcal{F}$ , then there exists a unique  $\mathcal{G}$ -measurable r.v., denote by  $E(Y|\mathcal{G}) \in \mathcal{G}$ , such that*

$$\int_{\Lambda} Y dP = \int_{\Lambda} E(Y|\mathcal{G}) dP \quad \forall \Lambda \in \mathcal{G}.$$

**Proof.** Consider the set-valued function

$$\nu(\Lambda) = \int_{\Lambda} Y dP \quad \Lambda \in \mathcal{G}.$$

Then  $\nu$  is a “signed measure” on  $\mathcal{G}$ . It satisfies

$$P(\Lambda) = 0 \implies \nu(\Lambda) = 0.$$

Hence  $\nu$  is absolutely continuous with respect to  $P$ . By the Radon-Nikodym theorem, the derivative  $g = \frac{d\nu}{dP} \in \mathcal{G}$  and

$$\int_{\Lambda} Y dP = \nu(\Lambda) = \int_{\Lambda} g dP \quad \forall \Lambda \in \mathcal{G}.$$

This  $g$  is unique: for if we have  $g_1 \in \mathcal{G}$  satisfies the same identity,

$$\int_{\Lambda} Y dP = \nu(\Lambda) = \int_{\Lambda} g_1 dP \quad \forall \Lambda \in \mathcal{G}.$$

Let  $\Lambda = \{g > g_1\} \in \mathcal{G}$ , then  $\int_{\Lambda} (g - g_1) dP = 0$  implies that  $P(\Lambda) = 0$ . We can reverse  $g$  and  $g_1$  and hence we have  $P(g \neq g_1) = 0$ . It follows that  $g = g_1$   $\mathcal{G}$ -a.e.

**Definition 1.2.2.** Given an integrable r.v.  $Y$  and a sub- $\sigma$ -field  $\mathcal{G}$ , we say that  $E(Y|\mathcal{G})$  is the conditional expectation of  $Y$  with respect to  $\mathcal{G}$  (also denote by  $E_{\mathcal{G}}(Y)$ ) if it satisfies

$$(a) \quad E(Y|\mathcal{G}) \in \mathcal{G};$$

$$(b) \quad \int_{\Lambda} Y dP = \int_{\Lambda} E(Y|\mathcal{G}) dP \quad \forall \Lambda \in \mathcal{G}.$$

If  $Y = \chi_{\Delta} \in \mathcal{F}$ , we define  $P(\Delta|\mathcal{G}) = E(\chi_{\Delta}|\mathcal{G})$  and call this the conditional probability with respect to  $\mathcal{G}$ .

Note that the conditional probability can be put in the following way:

$$(a)' \quad P(\Delta|\mathcal{G}) \in \mathcal{G};$$

$$(b)' \quad P(\Delta \cap \Lambda) = \int_{\Lambda} P(\Delta|\mathcal{G}) dP \quad \forall \Lambda \in \mathcal{G}.$$

It is a simple exercise to show that the original definition of  $P(\Delta|\Lambda)$  agrees with this new definition by taking  $\mathcal{G} = \{\emptyset, \Lambda, \Lambda^c, \Omega\}$ .

Note that  $E(Y|\mathcal{G})$  is “almost everywhere” defined, and we call one such function as a “version” of the conditional expectation. For brevity we will not mention the “a.e.” in the conditional expectation unless necessary. If  $\mathcal{G}$  is the sub- $\sigma$ -field  $\mathcal{F}_X$  generated by a r.v.  $X$ , we write  $E(Y|X)$  instead of  $E(Y|\mathcal{F}_X)$ . Similarly we can define  $E(Y|X_1, \dots, X_n)$ .

**Proposition 1.2.3.** *For  $E(Y|X) \in \mathcal{F}_X$ , there exists an extended-valued Borel measurable  $\varphi$  such that  $E(Y|X) = \varphi(X)$ , and  $\varphi$  is given by*

$$\varphi = \frac{d\lambda}{d\mu},$$

where  $\lambda(B) = \int_{X^{-1}(B)} Y dP$ ,  $B \in \mathcal{B}$ , and  $\mu$  is the associated probability of the r.v.  $X$  on  $\mathbb{R}$ .

**Proof.** Since  $E(Y|X) \in \mathcal{F}_X$ , we can write  $E(Y|X) = \varphi(X)$  for some Borel measurable  $\varphi$  (see §1). For  $\Lambda \in \mathcal{F}$ , there exists  $B \in \mathcal{B}$  such that  $\Lambda = X^{-1}(B)$ . Hence

$$\int_{\Lambda} E(Y|X) dP = \int_{\Omega} \chi_B(X) \varphi(X) dP = \int_{\mathbb{R}} \chi_B(X) \varphi(X) d\mu = \int_B \varphi(x) d\mu$$

On the other hand by the definition of conditional probability,

$$\int_{\Lambda} E(Y|X) dP = \int_{X^{-1}(B)} Y dP = \lambda(B).$$

It follows that  $\lambda(B) = \int_B \varphi(x) d\mu$  for all  $B \in \mathcal{B}$ . Hence  $\varphi = \frac{d\lambda}{d\mu}$ .  $\square$

The following are some simple facts of the conditional expectation:

- If  $\mathcal{G} = \{\phi, \Omega\}$ , then  $E(Y|\mathcal{G})$  is a constant function and equals  $E(Y)$ .
- If  $\mathcal{G} = \{\phi, \Lambda, \Lambda^c, \Omega\}$ , then  $E(Y|\mathcal{G})$  is a simple function which equals  $E(Y|\Lambda)$  on  $\Lambda$ , and equals  $E(Y|\Lambda^c)$  on  $\Lambda^c$ ,



- If  $\mathcal{G} = \mathcal{F}$  or  $Y \in \mathcal{G}$ , then  $E(Y|\mathcal{G}) = Y$ .
- If  $(X, Y)$  has a joint density function, then  $E(Y|X)$  coincides with the expression in (1.2.1).

Using the defining relationship of conditional expectation, we can show that the linearity, the basic inequalities and the convergence theorems for  $E(\cdot)$  also hold for  $E(\cdot | \mathcal{G})$ . For example we have

**Proposition 1.2.4.** (*Jensen inequality*) *If  $\varphi$  is a convex function on  $\mathbb{R}$ , and  $Y$  and  $\varphi(Y)$  are integrable r.v., then for each sub- $\sigma$ -algebra  $\mathcal{G}$ ,*

$$\varphi(E(Y|\mathcal{G})) \leq E(\varphi(Y)|\mathcal{G})$$

**Proof.** If  $Y$  is a simple r.v., then  $Y = \sum_{j=1}^n y_j \chi_{\Lambda_j}$  with  $\Lambda \in \mathcal{F}$ . It follows that

$$E(Y|\mathcal{G}) = \sum_{j=1}^n y_j E(\chi_{\Lambda_j}|\mathcal{G}) = \sum_{j=1}^n y_j P(Y_{\Lambda_j}|\mathcal{G})$$

and

$$E(\varphi(Y)|\mathcal{G}) = \sum_{j=1}^n \varphi(y_j) P(Y_{\Lambda_j}|\mathcal{G}).$$

Since  $\sum_{j=1}^n P(\Lambda_j|\mathcal{G}) = 1$ , the inequality holds by the convexity of  $\varphi$ .

In general we can find a sequence of simple r.v.  $\{Y_m\}$  with  $|Y_m| \leq |Y|$  and  $Y_m \rightarrow Y$ , then apply the above together with the dominated convergence theorem.  $\square$

**Proposition 1.2.5.** *Let  $Y$  and  $YZ$  be integrable r.v. and  $Z \in \mathcal{G}$ , then we have*

$$E(YZ|\mathcal{G}) = ZE(Y|\mathcal{G}).$$

**Proof.** It suffices to show that for  $Y, Z \geq 0$

$$\int_{\Lambda} Z E(Y|\mathcal{G}) dP = \int_{\Lambda} ZY dP \quad \forall \Lambda \in \mathcal{G}.$$

Obviously, this is true for  $Z = \chi_{\Delta}$ ,  $\Delta \in \mathcal{G}$ . We can pass it to the simple r.v. Then use the monotone convergence theorem to show that it hold for all  $Z \geq 0$ , and then the general integrable r.v.  $\square$

**Proposition 1.2.6.** *Let  $\mathcal{G}_1$  and  $\mathcal{G}_2$  be sub- $\sigma$ -fields of  $\mathcal{F}$  and  $\mathcal{G}_1 \subseteq \mathcal{G}_2$ . Then for  $Y$  integrable r.v.*

$$E(E(Y|\mathcal{G}_2)|\mathcal{G}_1) = E(Y|\mathcal{G}_1) = E(E(Y|\mathcal{G}_1)|\mathcal{G}_2). \quad (1.2.3)$$

Moreover

$$E(Y|\mathcal{G}_1) = E(Y|\mathcal{G}_2) \quad \text{iff} \quad E(Y|\mathcal{G}_2) \in \mathcal{G}_1. \quad (1.2.4)$$

**Proof.** Let  $\Lambda \in \mathcal{G}_1$ , then  $\Lambda \in \mathcal{G}_2$ . Hence

$$\int_{\Lambda} E(E(X|\mathcal{G}_2)|\mathcal{G}_1) dP = \int_{\Lambda} E(Y|\mathcal{G}_2) dP = \int_{\Lambda} Y dP = \int_{\Lambda} E(Y|\mathcal{G}_1) dP,$$

and the first identity in (1.2.3) follows. The second identity is by  $E(Y|\mathcal{G}_1) \in \mathcal{G}_2$  (recall that  $Z \in \mathcal{G}$  implies  $E(Z|\mathcal{G}) = Z$ ).

For the last part, the necessity is trivial, and the sufficiency follows from the first identity.  $\square$

As a simple consequence, we have

**Corollary 1.2.7.**  $E(E(Y|X_1, X_2)|X_1) = E(Y|X_1) = E(E(Y|X_1)|X_1, X_2)$ .

**Exercises**

1. (Bayes' rule) Let  $\{\Lambda_n\}$  be a  $\mathcal{F}$ -measurable partition of  $\Omega$  and let  $E \in \mathcal{F}$  with  $P(E) > 0$ . Then

$$P(\Lambda_n|E) = \frac{P(\Lambda_n) P(E|\Lambda_n)}{\sum_n P(\Lambda_n) P(E|\Lambda_n)} .$$

2. If the random vector  $(X, Y)$  has probability density  $p(x, y)$  and  $X$  is integrable, then one version of  $E(X|X + Y = z)$  is given by

$$\int xp(x, z - x)dx / \int p(x, z - x)dx .$$

3. Let  $X$  be a r.v. such that  $P(X > t) = e^{-t}$ ,  $t > 0$ . Compute  $E(X|X \vee t)$  and  $E(X|X \wedge t)$  for  $t > 0$ . ( Here  $\vee$  and  $\wedge$  mean maximum and minimum respectively.)

4. If  $X$  is an integrable r.v.,  $Y$  is a bounded r.v., and  $\mathcal{G}$  is a sub- $\sigma$ -field, then

$$E(E(X|\mathcal{G})Y) = E(XE(Y|\mathcal{G})).$$

5. Prove that  $\text{var}(E(Y|\mathcal{G})) \leq \text{var}(Y)$ .

6. Let  $X, Y$  be two r.v., and let  $\mathcal{G}$  be a sub- $\sigma$ -field. Suppose

$$E(Y^2|\mathcal{G}) = X^2, \quad E(Y|\mathcal{G}) = X,$$

then  $Y = X$  a.e.

7. Give an example that  $E(E(Y|X_1)|X_2) \neq E(E(Y|X_2)|X_1)$ . (*Hint: it suffices to find an example  $E(X|Y) \neq E(E(X|Y)|X)$  for  $\Omega$  to have three points*).

### 1.3 Markov Property

Let  $A$  be an index set and let  $\{\mathcal{F}_\alpha : \alpha \in A\}$  be family of sub- $\sigma$ -fields of  $\mathcal{F}$ . We say that the family of  $\mathcal{F}_\alpha$ 's are *conditionally independent* relative to  $\mathcal{G}$  if for any  $\Lambda_i \in \mathcal{F}_{\alpha_i}$   $i = 1, \dots, n$ ,

$$P\left(\bigcap_{j=1}^n \Lambda_j \mid \mathcal{G}\right) = \prod_{j=1}^n P(\Lambda_j \mid \mathcal{G}). \quad (1.3.1)$$

**Proposition 1.3.1.** *For  $\alpha \in A$ , let  $\mathcal{F}^{(\alpha)}$  denote the sub- $\sigma$ -field generated by  $\mathcal{F}_\beta$ ,  $\beta \in A \setminus \{\alpha\}$ . Then the family  $\{\mathcal{F}_\alpha\}_\alpha$  are conditionally independent relative to  $\mathcal{G}$  if and only if*

$$P(\Lambda \mid \mathcal{F}^{(\alpha)} \vee \mathcal{G}) = P(\Lambda \mid \mathcal{G}), \quad \Lambda \in \mathcal{F}_\alpha$$

where  $\mathcal{F}^{(\alpha)} \vee \mathcal{G}$  is the sub- $\sigma$ -field generated by  $\mathcal{F}^{(\alpha)}$  and  $\mathcal{G}$ .

**Proof.** We only prove the case  $A = \{1, 2\}$ , i.e.,

$$P(\Lambda \mid \mathcal{F}_2 \vee \mathcal{G}) = P(\Lambda \mid \mathcal{G}), \quad \Lambda \in \mathcal{F}_1. \quad (1.3.2)$$

The general case follows from the same argument. To prove the sufficiency, we assume (1.3.2). To check (1.3.1), let  $\Lambda \in \mathcal{F}_1$ , then for  $M \in \mathcal{F}_2$ ,

$$\begin{aligned} P(\Lambda \cap M \mid \mathcal{G}) &= E(P(\Lambda \cap M \mid \mathcal{F}_2 \vee \mathcal{G}) \mid \mathcal{G}) \\ &= E(P(\Lambda \mid \mathcal{F}_2 \vee \mathcal{G}) \chi_M \mid \mathcal{G}) \\ &= E(P(\Lambda \mid \mathcal{G}) \chi_M \mid \mathcal{G}) \quad (\text{by (1.3.2)}) \\ &= P(\Lambda \mid \mathcal{G}) P(M \mid \mathcal{G}). \end{aligned}$$

Hence  $\mathcal{F}_1$  and  $\mathcal{F}_2$  are  $\mathcal{G}$ -independent.

To prove the necessity, suppose (1.3.1) holds, we claim that for  $\Delta \in \mathcal{G}$ ,  $\Lambda \in \mathcal{F}_1$  and  $M \in \mathcal{F}_2$ ,

$$\int_{M \cap \Delta} P(\Lambda \mid \mathcal{G}) dP = \int_{M \cap \Delta} P(\Lambda \mid \mathcal{F}_2 \vee \mathcal{G}) dP$$

Since the sets of the form  $M \cap \Delta$  generate  $\mathcal{G} \vee \mathcal{F}_2$ , we have  $P(\Lambda|\mathcal{G}) = P(\Lambda|\mathcal{F}_2 \vee \mathcal{G})$ . i.e., (1.3.2) holds.

The claim follows from the following: let  $\Lambda \in \mathcal{F}_1$ ,  $M \in \mathcal{F}_2$ , then

$$\begin{aligned} E(P(\Lambda|\mathcal{G})\chi_M|\mathcal{G}) &= P(\Lambda|\mathcal{G})P(M|\mathcal{G}) \\ &= P(\Lambda \cap M | \mathcal{G}) \quad (\text{by (1.3.1)}) \\ &= E(P(\Lambda|\mathcal{F}_2 \vee \mathcal{G})\chi_M|\mathcal{G}) \quad \square \end{aligned}$$

**Corollary 1.3.2.** *Let  $\{X_\alpha\}_{\alpha \in A}$  be a family of r.v. and let  $\mathcal{F}_\alpha$  be the sub- $\sigma$ -field generated by  $X_\alpha$ . Then the  $X_\alpha$ 's are independent if and only if for any Borel set  $B$ ,*

$$P(X_\alpha \in B | \mathcal{F}^{(\alpha)}) = P(X_\alpha \in B).$$

Moreover the above condition can be replaced by: for any integrable  $Y \in \mathcal{F}_\alpha$ ,

$$E(Y | \mathcal{F}^{(\alpha)}) = E(Y).$$

**Proof.** The first identity follows from Proposition 1.3.1 by taking  $\mathcal{G}$  as the trivial  $\sigma$ -field. The second one follows from an approximation by simple function and use the first identity.  $\square$

To consider the Markov property, we first consider an important basic case.

**Theorem 1.3.3.** *Let  $\{X_n\}_{n=1}^\infty$  be a sequence of independent r.v. and each  $X_n$  has a distribution  $\mu_n$  on  $\mathbb{R}$ . Let  $S_n = \sum_{j=1}^n X_j$ . Then for  $B \in \mathcal{B}$ ,*

$$P(S_n \in B | S_1, \dots, S_{n-1}) = P(S_n \in B | S_{n-1}) = \mu_n(B - S_{n-1})$$

(Hence  $S_n$  is independent of  $S_1, \dots, S_{n-2}$  given  $S_{n-1}$ .)

**Proof.** We divide the proof into two steps.

*Step 1.* We show that

$$P(X_1 + X_2 \in B \mid X_1) = \mu_2(B - X_1)$$

First observe that  $\mu_2(B - X_1)$  is in  $\mathcal{F}_{X_1}$ . Let  $\Lambda \in \mathcal{F}_{X_1}$ , then  $\Lambda = X_1^{-1}(A)$  for some  $A \in \mathcal{B}$ , and

$$\begin{aligned} \int_{\Lambda} \mu_2(B - X_1) dP &= \int_A \mu_2(B - x_1) d\mu_1(x_1) \\ &= \int_A \left( \int_{x_1+x_2 \in B} d\mu_2(x_2) \right) d\mu_1(x_1) \\ &= \iint_{x_1 \in A, x_1+x_2 \in B} d(\mu_1 \times \mu_2)(x_1, x_2) \\ &= P(X_1 \in A, X_1 + X_2 \in B) \\ &= \int_{\Lambda} P(X_1 + X_2 \in B \mid \mathcal{F}_{X_1}) dP \end{aligned}$$

This implies that  $\mu_2(B - X_1) = P(X_1 + X_2 \in B \mid X_1)$  .

*Step 2.* The second equality in the proposition follows from Step 1 by applying to  $S_{n-1}$  and  $X_n$ . To prove the first identity, we let  $\mu^n = \mu_1 \times \cdots \times \mu_n = \mu^{n-1} \times \mu_n$ . Let  $B_j \in \mathcal{B}$ ,  $1 \leq j \leq n-1$ , and let  $\Lambda = \bigcap_{j=1}^{n-1} S_j^{-1}(B_j) \in \mathcal{F}(S_1, \dots, S_{n-1})$ . We show as in Step 1,

$$\int_{\Lambda} \mu_n(B - S_{n-1}) dP = \int_{\Lambda} P(S_n \in B \mid S_1, \dots, S_{n-1}) dP$$

and the identity  $\mu_n(B - S_{n-1}) = P(S_n \in B \mid S_1, \dots, S_{n-1})$  follows.  $\square$

**Definition 1.3.4.** We call a sequence of random variables  $\{X_n\}_{n=0}^{\infty}$  a (discrete time) stochastic process. It is called a Markov process (Markov chain if the state space is countable or finite) if for any  $n$  and  $B \in \mathcal{B}$ ,

$$P(X_{n+1} \in B \mid X_0, \dots, X_n) = P(X_{n+1} \in B \mid X_n).$$

Let  $I \subseteq \mathbb{N}_0 := \mathbb{N} \cup \{0\}$  and let  $\mathcal{F}_I$  denote the sub- $\sigma$ -field generated by  $\mathcal{F}_n$ ,  $n \in I$ . Typically,  $I = \{n\}$ , or  $[0, n]$ , or  $(n, \infty)$ ;  $\mathcal{F}_{\{n\}}$  denotes the events at the present,  $\mathcal{F}_{[0, n]}$  denotes the events from the past up to the present, and  $\mathcal{F}_{(n, \infty)}$  denotes the events in the future. The above Markov property means the future depends on the present and is independent of the past.

One of the most important examples of Markov process is the sequence  $\{S_n\}_{n=0}^\infty$  in Theorem 1.2.3.

**Theorem 1.3.5.** *Let  $\{X_n\}_{n=0}^\infty$  be a stochastic process, then the following are equivalent:*

(a)  $\{X_n\}_{n=0}^\infty$  has the Markov property;

(b)  $P(M|\mathcal{F}_{[0, n]}) = P(M|X_n)$  for all  $n \in \mathbb{N}$  and  $M \in \mathcal{F}_{(n, \infty)}$ ;

(c)  $P(M_1 \cap M_2 | X_n) = P(M_1|X_n) P(M_2|X_n)$  for all  $M_1 \in \mathcal{F}_{[0, n]}$ ,  $M_2 \in \mathcal{F}_{(n, \infty)}$  and  $n \in \mathbb{N}$ .

The conditions remain true if  $\mathcal{F}_{(n, \infty)}$  is replaced by  $\mathcal{F}_{[n, \infty)}$  (Exercise). Condition (c) can be interpreted as conditioning on the present, the past and the future are independent.

**Proof.** (b)  $\Rightarrow$  (c). Let  $Y_i = \chi_{M_i}$  with  $M_1 \in \mathcal{F}_{[0, n]}$ ,  $M_2 \in \mathcal{F}_{(n, \infty)}$ , then

$$\begin{aligned} P(M_1|X_n) P(M_2|X_n) &= E(Y_1|X_n) E(Y_2|X_n) = E(Y_1 E(Y_2|X_n)|X_n) \\ &= E(Y_1 E(Y_2|\mathcal{F}_{[0, n]})|X_n) = E(E(Y_1 Y_2|\mathcal{F}_{[0, n]})|X_n) \\ &= E(Y_1 Y_2|X_n) = P(M_1 \cap M_2 | X_n). \end{aligned}$$

(c)  $\Rightarrow$  (b). Let  $\Lambda \in \mathcal{F}_{[0, n]}$  be the test set, and let  $Y_1 = \chi_\Lambda$ ,  $Y_2 = \chi_M \in$

$\mathcal{F}_{(0,\infty)}$ . Then

$$\begin{aligned} \int_{\Lambda} P(M|X_n) dP &= E(Y_1 E(Y_2|X_n)) = E(E(Y_1 E(Y_2|X_n))|X_n) \\ &= E(E(Y_1|X_n)E(Y_2|X_n)) = E(E(Y_1 Y_2|X_n)) \\ &= \int_{\Omega} P(\Lambda \cap M|X_n) dP = P(\Lambda \cap M). \end{aligned}$$

This implies  $P(M|X_n) = P(M|\mathcal{F}_{[0,n]})$ .

(b)  $\Rightarrow$  (a) is trivial.

(a)  $\Rightarrow$  (b). We claim that for each  $n$ ,

$$E(Y|\mathcal{F}_{[0,n]}) = E(Y|X_n) \quad \forall Y \in \mathcal{F}_{[n+1,n+k]}, \quad k = 1, 2, \dots \quad (1.3.3)$$

This will establish (b) for  $M \in \bigcup_{k=1}^{\infty} \mathcal{F}_{(n,n+k)}$ ; this family of  $M$  generates  $\mathcal{F}_{(0,\infty)}$ .

Note that the Markov property implies (1.3.3) is true for  $k = 1$ . Suppose the statement is true for  $k$ , we consider  $Y = Y_1 Y_2 \in \mathcal{F}_{[n+1,n+k+1]}$ , where  $Y_1 \in \mathcal{F}_{[n+1,n+k]}$  and  $Y_2 \in \mathcal{F}_{n+k+1}$ . Then

$$\begin{aligned} E(Y|\mathcal{F}_{[0,n]}) &= E(E(Y|\mathcal{F}_{[0,n+k]}) | \mathcal{F}_{[0,n]}) \\ &= E(Y_1 E(Y_2|\mathcal{F}_{[0,n+k]}) | \mathcal{F}_{[0,n]}) \\ &= E(Y_1 E(Y_2|\mathcal{F}_{n+k}) | \mathcal{F}_{[0,n]}) \quad (\text{by Markov}) \\ &= E(Y_1 E(Y_2|\mathcal{F}_{n+k}) | \mathcal{F}_n) \quad (\text{by induction}) \\ &= E(Y_1 E(Y_2|\mathcal{F}_{[n,n+k]}) | \mathcal{F}_{[0,n]}) \quad (\text{by Markov}) \\ &= E(E(Y_1 Y_2|\mathcal{F}_{[n,n+k]}) | \mathcal{F}_{[0,n]}) \\ &= E(Y_1 Y_2|\mathcal{F}_n) \\ &= E(Y|\mathcal{F}_n). \end{aligned}$$

This implies the inductive step for  $Y = \chi_{M_1 \cap M_2} = \chi_{M_1} \chi_{M_2}$  with  $M_1 \in \mathcal{F}_{[n+1,n+k]}$  and  $M_2 \in \mathcal{F}_{n+k+1}$ . But the class of all such  $Y$  generates  $\mathcal{F}_{[n+1,n+k]}$ .

This implies the claim and completes the proof of the theorem.  $\square$



The following random variable plays a central role in stochastic process.

**Definition 1.3.6.** A r.v.  $\alpha : \Omega \rightarrow \mathbb{N}_0 \cup \{\infty\}$  is called a stopping time (or Markov time or optional r.v. ) with respect to  $\{X_n\}_{n=0}^\infty$  if

$$\{\omega : \alpha(\omega) = n\} \in \mathcal{F}_{[0,n]} \quad \text{for each } n \in \mathbb{N}_0 \cup \{\infty\}.$$

It is easy to see the definition can be replaced by  $\{\omega : \alpha(\omega) \leq n\} \in \mathcal{F}_{[0,n]}$ . In practice, the most important example is: for a given  $A \in \mathcal{B}$ , let

$$\alpha_A(\omega) = \min\{n \geq 0 : X_n(\omega) \in A\}.$$

( $\alpha_A(\omega) = \infty$  if  $X_n(\omega) \notin A$  for all  $n$ .) This is the r.v. of the first time the process  $\{X_n\}_{n=0}^\infty$  enters  $A$ . It is clear that

$$\{\omega : \alpha_A(\omega) = n\} = \bigcap_{j=0}^{n-1} \{\omega : X_j(\omega) \in A^c, X_n(\omega) \in A\} \in \mathcal{F}_{[0,n]},$$

and similarly for  $n = \infty$ . Hence  $\alpha_A$  is a stopping time.

Very often  $\alpha$  represents the random time that a specific event happens, and  $\{X_{\alpha+n}\}_{n=1}^\infty$  is the process after the event has occurred. We will use the following terminologies:

– The pre- $\alpha$  field  $\mathcal{F}_\alpha$  is the sets  $\Lambda \in \mathcal{F}_{[0,\infty)}$  of the form

$$\Lambda = \bigcup_{0 \leq n < \infty} \{\{\alpha = n\} \cap \Lambda_n\}, \quad \Lambda_n \in \mathcal{F}_{[0,n]}. \quad (1.3.4)$$

It follows that  $\Lambda \in \mathcal{F}_\alpha$  if and only if  $\{\alpha = n\} \cap \Lambda \in \mathcal{F}_n$  for each  $n$ .

– The post  $\alpha$ -process is  $\{X_{\alpha+n}\}_{n=1}^\infty$  where  $X_{\alpha+n}(\omega) = X_{\alpha(\omega)+n}(\omega)$ . The post- $\alpha$  field  $\mathcal{F}'_\alpha$  is the sub- $\sigma$ -field generated by the post- $\alpha$  process.

**Proposition 1.3.7.** *Let  $\{X_n\}_{n=0}^\infty$  be a stochastic process and let  $\alpha$  be a stopping time. Then  $\alpha \in \mathcal{F}_\alpha$  and  $X_\alpha \in \mathcal{F}_\alpha$ .*

**Proof.** For  $\alpha$  to be  $\mathcal{F}_\alpha$ -measurable, we need to show that  $\{\alpha = k\} \in \mathcal{F}_\alpha$ . This follows from (1.3.4) by taking  $\Lambda_n = \emptyset$  for  $n \neq k$  and  $\Lambda_k = \Omega$ .

That  $X_\alpha \in \mathcal{F}_\alpha$  follows from

$$\{\omega : X_\alpha(\omega) \in B\} = \bigcup_n \{\omega : \alpha(\omega) = n, X_n(\omega) \in B\} \in \mathcal{F}_\alpha$$

for any Borel set  $B \in \mathcal{B}$ .  $\square$

**Theorem 1.3.8.** *Let  $\{X_n\}_{n=0}^\infty$  be a Markov-process and  $\alpha$  is an a.e. finite stopping time, then for each  $M \in \mathcal{F}'_\alpha$ ,*

$$P(M|\mathcal{F}_\alpha) = P(M|\alpha, X_\alpha). \quad (1.3.5)$$

We call this property the *strong* Markov-property.

**Proof.** Note that the generating sets of  $\mathcal{F}'_\alpha$  are  $M = \bigcap_{j=1}^l X_{\alpha+j}^{-1}(B_j)$ ,  $B_j \in \mathcal{B}$ .

Let  $M_n = \bigcap_{j=1}^l X_{n+j}^{-1}(B_j) \in \mathcal{F}_{(n,\infty)}$ , We claim that

$$P(M|\alpha, X_\alpha) = \sum_{n=1}^{\infty} P(M_n|X_n)\chi_{\{\alpha=n\}}. \quad (1.3.6)$$

Indeed if we consider  $P(M_n|X_n) = \varphi_n(X_n)$ , then it is clear  $\sum_{n=1}^{\infty} \varphi_n(X_n)\chi_{\{\alpha=n\}}$  is measurable with respect to the  $\sigma$ -field generated by  $\alpha$  and  $X_\alpha$ . By making use of Theorem 1.3.5(b), we have

$$\begin{aligned} \int_{\{\alpha=m, X_\alpha \in B\}} \sum_{n=1}^{\infty} P(M_n|X_n)\chi_{\{\alpha=n\}} dP &= \int_{\{\alpha=m, X_m \in B\}} P(M_m|X_m) dP \\ &= \int_{\{\alpha=m, X_m \in B\}} P(M_m|\mathcal{F}_{[0,m]}) dP \\ &= P(\{\alpha = m, X_m \in B\} \cap M_m) \\ &= P(\{\alpha = m, X_\alpha \in B\} \cap M). \end{aligned}$$

(The last equality is due to  $M_m \cap \{\alpha = m\} = M \cap \{\alpha = m\}$ ). Hence the claim follows.

Now to prove the theorem, let  $\Lambda \in \mathcal{F}_\alpha$ ,  $\Lambda = \bigcup_{n=0}^{\infty} (\{\alpha = n\} \cap \Lambda_n)$ , then

$$\begin{aligned}
 P(\Lambda \cap M) &= \sum_{n=0}^{\infty} P(\{\alpha = n, \Lambda_n\} \cap M_n) \\
 &= \sum_{n=0}^{\infty} \int_{\{\alpha=n\} \cap \Lambda_n} P(M_n | \mathcal{F}_{[0,n]}) dP \\
 &= \sum_{n=0}^{\infty} \int_{\Lambda} P(M_n | X_n) \chi_{\{\alpha=n\}} dP \quad (\text{by Theorem 1.3.5(b)}) \\
 &= \int_{\Lambda} P(M_n | \alpha, X_\alpha) dP \quad (\text{by (1.3.6)}).
 \end{aligned}$$

The theorem follows from this.  $\square$

We remark that when  $\alpha$  is the constant  $n$ , then we can omit the  $\alpha$  in (1.3.5) and it reduces to the Markov property as in Theorem 1.3.5. Also if the process is homogeneous (i.e., invariant on the time  $n$ ), then we can omit the  $\alpha$  there. It is because in (1.3.6), the right side can be represented as  $\sum_{n=1}^{\infty} \varphi(X_n) \chi_{\{\alpha=n\}}$  (instead of  $\varphi_n(X_n)$ ) which is  $\mathcal{F}_\alpha$ -measurable. In this case we can rewrite (1.3.5) as

$$P(X_{\alpha+1} \in B | \mathcal{F}_\alpha) = P(X_{\alpha+1} \in B | X_\alpha) \quad \forall B \in \mathcal{B},$$

a direct analog of the definition of Markov property.

There is a constructive way to obtain Markov processes. For a Markov chain  $\{X_n\}_{n=0}^{\infty}$ , we mean a stochastic process that has a state space  $S = \{a_1, a_2, \dots, a_N\}$  (finite or countable) and a transition matrix

$$P = \begin{pmatrix} p_{11} & \cdots & p_{1N} \\ \vdots & \cdots & \vdots \\ p_{N1} & \cdots & p_{NN} \end{pmatrix}$$

where  $p_{ij} \geq 0$  and the row sum is 1; the  $p_{ij}$  is the probability from  $i$  to  $j$ . Suppose the process starts at  $X_0$  with initial distribution  $\mu = (\mu_1, \dots, \mu_N)$ , let  $X_n$  denote the location of the chain at the  $n$ -th time according to the transition matrix  $P$ , then  $\{X_n\}_{n=0}^\infty$  satisfies the Markov property:

$$P(X_{n+1} = x_{n+1} | X_0 = x_0, \dots, X_n = x_n) = P(X_{n+1} = x_{n+1} | X_n = x_n) = p_{ij}.$$

Also it follows that

$$\begin{aligned} & P(X_0 = x_0, X_1 = x_1, \dots, X_n = x_n) \\ &= P(X_0 = x_0)P(X_1 = x_1 | X_0 = x_0) \cdots P(X_n = x_n | X_{n-1} = x_{n-1}) \\ &= \mu_{x_0} p_{x_0 x_1} \cdots p_{x_{n-1} x_n}. \end{aligned}$$

More generally, we consider the state space to be  $\mathbb{R}$ . Let  $\mu : \mathbb{R} \times \mathcal{B} \rightarrow [0, 1]$  satisfies

- (a) for each  $x$ ,  $\mu(x, \cdot)$  is a probability measure;
- (b) for each  $B$ ,  $\mu(\cdot, B)$  is a Borel measurable function.

Let  $\{X_n\}_{n=0}^\infty$  be a sequence of r.v. with finite dimensional joint distributions  $\mu^{(n)}$  for  $X_0, \dots, X_n$  given by

$$\begin{aligned} P\left(\bigcap_{j=0}^n \{X_j \in B_j\}\right) &= \mu^{(n)}(B_0 \times \cdots \times B_n) \\ &:= \int \cdots \int_{B_0 \times \cdots \times B_n} \mu_0(dx_0) \mu(x_0, dx_1) \cdots \mu(x_{n-1}, dx_n). \end{aligned}$$

where  $\mu_0$  is the distribution function of  $X_0$ .

It is direct to check from definition that

$$P(X_{n+1} \in B | X_n) = \mu(X_n, B),$$

i.e.,

$$P(X_{n+1} \in B | X_n = x) = \mu(x, B).$$

Hence  $\mu(x, B)$  represents the probability that in the  $(n + 1)$ -step the chain is in  $B$ , starting at  $x$  in the  $n$ -th step. To see that  $\{X_n\}_{n=0}^\infty$  satisfies the Markov property, we let  $\Lambda = \bigcap_{j=0}^n \{X_j \in B_j\}$ , then

$$\begin{aligned} \int_{\Lambda} P(X_{n+1} \in B | X_n) dP &= \int \cdots \int_{B_0 \times \cdots \times B_n} \mu(x_n, B) d\mu^{(n)}(x_0, \cdots, x_n) \\ &= \int \cdots \int_{B_0 \times \cdots \times B_n \times B} \mu_0(dx_0) \prod_{j=1}^{n+1} \mu(x_{j-1}, dx_j) \\ &= P(\Lambda \cap \{X_{n+1} \in B\}). \end{aligned}$$

This implies

$$P(X_{n+1} \in B | X_n) = P(X_{n+1} \in B | X_1, \cdots, X_n)$$

and the Markov property follows.

We call the above  $\{X_n\}_{n=0}^\infty$  a *stationary* (or *homogeneous*) Markov process and  $\mu(x, B)$  the transition probability.

**Exercises**

1. Let  $\{X_n\}_{n=0}^{\infty}$  be a Markov process. Let  $f$  be a one-to-one Borel measurable function on  $\mathbb{R}$  and let  $Y_n = f(X_n)$ . Show that  $\{Y_n\}_{n=0}^{\infty}$  is also a Markov process (with respect to the fields generated by  $f(X_n)$ ); but the conclusion does not hold if we do not assume  $f$  is one-to-one.
2. Prove the strong Markov property in the form of Theorem 1.3.5(c).
3. If  $\alpha_1$  and  $\alpha_2$  are both stopping times, so are  $\alpha_1 \wedge \alpha_2$ ,  $\alpha_1 \vee \alpha_2$  and  $\alpha_1 + \alpha_2$ . However  $\alpha_1 - \alpha_2$  is not necessarily a stopping time.
4. Let  $\{X_n\}_{n=1}^{\infty}$  be a sequence of i.i.d.r.v. Let  $\{\alpha_k\}_{k=1}^{\infty}$  be a sequence of strictly increasing finite stopping times. Then  $\{X_{\alpha_k+1}\}_{k=1}^{\infty}$  is also a sequence of i.i.d.r.v. (This is the gambling-system theorem given by Doob).
5. A sequence  $\{X_n\}_{n=0}^{\infty}$  is a Markov chain of *second order* if

$$P(X_{n+1} = j | X_0 = i_0, \dots, X_n = i_n) = P(X_{n+1} = j | X_{n-1} = i_{n-1}, X_n = i_n).$$

Show that nothing really new is involved because the sequence  $(X_n, X_{n+1})$  is a Markov chain.

6. Let  $\mu^{(n)}(x, B)$  be the  $n$ -step transition probability in the stationary Markov process. Prove the Chapman-Kolmogorov equation

$$\mu^{(m+n)}(x, B) = \int_{\mathbb{R}} \mu^{(m)}(x, dy) \mu^{(n)}(y, B) \quad \forall m, n \in \mathbb{N}.$$

## 1.4 Martingales

We first consider a simple example in analysis. Let  $f$  be an integrable function on  $[0, 1]$ , let  $\mathcal{P}_n = \{0 = \frac{1}{2^n} \leq \dots \leq \frac{k}{2^n} \dots \leq 1\}$  be a partition of  $[0, 1]$  and let  $I_{n,k} = [\frac{k}{2^n}, \frac{k+1}{2^n})$ . We define the average function  $f_n$  of  $f$  on the partition  $\mathcal{P}_n$ :

$$f_n(x) = \sum_{k=0}^{2^n-1} a_{n,k} \chi_{I_{n,k}}, \quad x \in I_{n,k}. \quad (1.4.1)$$

where  $a_{n,k} = \frac{1}{|I_{n,k}|} \int_{I_{n,k}} f(x) dx$ . Then  $\{f_n\}_n$  converges to  $f$  in  $L^1$ . Moreover  $\{f_n\}_n$  has the following consistency property: for  $m > n$

$$f_n(x) = \frac{1}{|I_{n,k}|} \int_{I_{n,k}} f_m(y) dy \quad x \in I_{n,k}. \quad (1.4.2)$$

This property has been reformulated by Doob in the more general probability setting.

**Definition 1.4.1.** Let  $\{(X_n, \mathcal{F}_n)\}_{n=1}^\infty$  be a sequence of r.v. such that  $X_n \in \mathcal{F}_n$ . It is called a martingale if

- (a)  $\mathcal{F}_n \subset \mathcal{F}_{n+1}$ ;
- (b)  $E(|X_n|) < \infty$ ;
- (c)  $X_n = E(X_{n+1} | \mathcal{F}_n)$ .

It is called a supermartingale (or submartingale) if  $\geq$  (or  $\leq$  respectively) in (c) holds. We will call  $\{X_n\}_n$  a s-martingale if it is any one of the three cases.

Condition (c) can be strengthened as  $X_n = E(X_m | \mathcal{F}_n)$  for  $m > n$ . It follows from

$$E(X_m | \mathcal{F}_n) = E(E(X_m | \mathcal{F}_{m-1}) | \mathcal{F}_n) = E(X_{m-1} | \mathcal{F}_n) = \dots = E(X_n | \mathcal{F}_n) = X_n .$$

Martingale has its intuitive background in gambling. If  $X_n$  is interpreted as the gambler's capital at time  $n$ , then the defining property says that his

expected capital after next game, played with the knowledge of the entire past and present, is exactly equal to his current capital. In other words, his expected gain is zero, and is in this sense the game is said to be “fair”. The supermartingale and submartingale can be interpreted similarly.

**Example 1.** As a direct analog of the above function case, we let  $X$  be an integrable r.v. and let  $\{\mathcal{F}_n\}_{n=1}^\infty$  be an increasing sequence of sub- $\sigma$ -fields (e.g., take  $\mathcal{F}_n$  to be a partition). Let  $X_n = E(X|\mathcal{F}_n)$ . Then  $\{X_n\}_{n=1}^\infty$  is a martingale. Indeed we see that

$$E(|X_n|) = E(|E(X|\mathcal{F}_n)|) \leq E(E(|X||\mathcal{F}_n)) = E(|X|) < \infty$$

and (b) follows. For (c), we observe that

$$E(X_{n+1}|\mathcal{F}_n) = E(E(X|\mathcal{F}_{n+1})|\mathcal{F}_n) = E(X|\mathcal{F}_n) = X_n.$$

**Example 2.** Let  $\{X_n\}_{n=1}^\infty$  be a sequence of independent integrable r.v. with mean zero. Let  $S_n = \sum_{j=1}^n X_j$  and  $\mathcal{F}_n = \mathcal{F}(X_1, \dots, X_n)$ . Then

$$\begin{aligned} E(S_{n+1}|\mathcal{F}_n) &= E(S_n + X_{n+1}|\mathcal{F}_n) \\ &= S_n + E(X_{n+1}|\mathcal{F}_n) \\ &= S_n + E(X_{n+1}) \\ &= S_n. \end{aligned}$$

Hence  $\{(S_n, \mathcal{F}_n)\}$  is a martingale.

**Proposition 1.4.2.** *If  $\{(X_n, \mathcal{F}_n)\}_{n=1}^\infty$  is a submartingale, and  $\varphi$  is increasing and convex in  $\mathbb{R}$ . If  $\{\varphi(X_n)\}$  is integrable, then  $\{(\varphi(X_n), \mathcal{F}_n)\}$  is also a submartingale.*

**Proof.** Since  $X_n \leq E(X_{n+1}|\mathcal{F}_n)$ , by the property of  $\varphi$ , we have

$$\varphi(X_n) \leq \varphi(E(X_{n+1}|\mathcal{F}_n)) \leq E(\varphi(X_{n+1})|\mathcal{F}_n) \quad \square$$



It follows that if  $\{X_n\}_{n=0}^\infty$  is a martingale (or submartingale), then  $\{|X_n|^p\}_{n=0}^\infty$ ,  $p \geq 1$  (provided that  $X_n \in L^p$ ) and  $\{X_n^+\}_{n=0}^\infty$  are submartingales. Also if  $\{X_n\}$  is a supermartingale, so does  $\{X_n \wedge a\}_n$  for any  $a \in \mathbb{R}$ .

**Theorem 1.4.3.** (*Doob's decomposition Theorem*) For any submartingale  $\{(X_n, \mathcal{F}_n)\}_{n=1}^\infty$ ,  $X_n$  can be decomposed as

$$X_n = Y_n + Z_n$$

where  $\{(Y_n, \mathcal{F}_n)\}_{n=1}^\infty$  is a martingale and  $\{Z_n\}$  is a non-negative increasing process.

**Proof.** We define the difference r.v.

$$D_1 = X_1, \quad D_j = X_j - X_{j-1}, \quad j \geq 2.$$

Then  $X_n = \sum_{j=1}^n D_j$ , and the defining relation of submartingale yields

$$E(D_j | \mathcal{F}_{j-1}) \geq 0, \quad j \geq 2. \tag{1.4.3}$$

We consider yet another difference

$$S_1 = D_1, \quad S_j = D_j - E(D_j | \mathcal{F}_{j-1}),$$

and let

$$Y_n = \sum_{j=1}^n S_j, \quad Z_n = \sum_{j=1}^n E(D_j | \mathcal{F}_{j-1}).$$

It is clear that  $X_n = Y_n + Z_n$ ,  $X_1 = Y_1$ ,  $Z_1 = 0$  and  $\{Z_n\}_{n=1}^\infty$  is a non-negative increasing process (by (1.4.3)). On the other hand, note that  $E(S_j | \mathcal{F}_{j-1}) = 0$ , it follows that

$$E(Y_n | \mathcal{F}_{n-1}) = \sum_{j=1}^{n-1} S_j = Y_{n-1}$$

and hence a martingale.  $\square$

For an increasing family of sub- $\sigma$ -fields  $\{\mathcal{F}_n\}_{n=1}^\infty$ , let  $\mathcal{F}_\infty = \bigcup_{n=1}^\infty \mathcal{F}_n$  and let  $\alpha$  be a stopping time with respect to  $\{\mathcal{F}_n\}_{n=1}^\infty$ , i.e.,

$$\alpha : \Omega \rightarrow \mathbb{N} \cup \{\infty\} \quad \text{such that} \quad \{\alpha = n\} \in \mathcal{F}_n$$

As in last section, the pre- $\alpha$  field  $\mathcal{F}_\alpha$  is the family of sets

$$\Lambda = \bigcup_n (\{\alpha = n\} \cap \Lambda_n), \quad \Lambda_n \in \mathcal{F}_n.$$

The following theorems aim at replacing the constant time of a martingale by a stopping time.

**Theorem 1.4.4.** *Let  $Y$  be integrable r.v. and let  $X_n = E(Y|\mathcal{F}_n)$  where  $\mathcal{F}_n$  is an increasing family of sub- $\sigma$ -fields (it is a martingale). Then for any stopping time  $\alpha$ , we have  $X_\alpha = E(Y|\mathcal{F}_\alpha)$ .*

*Moreover if  $\beta$  is also a stopping time and  $\alpha \leq \beta$ , then  $\{(X_\alpha, \mathcal{F}_\alpha), (X_\beta, \mathcal{F}_\beta)\}$  is a two term martingale (i.e.,  $X_\alpha = E(X_\beta|\mathcal{F}_\alpha)$ ).*

**Proof.** Note that  $X_\alpha \in \mathcal{F}_\alpha$ . We claim that it is also integrable. Indeed as

$$|X_n| = |E(Y|\mathcal{F}_n)| \leq E(|Y||\mathcal{F}_n),$$

we have

$$\int_\Omega |X_\alpha| dP = \sum_n \int_{\{\alpha=n\}} |X_n| dP \leq \sum_n \int_{\{\alpha=n\}} |Y| dP = \int_\Omega |Y| dP < \infty.$$

Now if  $\Lambda \in \mathcal{F}_\alpha$ , let  $\Lambda_n = \Lambda \cap \{\alpha = n\}$ , then

$$\int_\Lambda X_\alpha dP = \sum_n \int_{\Lambda_n} X_n dP = \sum_n \int_{\Lambda_n} Y dP = \int_\Lambda Y dP.$$

Hence  $X_\alpha = E(Y|\mathcal{F}_\alpha)$ .

For the last statement, note that  $\mathcal{F}_\alpha \subset \mathcal{F}_\beta$ , hence

$$E(X_\beta|\mathcal{F}_\alpha) = E(E(Y|\mathcal{F}_\beta)|\mathcal{F}_\alpha) = X_\alpha. \quad \square$$

**Corollary 1.4.5.** *Under the above assumption and suppose  $\{\alpha_i\}_{i=1}^{\infty}$  is an increasing sequence of stopping times. If  $\{(X_n, \mathcal{F}_n)\}_n$  is an  $s$ -martingale, then  $\{(X_{\alpha_i}, \mathcal{F}_{\alpha_i})\}_i$  is an  $s$ -martingale.*

Unlike Theorem 1.4.4, in the following theorem, we do not assume that the  $\{X_n\}$  is the conditional expectation of an integrable  $Y$ .

**Theorem 1.4.6.** *Let  $\{(X_n, \mathcal{F}_n)\}_n$  be a  $s$ -martingale. Let  $\alpha \leq \beta$  be two bounded stopping times, then  $\{(X_\alpha, \mathcal{F}_\alpha), (X_\beta, \mathcal{F}_\beta)\}$  is also an  $s$ -martingale (of the same type).*

**Proof.** We prove the theorem for supermartingale. For submartingale, we consider  $\{-X_n\}$  instead.

Let  $\Lambda \in \mathcal{F}_\alpha$ , and let  $\Lambda_j = \Lambda \cap \{\alpha = j\}$  ( $\in \mathcal{F}_j$ ). Then for  $k \geq j$ ,  $\Lambda_j \cap \{\beta > k\} \in \mathcal{F}_k$ , hence

$$\begin{aligned} \int_{\Lambda_j \cap \{\beta \geq k\}} X_k dP &= \int_{\Lambda_j \cap \{\beta > k\}} X_k dP + \int_{\Lambda_j \cap \{\beta = k\}} X_k dP \\ &\geq \int_{\Lambda_j \cap \{\beta > k\}} X_{k+1} dP + \int_{\Lambda_j \cap \{\beta = k\}} X_k dP \end{aligned}$$

i.e.,

$$\int_{\Lambda_j \cap \{\beta \geq k\}} X_k dP - \int_{\Lambda_j \cap \{\beta \geq k+1\}} X_{k+1} dP \geq \int_{\Lambda_j \cap \{\beta = k\}} X_\beta dP$$

Summing over  $k$ ,  $j \leq k \leq m$ , where  $m$  is the upper bound of  $\beta$ , then

$$\int_{\Lambda_j \cap \{\beta \geq j\}} X_\alpha dP - \int_{\Lambda_j \cap \{\beta \geq m+1\}} X_{m+1} dP \geq \int_{\Lambda_j \cap \{j \leq \beta \leq m\}} X_\beta dP$$

Hence

$$\int_{\Lambda_j} X_\alpha dP \geq \int_{\Lambda_j} X_\beta dP$$

Summing over  $1 \leq j \leq m$ , we have

$$\int_{\Lambda} X_\alpha dP \geq \int_{\Lambda} X_\beta dP \quad \forall \Lambda \in \mathcal{F}_\alpha. \quad \square$$

**Corollary 1.4.7.** *If  $\{(X_n, \mathcal{F}_n)\}$  is a martingale or a supermartingale, then the same is for  $\{(X_{\alpha \wedge n}, \mathcal{F}_{\alpha \wedge n})\}$  for any stopping time  $\alpha$ .*

The theorem still holds if  $\alpha, \beta$  are unbounded. For this we need to associate a random variable  $X_\infty$  at  $\infty$ .

**Theorem 1.4.8.** *Assume  $\lim_{n \rightarrow \infty} X_n = X_\infty$  exists and is integrable. Let  $\alpha, \beta$  be two arbitrary stopping times. Then Theorem 1.4.6 still hold if  $\{(X_n, \mathcal{F}_n)\}_{n \in \mathbb{N}_\infty}$  is a supermartingale.*

Proof. We first assume that  $X_n \geq 0$  and  $X_\infty = 0$ . Then  $X_\alpha \leq \liminf_{n \rightarrow \infty} X_{\alpha \wedge n}$ , and hence  $X_\alpha$  is integrable by Fatou's lemma. The same is for  $X_\beta$ .

From the proof of Theorem 1.4.6, we can conclude that for any  $m$

$$\int_{\Lambda \cap \{\alpha=j\}} X_\alpha dP \geq \int_{\Lambda \cap \{\alpha=j\} \cap \{\beta \leq m\}} X_\beta dP .$$

By letting  $m \rightarrow \infty$  and summing over all  $j$ , we have

$$\int_{\Lambda \cap \{\alpha < \infty\}} X_\alpha dP \geq \int_{\Lambda \cap \{\beta < \infty\}} X_\beta dP .$$

In addition we have  $X_\alpha = X_\infty = 0$  on  $\{\alpha = \infty\}$ , and  $X_\beta = X_\infty = 0$  on  $\{\beta = \infty\}$ , We conclude that

$$\int_{\Lambda} X_\alpha dP = \int_{\Lambda} X_\beta dP$$

and hence  $\{(X_\alpha, \mathcal{F}_\alpha), (X_\beta, \mathcal{F}_\beta)\}$  is a supermartingale.

For the general case we let

$$X'_n = E(X_\infty | \mathcal{F}_n), \quad X''_n = X_n - X'_n.$$

Then  $\{X'_n\}$  is a martingale, and  $X_n \geq X'_n$  by the defining property of supermartingale apply to  $X_n$  and  $X_\infty$ . We can apply the above proved case to  $X''_n$ , and conclude that  $\{X_n\}$  is a supermartingale.  $\square$

The above theorems are referred as Doob's *optional sampling theorems*. In terms of gambling, one would hope to devise a strategy to gain advantage of the outcome, but the theorems say that such a strategy does not exist, at least mathematically. The reader can refer to [1, p.327](and the exercises there) for a discussion of the *gambler's ruin problem*.

We use the above stopping time consideration to prove a useful inequality for sub-martingales.

**Theorem 1.4.9.** *If  $\{(X_j, \mathcal{F}_j)\}_{j=1}^n$  is a submartingale, then for any real  $\lambda$ , we have*

$$\lambda P(\max_{1 \leq j \leq n} X_j > \lambda) \leq \int_{\{\max_{1 \leq j \leq n} X_j > \lambda\}} X_n dP \leq E(X_n^+).$$

**Proof.** Let  $\alpha$  be the first  $j$  such that  $X_j \geq \lambda$  if such  $1 \leq j \leq n$  exists, otherwise let  $\alpha = n$ . It is clear that  $\alpha$  is a stopping time, and hence  $\{X_\alpha, X_n\}$  is a submartingale (Theorem 1.4.6). If we write

$$M = \{\max_{1 \leq j \leq n} X_j \geq \lambda\},$$

then  $M \in \mathcal{F}_\alpha$  and  $X_\alpha \geq \lambda$  on  $M$ , hence the first inequality follows from

$$\lambda P(M) \leq \int_M X_\alpha dP \leq \int_M X_n dP.$$

The second inequality is clear.  $\square$ .

**Corollary 1.4.10.** *If  $\{(X_n, \mathcal{F}_n)\}_{n=1}^\infty$  is a martingale, then for any  $\lambda > 0$ , we have*

$$P(\max_{1 \leq j \leq n} |X_j| > \lambda) \leq \int_{\{\max_{1 \leq j \leq n} |X_j| > \lambda\}} |X_n| dP \leq \frac{1}{\lambda} E(|X_n|).$$

*In addition if  $E(|X_n|^2) < \infty$ , then we also have*

$$P(\max_{1 \leq j \leq n} |X_j| > \lambda) \leq \frac{1}{\lambda^2} E(|X_n|^2).$$

For a sequence of independent r.v.  $\{X_n\}_{n=0}^\infty$  with zero mean and finite variance, we let  $S_n = \sum_{j=1}^n X_j$ . It is well known (Kolmogorov's inequality [1, p. 116]) that for any  $\lambda > 0$ ,

$$P(\max_{1 \leq j \leq n} |S_j| > \lambda) \leq \frac{1}{\lambda^2} E(|S_n|^2).$$

We see that the inequality follows directly from the above corollary.

To conclude this section, we prove a deep theorem on the convergence of the  $\{X_n\}_n$ , which is also due to Doob. It involves an ingenious method in the proof.

**Theorem 1.4.11.** *If  $\{(X_n, \mathcal{F}_n)\}_{n=0}^\infty$  is an  $L^1$ -bounded submartingale, then  $\{X_n\}_{n=0}^\infty$  converges a.e. to a finite limit.*

**Proof.** First we define, for any pair of rationals  $a, b$ , let

$$\Lambda_{[a,b]} = \{\omega : \liminf_{n \rightarrow \infty} X_n(\omega) < a < b < \limsup_{n \rightarrow \infty} X_n(\omega)\} \quad (1.4.4)$$

We show that  $\Lambda_{[a,b]}$  is a zero set for any  $a, b \in \mathbb{Q}$ . It follows that

$$\{\omega : \liminf_{n \rightarrow \infty} X_n(\omega) < \limsup_{n \rightarrow \infty} X_n(\omega)\} = \bigcup_{a, b \in \mathbb{Q}, a < b} \Lambda_{[a,b]}$$

is a zero set. Note that  $\liminf_{n \rightarrow \infty} X_n$  is finite almost everywhere (by Fatou lemma and the  $L^1$ -boundedness assumption,  $E(\liminf |X_n|) \leq \liminf E(|X_n|) < \infty$ ), hence the theorem follows.

It remains to prove (1.4.4). We first introduce some notations. Let  $\{x_1, \dots, x_n\}$  be a numerical sequence, for  $a < b$ , let

$$\alpha_1 = \min\{j : 1 \leq j \leq n, x_j \leq a\},$$


$$\alpha_2 = \min\{j : \alpha_1 < j \leq n, x_j \geq b\}.$$

Inductively we define

$$\alpha_{2k-1} = \min\{j : \alpha_{2k-2} < j \leq n, x_j \leq a\},$$

$$\alpha_{2k} = \min\{j : \alpha_{2k-1} < j \leq n, x_j \geq b\}.$$

Let  $\alpha_l$  be the last one defined. We can think of connecting the consecutive  $x_i$  by line segments, Let  $\nu$  be the number of times the line segments comes from  $\leq a$  to  $\geq b$ , i.e., the number of upcrossing through the interval  $[a, b]$ , it is seen that  $\nu = \lfloor l/2 \rfloor$ .



**Lemma 1.4.12.** Let  $\{(X_j, \mathcal{F}_j)\}_{j=1}^n$  be a submartingale and assume that  $X_j \geq 0$ . Let  $\nu_{[0,b]}^{(n)}$  be the r.v. of the number of upcrossing of  $[0, b]$  by the sample sequence  $\{X_j(\omega) : 1 \leq j \leq n\}$ . Then

$$E(\nu_{[0,b]}^{(n)}) \leq \frac{E(X_n - X_1)}{b}.$$

**Proof.** For convenience, we let  $\alpha_0 = 1$  and  $\alpha_{l+1} = \alpha_{l+2} = \dots = \alpha_n = n$ . Then we have a sequence of stopping times with

$$1 = \alpha_0 \leq \alpha_1 < \dots < \alpha_l \leq \alpha_{l+1} \dots \leq \alpha_n = n.$$

We write

$$X_n - X_1 = X_{\alpha_n} - X_{\alpha_0} = \sum_{j=1}^{n-1} (X_{\alpha_{j+1}} - X_{\alpha_j}) = \left( \sum_{j \text{ odd}} + \sum_{j \text{ even}} \right) (X_{\alpha_{j+1}} - X_{\alpha_j}).$$

It follows that

$$\sum_{j \text{ odd}} (X_{\alpha_{j+1}}(\omega) - X_{\alpha_j}(\omega)) \geq [l(\omega)/2] \cdot b = \nu_{[0,b]}^{(n)}(\omega) \cdot b.$$

On the other hand by Theorem 1.4.6,  $\{X_{\alpha_j} : 0 \leq j \leq n\}$  is a submartingale, so that for each  $0 \leq j \leq n-1$ ,  $E(X_{\alpha_{j+1}} - X_{\alpha_j}) \geq 0$ . Consequently

$$E\left(\sum_{j \text{ even}} (X_{\alpha_{j+1}} - X_{\alpha_j})\right) \geq 0.$$

Therefore  $E(X_n - X_1) \geq E(\nu_{[0,b]}^{(n)}) \cdot b$  which yields the lemma.  $\square$ .

Now to complete the proof of (1.4.4), we consider the upcrossing on any  $[a,b]$ . We replace the r.v. in the lemma by  $(X_n - a)^+$ . The sequence  $\{(X_n - a)^+\}_n$  is still submartingale and by the lemma,

$$E(\nu_{[a,b]}^{(n)}) \leq \frac{E(X_n - a)^+ - E(X_1 - a)^+}{b - a} \leq \frac{E(X_n^+) + |a|}{b - a}.$$

Let  $\nu_{[a,b]} = \lim_{n \rightarrow \infty} \nu_{[a,b]}^{(n)}$ . The  $L^1$ -boundedness of  $\{X_n\}_n$  implies that  $E(\nu_{[a,b]}) < \infty$ . Hence  $\nu_{[a,b]}$  is finite with probability 1. Note that

$$\begin{aligned} \Lambda_{[a,b]} &= \{\omega : \liminf_{n \rightarrow \infty} X_n(\omega) \leq a < b \leq \limsup_{n \rightarrow \infty} X_n(\omega)\} \\ &\subseteq \{\omega : \nu_{[a,b]}(\omega) = \infty\}, \end{aligned}$$

hence  $\Lambda_{[a,b]}$  is a zero set and (1.4.4) follows. This completes the proof of the theorem.  $\square$

**Corollary 1.4.13.** *Every uniformly bounded  $s$ -martingale converges a.e. Also every positive supermartingale and every negative submartingale converges a.e.*

**Proof.** The first statement follows directly from Theorem 1.4.8 and that  $\{X_n\}$  is a submartingale if and only if  $\{-X_n\}$  is a supermartingale.

For the second part we use Doob's decomposition theorem (Theorem 1.4.3). Let  $\{X_n\}_n$  be a positive supermartingale, then  $X_n = Y_n - Z_n$  where  $\{Y_n\}$  is a



martingale and  $Z_n \geq 0$ ,  $\{Z_n\} \nearrow$ . Since  $X_n \geq 0$ , it follows that  $0 \leq Z_n \leq Y_n$ . Let  $Z_\infty = \lim_{n \rightarrow \infty} Z_n$ . It is finite a.e. because

$$E(Z_\infty) = \lim_{n \rightarrow \infty} E(Z_n) \leq E(Y_1) < \infty.$$

Also since  $\{X_n\}_n$  is a supermartingale,

$$E(Y_n) = E(X_n) + E(Z_n) \leq E(X_1) + E(Z_\infty).$$

This implies  $\{Y_n\}_n$  is  $L^1$ -uniformly bounded and  $\{Y_n\}_n$  converges to a finite limit a.e. (Theorem 1.4.8). The same holds for  $\{X_n\}_n$ .  $\square$

Recall that a sequence of r.v.  $\{X_n\}_{n=1}^\infty$  is called uniformly integrable if

$$\lim_{k \rightarrow \infty} \int_{|X_n| \geq k} |X_n| dP = 0 \quad \text{uniformly on } n.$$

It is clear that it implies that  $\{X_n\}_{n=1}^\infty$  is  $L^1$ -bounded. Also, if  $X_n \rightarrow X$  a.e., then the uniform boundedness implies that  $X_n \rightarrow X$  in  $L^1$  ([1, p.96-97]).

**Corollary 1.4.14.** *If  $\{(X_n, \mathcal{F}_n)\}_{n=1}^\infty$  is a submartingale and is uniformly integrable, then  $X_\infty = \lim_{n \rightarrow \infty} X_n$  a.e. and in  $L^1$ .*

**Remark.** Theorem 1.4.11 and Corollary 1.4.14 are more or less that the converse of Example 1. However for Example 2, the sum  $\{S_n\}_{n=1}^\infty$  of i.i.d.r.v.  $\{X_n\}_{n=0}^\infty$  with zero mean forms a martingale, but does not converge; it is because the  $L^1$ -bounded condition is not satisfied. In fact, we can show that

$$\lim_{n \rightarrow \infty} E\left(\frac{|S_n|}{\sqrt{n}}\right) = \sqrt{\frac{2}{\pi}}\sigma$$

where  $\sigma$  is the variance of  $X_n$ . For more detail, the reader can refer to [1, Chapter 5, 6] for the law of large number and the central limit theorem for  $\{S_n\}_{n=1}^\infty$ .

**Exercises**

1. Suppose  $\{(X_n^{(k)}, \mathcal{F}_n)\}_n$ ,  $k = 1, 2$  are two martingales,  $\alpha$  is a finite stopping time and  $X_\alpha^{(1)} = X_\alpha^{(2)}$ . Define  $X_n = X_n^{(1)}\chi_{\{n \leq \alpha\}} + X_n^{(2)}\chi_{\{n > \alpha\}}$ . Show that  $\{(X_n, \mathcal{F}_n)\}_n$  is a martingale.

2 If  $\{(X_n, \mathcal{F}_n)\}_n$ ,  $\{(Y_n, \mathcal{F}_n)\}_n$  are martingales, then  $\{(X_n + Y_n, \mathcal{F}_n)\}_n$  is again a martingale. However it may happen that  $\{X_n\}_n$ ,  $\{Y_n\}_n$  are martingales, but  $\{X_n + Y_n\}_n$  is not a martingale. (Note the the  $\sigma$ -field generated by  $X_n + Y_n$  may not have the same  $\sigma$ -field  $\mathcal{F}_n$ .)

3 Prove that for any  $L^1$ -bounded s-martingale  $\{(X_n, \mathcal{F}_n)\}_n$ , and for any  $\alpha$  stopping time, then  $E(|X_\alpha|) < \infty$ .

4. If  $X$  is an integrable r.v., then the collection of r.v.,  $D(X|\mathcal{G})$  with  $\mathcal{G}$  ranging over all Borel subfields of  $\mathcal{F}$ , is uniformly integrable.

5. Find an example of a positive martingale that is not uniformly integrable.

6. Find an example of a martingale  $\{X_n\}_n$  such that  $X_n \rightarrow -\infty$ . This implies that in a fair game one player may lose an arbitrary large amount if he stays on long enough. (Hint: Try sums of independent but not identically distributed r.v. with mean 0.)

7. If  $\{X_n\}_n$  is a uniformly integrable submartingale, then for any stopping time  $\alpha$ ,  $\{X_{\alpha \wedge n}\}_n$  is again a uniformly integrable submartingale and

$$E(X_1) \leq E(X_\alpha) \leq \sup_n E(X_n).$$

8 Prove that for any s-martingale, we have for each  $\lambda > 0$ ,

$$\lambda P(\sup_n |X_n| \geq \lambda) \leq 3 \sup_n E(|X_n|).$$

For a martingale or a positive or nonnegative s-martingale the constant 3 may be replaced by 1.

**9.** Let  $\{X_n\}_n$  be a positive supermartingale. Then for almost every  $\omega$ ,  $X_k(\omega) = 0$  implies  $X_n(\omega) = 0$  for all  $n \geq k$ .

**10.** Every  $L^1$ -martingale is the difference of two positive  $L^1$ -bounded martingales. (Hint, take one of them to be  $\lim_{k \rightarrow \infty} E(X_k^+ | \mathcal{F}_n)$ ).

**11.** If  $\{X_n\}$  is a martingale or positive submartingale such that  $\sup_n E(X_n^2) \leq \infty$ , then  $\{X_n\}_n$  converges in  $L^2$  as well as a.e.

**12.** Show that if  $\{(X_n, \mathcal{F}_n)\}_n$  is a submartingate,  $X_n \geq 0$ , then for  $p > 1$ ,

$$\| \max_{\{1 \leq k \leq n\}} \|_p \leq \frac{p}{p-1} \|X_n\|_p .$$

(Hint: Show that for  $Y \geq 0$ ,  $E(Y^p) = p \int_0^\infty \lambda^{p-1} P(Y \geq \lambda) d\lambda$ .)



# Chapter 2

## Brownian Motion

### 2.1 Continuous time stochastic processes

We call a family of random variables  $\{X_t\}_{t \geq 0}$  on  $(\Omega, \mathcal{F}, P)$  a *continuous time* stochastic process. For each  $\omega \in \Omega$ ,  $X(\cdot, \omega) = X_{(\cdot)}(\omega)$  is called a sample path. Usually we treat  $X(\cdot, \omega) = \omega(t)$  (this can be justified).

There are two most important classes of continuous time stochastic processes. The first one is the **Poisson process**  $\{N_t\}_{t \geq 0}$ , the number of arrivals in time  $[0, t]$  according to an arrival rate  $\lambda$  per unit time.




Figure 2.1:

Recall that a Poisson random variable  $X$  with rate  $\lambda$  has distribution

$$P(X = k) = e^{-\lambda} \frac{\lambda^k}{k!}, \quad k = 0, 1, 2, \dots$$

Hence  $N_t$  has distribution

$$P(N_t = k) = e^{-\lambda t} \frac{(\lambda t)^k}{k!}, \quad k = 0, 1, 2, \dots$$

A Poisson process is characterized by

1.  $N_0 = 0$ ;
2. Independent increment: for  $0 < t_1 < t_2 < \dots < t_n$ ,

$$N_{t_1}, N_{t_2} - N_{t_1}, N_{t_3} - N_{t_2}, \dots, N_{t_n} - N_{t_{n-1}}$$

are independent.

3. Poisson increment: for  $t > s$ ,  $N_t - N_s \sim N_{(t-s)}$ , i.e., it has a Poisson distribution with rate  $\lambda(t - s)$ .

The next one is the **Brownian motion**  $\{B_t\}_{t \geq 0}$ . It is also called a Wiener process due to the pioneer work of Wiener in the 20's. Recall that a one dimension normal distribution  $N(\mu, \sigma^2)$  has density function

$$\frac{1}{\sqrt{2\pi}\sigma} e^{-\frac{(x-\mu)^2}{2\sigma^2}}, \quad x \in \mathbb{R}$$

and  $N(0, 1)$  is called the standard normal distribution. The Brownian motion is defined by

1.  $B_0 = 0$ ;
2. Independent increment: for  $0 < t_1 < t_2 < \dots < t_n$ ,

$$B_{t_1}, B_{t_2} - B_{t_1}, B_{t_3} - B_{t_2}, \dots, B_{t_n} - B_{t_{n-1}}$$

are independent;




Figure 2.2:

3. Normal increment: for  $t > s$ ,  $B_t - B_s$ , has normal distribution  $N(0, t - s)$ .

We will see in the next section that almost all sample paths are continuous, but not differentiable anywhere. We can also define in the same way the higher dimensional Brownian motion, i.e.,  $\{B_t\}_{t \geq 0}$  has range in  $\mathbb{R}^d$ ; the corresponding density function in (3) is

$$\frac{1}{(2\pi(t-s))^{d/2}} e^{-\frac{|x|^2}{2(t-s)}}, \quad x \in \mathbb{R}^d.$$

The Brownian motion was first formulated by Einstein to study diffusion. Heuristically we can realize it as the following: it is direct to check that  $p(t, x) = (2\pi t)^{-\frac{d}{2}} e^{-|x|^2/2t}$  satisfies

$$\frac{\partial p(t, x)}{\partial t} = \frac{1}{2} \Delta p(t, x)$$

where  $\Delta = \sum_{i=1}^d \frac{\partial^2}{\partial x_i^2}$  is the Laplacian. Hence it satisfies the *heat equation*

$$\frac{\partial u}{\partial t} = \frac{1}{2} \Delta u \quad \text{on } \mathbb{R}^d.$$

If we are given an initial condition  $u(x, 0) = f(x)$ , it is known that the solution is given by

$$u(x, t) = \int_{\mathbb{R}^d} f(y) p(t, x - y) dy = (f * p_t)(x) = E_x(f(B_t)).$$

Equivalently, we can put it in terms of the Brownian motion  $u(x, t) = E(f(x - B_t))$ . The study of heat equation can be put entirely into a probabilistic setting.

In view of the two definitions above, there is another more general type of stochastic processes called *Lévy processes*. They are  $\{X_t\}_{t \geq 0}$  defined by replacing (3) with a *stationary increment* condition, i.e., for  $t > s$ ,  $X_t - X_s$  has the same distribution as  $X_{(t-s)}$ . The reader can refer to [Ito, Stochastic Process, Springer, 2004] for detail.

In the following we outline the theoretical existence of a probability space  $(\Omega, \mathcal{F}, P)$  for a stochastic process  $\{X_t\}_{t \geq 0}$ , and the measurability problem arised. The space and the  $\sigma$ -field are constructed by the family of finite dimensional distributions as for the discrete time case  $\{X_n\}_{n=1}^\infty$ .

Let  $T = [0, \infty)$  and let  $\mathbb{R}^T$  denote all functions  $\omega : T \rightarrow \mathbb{R}$ . For  $t_1 < \dots < t_n$ , the  $n$ -variate r.v.  $(X_{t_1}, \dots, X_{t_n})$  induces a distribution  $\mu_{t_1 \dots t_n}$  on  $\mathbb{R}^d$ . Let  $\mathcal{F}$  be the  $\sigma$ -field generated by  $(X_{t_1}, \dots, X_{t_n})$ , i.e., by sets (cylinder sets) of the form

$$E_{t_1 \dots t_n} = \{\omega : \omega(t_i) \in E_i\} \quad \text{with } E_i \text{ Borel sets, } 0 \leq t_1 \dots < t_n.$$

It can be checked that the family  $\{\mu_{t_1 \dots t_n}\}_{t_1 < \dots < t_n}$  satisfies the consistency condition:

$$\mu_{t_1 \dots t_{i-1} t_{i+1} \dots t_n} = \mu_{t_1 \dots t_n} \circ \varphi_i^{-1}$$

where  $\varphi_i : \mathbb{R}^n \rightarrow \mathbb{R}^{n-1}$ ,  $(x_1 \dots x_n) \rightarrow (x_1 \dots x_{i-1}, x_{i+1} \dots x_n)$  is the projection map. In this case  $\mu_{t_1 \dots t_{i-1} t_{i+1} \dots t_n}$  is the marginal distribution of  $\mu_{t_1 \dots t_n}$ . By the Kolmogorov extension theorem, there exists a probability  $P$  on  $(\Omega, \mathcal{F})$  and  $\{X_t\}_{t \geq 0}$  is the stochastic process with respect to  $(\Omega, \mathcal{F}, P)$ .

The probability space defined in this way is, however, still needed to be refined. One of the problems we often encounter is the measurability of union



of uncountably many sets with indices from  $T = [0, \infty)$ . Another problem is that the  $\sigma$ -field  $\mathcal{F}$  thus defined does not impose any condition on the continuity of the sample paths on  $[0, \infty)$  as is seen in the following example.

**Example.** Consider  $(\Omega, \mathcal{F}, P)$  on which there is a continuous random variable  $\tau$  with values in  $[0, T)$  (i.e.,  $P(\tau = t) = 0$  for all  $t \geq 0$ ). Define  $X_t(\omega) \equiv 0$  for all  $t \geq 0$ , and

$$Y_t(\omega) = \begin{cases} 1 & \text{if } \tau(\omega) = t \\ 0 & \text{if } \tau(\omega) \neq t \end{cases}$$

Then the only sample path of  $X(\cdot, \omega)$  is 0, but each sample path of  $Y(\cdot, \omega)$  has a jump at  $\tau(\omega) = t$ . On the other hand, it follows from the assumption on  $\tau$  that  $P(Y_t = 1) = P(\tau = t) = 0$  for each  $t$ , hence  $P(X_t = Y_t) = 1$  for each  $t$ . Therefore  $\{X_t\}_{t \geq 0}, \{Y_t\}_{t \geq 0}$  have the same finite dimensional distribution, they equals the point mass with probability 1 at the path  $\omega \equiv 0$ .

We will resolve the problem as follows:

**Definition 2.1.1.** *Two stochastic processes  $\{X_t\}_{t \geq 0}, \{Y_t\}_{t \geq 0}$  on  $(\Omega, \mathcal{F}, P)$  is called a version of each other if  $P(X_t = Y_t) = 1$  for all  $t \geq 0$ .*

Note that if we let  $N_t = \{X_t \neq Y_t\}$ , they are zero set with respect to  $P$ . We would like to have  $\bigcup_{t \geq 0} N_t$  to be a zero set, however, it is not necessary measurable from the construction of probability space. We will use the following theoretical device to overcome this dilemma. Let  $D$  be a countable subset of  $T = [0, \infty)$ , a function  $x : T \rightarrow \mathbb{R}$  is called *separable* if for any  $t \in T$ , there exists a sequence  $\{t_n\} \subseteq D$ ,  $t_n \rightarrow t$  and  $x(t_n) \rightarrow x(t)$ . For example continuous functions or right continuous functions are separable with respect to the rationals.

**Definition 2.1.2.** *A stochastic process  $\{X_t\}_{t \geq 0}$  on  $(\Omega, \mathcal{F}, P)$  is separable with respect to  $D$  if there exists an  $\mathcal{F}$ -null set  $N$  such that  $X(\cdot, \omega)$  is separable with*

respect to  $D$  for all  $\omega \notin N$ .

The process  $\{Y_t\}_{t \geq 0}$  in the Example is not separable. For if otherwise let  $D$  be a countable set in the definition. For any  $\tilde{t} \notin D$ , let  $\omega$  be such that  $\tau(\omega) = \tilde{t}$ , then  $Y(\tilde{t}, \omega) = 1$  and  $Y(t, \omega) = 0$  for all  $t \neq \tilde{t}$ . Hence for  $\{t_n\} \subseteq D$  and  $t_n \rightarrow t$ ,  $Y(t_n, \omega) \not\rightarrow Y(\tilde{t}, \omega)$ .

The following is the main theorem

**Theorem 2.1.3.** . *Let  $\{X_t\}_{t \geq 0}$  be a process on  $(\Omega, \mathcal{F}, P)$ , then there exists on the same space a separable process  $\{X'_t\}_{t > 0}$  such that  $P(X'_t = X_t) = 1$  for every  $t > 0$ .*

*Sketch of proof* ([2, p.555-559]). Note that for any fixed  $t$  and for any countable set  $D(\subset [0, T))$ , the set of  $\omega$  for which  $X(\cdot, \omega)$  is separable with respect to  $D$  at  $t$  can be written as

$$\bigcap_{n=1}^{\infty} \bigcup_{\substack{|s-t| < \frac{1}{n} \\ s \in D}} \{\omega : |X(s, \omega) - X(t, \omega)| < \frac{1}{n}\}.$$

The main task is to construct  $D$  (independent of  $t$ ) so the above set has probability 1. To prove this, we take any interval  $I \subseteq T$  and  $J \subseteq \mathbb{R}$ , and let

$$p(C) = P\left(\bigcap_{s \in C} (X_s \notin J)\right)$$

for any countable set  $C \subset I$ . Observe that as  $C$  increases,  $p(C)$  decreases. we can choose  $C_n$  such that  $p(C_n) \rightarrow \inf_C p(C)$ , and let  $U_{(I,J)} = \bigcup_n C_n$ . Then

$$P(\{X_t \in J\} \cap \bigcap_{s \in C_{I,J}} (X_s \notin J)) = 0$$

(otherwise, we consider  $C_{I,J} \cup \{t\}$  and obtain a contradiction). Let  $D = \bigcup C_{(I,J)}$  where  $(I, J)$  runs through all intervals  $I$  and  $J$  with rational end points. If we let

$$N(t) = \bigcup_{I,J} \left( \{X_t \in J\} \cap \bigcap_{s \in C_{I,J}} (X_s \notin J) \right)$$

Then we have  $P(N(t)) = 0$ . It is direct to check that  $D$  has the property we want. Now we define the separable version of  $\{X_t\}_{t \geq 0}$  as

$$X'(t, \omega) = \begin{cases} X(t, \omega) & \text{if } t \in D \text{ or } t \in D \text{ \& } \omega \notin N(t), \\ \limsup_{n \rightarrow \infty} X(s_n(t), \omega) & \text{if } t \notin D \text{ and } \omega \in N(t) \end{cases}$$

where  $s_n(t)$  is a fixed sequence converges to  $t$ . It follows that for each  $t$ , and for  $\omega \notin N(t)$ ,  $X'(\cdot, \omega)$  is separable with respect to  $D$  at  $t$ .  $\square$

We introduce the following definitions on a probability space  $(\Omega, \mathcal{F}, P)$ :

- a family  $\mathbb{F} = \{\mathcal{F}_t\}_{t \geq 0}$  of sub- $\sigma$ -fields in  $\mathcal{F}$  is called a *filtration* if  $\{\mathcal{F}_t\}$  is an increasing sequence of  $\sigma$ -fields on  $t$ ;
- a process  $\{X_t\}_{t \geq 0}$  is said to be *adaptable* to  $\mathbb{F}$  if  $X_t \in \mathcal{F}_t$  for each  $t > 0$ ;
- a filtration  $\mathbb{F}$  is called *right continuous* if  $\mathcal{F}_{t+} = \mathcal{F}_t$  (by definition  $\mathcal{F}_{t+} = \bigcap_{s > t} \mathcal{F}_s$ ).

For any filtration  $\{\mathcal{F}_t\}$ , let  $\mathcal{G}_t = \mathcal{F}_{t+}$ , then  $\mathbb{G} = \{\mathcal{G}_t\}_{t \geq 0}$  is right continuous. It is clear that if  $\{X_t\}_{t \geq 0}$  is adaptable to  $\mathbb{F}$ , then it is also adaptable to  $\mathbb{G}$ . For reasons that will be obvious later, we assume without loss of generality that  $\mathbb{F}$  is right continuous. It is also convenient to enlarge  $\mathcal{F}_0$  (hence all  $\mathcal{F}_t$ ) to include all subsets of the zero sets (completion by null sets).

With the filtration  $\mathbb{F}$ , we can define the necessary terminologies as before:

- Markov property:  $P(X_{t+s} \in E \mid \mathcal{F}_t) = P(X_{t+s} \in E \mid X_t)$  for  $t, s > 0$ ;
- Martingale:  $X_s = E(X_t \mid \mathcal{F}_s)$  for  $t > s$ ;
- Stopping time  $\alpha : \Omega \rightarrow [0, \infty)$  such that  $\{\tau \leq t\} \in \mathcal{F}_t$ .

**Exercises**

1. Let  $\xi : \Omega \rightarrow [0, \infty)$  be a random variable which satisfies

$$P(\xi \geq t + s \mid \xi \geq s) = e^{-\lambda s} \quad \forall t, s \geq 0$$

(*lack of memory property*). Show that this property is equivalent to  $\xi$  being an exponential distribution, i.e.,  $P(\xi \geq t) = e^{-\lambda t}$ ,  $t > 0$ , the waiting time with arrival rate  $\lambda$ .

2. Let  $X(t)$  be a Poisson process, let  $S_i = \inf\{t > 0 : X(t) = i\}$  and let  $\xi_n = S_n - S_{n-1}$  be the waiting time of the interarrivals. Show that the  $\{\xi_n\}_{n=1}^\infty$  are i.i.d. exponential random variables.

3. Conversely, let  $\{\xi_n\}_{n=1}^\infty$  be i.i.d. exponential random variables. Let  $\tau_n = \xi_1 + \cdots + \xi_n$ , and let

$$X(t) = \max\{n : \tau_n \leq t\}, \quad t > 0.$$

Show that  $X(t)$  is a Poisson process. (This is an alternative way to define a Poisson process.) Use the picture of a sample path to realize  $\tau_n$  and  $X(t)$  are “inverse” of each other (like the inverse function).

4. Show that if  $X$  is measurable in the sub- $\sigma$ -field  $\sigma\{X_t : t \in T\}$ , then  $X$  is measurable in  $\sigma\{X_t : t \in S\}$  for some countable subset  $S \subset T$ .

5. Let  $\{X_t\}_{t \geq 0}$  be a stochastic process on  $(\Omega, \mathcal{F}, P)$  and  $A \in \mathcal{F}$ . Show that there is a countable set  $S \subset T$  such that  $P(A \mid X_t, t \in T) = P(A \mid X_t, t \in S)$ .

6. Let  $K(s, t)$  be a real function over  $T \times T$ . Suppose that  $K$  is symmetric and nonnegative definite on  $T$ . Show that there is a process  $\{X_t\}_{t \geq 0}$  for which  $(X_{t_1}, \dots, X_{t_n})$  has the central (zero mean) normal distribution with covariance  $\text{cov}(X_{t_i}, X_{t_j}) = K(t_i, t_j)$ ,  $i, j = 1, \dots, k$ .

## 2.2 Brownian motion and sample paths

For a normal r.v.  $X \sim N(0, \sigma^2)$ , the symmetry implies  $E(X^{2k+1}) = 0$ , and the integration by parts yields

$$E(X^{2k}) = 1 \cdot 3 \cdot 5 \cdots (2k - 1) \cdot \sigma^{2k} . \quad (2.2.1)$$

We also need the following elementary properties of the normal r.v.'s :

– Suppose  $X_1 \sim N(\mu_1, \sigma_1^2)$ ,  $X_2 \sim N(\mu_2, \sigma_2^2)$ , and they are independent, then  $X_1 + X_2 \sim N(\mu_1 + \mu_2, \sigma_1^2 + \sigma_2^2)$ .

– Suppose  $\mathbf{X} = (X_1, \dots, X_n)$  is a  $n$ -variate normal r.v. with distribution  $N(\boldsymbol{\mu}, \Sigma)$  where  $\Sigma$  is a symmetric, positive definite  $n \times n$ -matrix. The density function is given by

$$f(\mathbf{x}) = \frac{1}{(2\pi)^{\frac{n}{2}} (\det \Sigma)^{\frac{1}{2}}} \exp \left( -\frac{1}{2} (\mathbf{x} - \boldsymbol{\mu})^t \Sigma^{-1} (\mathbf{x} - \boldsymbol{\mu}) \right) .$$

By a direct calculation, we have  $\Sigma = [\text{cov}(X_i, X_j)]$  where  $\text{cov}(X_i, X_j) = E((X_i - \mu_i)(X_j - \mu_j))$ .

– Suppose  $\mathbf{X} \sim N(\boldsymbol{\mu}, \Sigma)$ . Let  $\mathbf{Y} = A\mathbf{X} + \mathbf{c}$  where  $A$  is non-singular, then by a change of variable,  $\mathbf{Y}$  has density  $g(\mathbf{y}) = |\det A|^{-1} f(A^{-1}\mathbf{y} - \mathbf{c})$ . A direct substitution yields  $\mathbf{Y} \sim N(A\boldsymbol{\mu} + \mathbf{c}, \Sigma')$  where  $\Sigma' = A\Sigma A^t$ .

Let  $\{B_t\}_{t \geq 0}$  be the Brownian motion defined as in Section 2.1, then the process is *stationary* in the sense that the distribution  $B_t - B_s$  depends only on the difference  $t - s$ . Since  $B_t$  has distribution  $N(0, t)$ , it follows that

$$E(B_t) = 0 , \quad E(B_t^2) = t.$$

Moreover, by using independence, we have

$$E(B_s B_t) = \min(s, t) \quad (2.2.2)$$

This can be checked directly: assume  $s < t$ , then

$$\begin{aligned} E(B_s B_t) &= E(B_s(B_s + (B_t - B_s))) \\ &= E(B_s^2) + E((B_s(B_t - B_s))) \\ &= E(B_s^2) + E(B_s)E(B_t - B_s) = s. \end{aligned}$$

For  $0 < t_1 < t_2 \cdots < t_n$ , the joint distribution of  $(B_{t_1}, B_{t_2} - B_{t_1}, \cdots, B_{t_n} - B_{t_{n-1}})$  is given by

$$\begin{aligned} f_{t_1 \cdots t_n}(x_1, \cdots, x_n) &= \prod_{i=0}^n \frac{1}{\sqrt{2\pi(t_i - t_{i-1})}} \exp\left(-\frac{(x_i - x_{i-1})^2}{2(t_i - t_{i-1})}\right) \\ &= \frac{1}{\sqrt{2\pi}(\det \Sigma)^{1/2}} \exp\left(-\frac{1}{2}(\mathbf{z}^t \Sigma^{-1} \mathbf{z})\right) \end{aligned}$$

where  $\mathbf{z} = (x_1, x_2 - x_1, \cdots, x_n - x_{n-1})$  and

$$\Sigma = \begin{pmatrix} t_1 & 0 & \cdots & 0 \\ 0 & t_2 - t_1 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & t_n - t_1 \end{pmatrix}.$$

On the other hand the distribution of the  $n$ -variate random vector  $(B_{t_1}, B_{t_2}, \cdots, B_{t_n})$  is given by

$$g_{t_1, \dots, t_n}(\mathbf{x}) = \frac{1}{\sqrt{2\pi}(\det \Sigma')^{1/2}} \exp\left(-\frac{1}{2}(\mathbf{x}^t \Sigma'^{-1} \mathbf{x})\right) \quad (2.2.3)$$

where

$$\Sigma' = \begin{pmatrix} t_1 & t_1 & t_1 & \cdots & t_1 \\ t_1 & t_2 & t_2 & \cdots & t_2 \\ t_1 & t_2 & t_3 & \cdots & t_3 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ t_1 & t_2 & t_3 & \cdots & t_n \end{pmatrix}.$$

This follows from the transformation of the above multivariate normal random

vector :

$$\begin{pmatrix} B_{t_1} \\ B_{t_2} \\ \vdots \\ B_{t_n} \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 & \cdots & 0 \\ 1 & 1 & 0 & \cdots & 0 \\ \vdots & & & \vdots & \vdots \\ 1 & 1 & 1 & \cdots & 1 \end{pmatrix} \begin{pmatrix} B_{t_1} \\ B_{t_2} - B_{t_1} \\ \vdots \\ B_{t_n} - B_{t_{n-1}} \end{pmatrix}$$

and notice that  $\text{cov}(B_{t_i}, B_{t_j}) = \min\{t_i, t_j\}$ . By using this and the construction of the probability space in the last section, we can conclude the existence of a probability space for the Brownian motion.

One of the main properties of the Brownian motion is the continuity of the sample paths.

**Theorem 2.2.1.** *Let  $\{B_t\}_{t \geq 0}$  be a Brownian motion, then there is a version such that with probability 1, the sample paths  $B(\cdot, \omega)$  are continuous.*

**Proof.** Let  $D$  denote the set of dyadic rationals in  $[0, \infty)$  and let  $I_{n,k} = [\frac{k}{2^n}, \frac{k+1}{2^n})$  be the dyadic intervals. Let

$$E_n = \left\{ \omega : \max_{0 \leq k \leq n2^n} \left( \sup_{r \in I_{n,k} \cap D} |B(r, \omega) - B(\frac{k}{2^n}, \omega)| \right) > \frac{1}{n} \right\}.$$

We divide the proof into three steps.

(i) We claim that  $\sum_{n=1}^{\infty} P(E_n) < \infty$ . This will be proved in Lemma 2.2.5.

(ii) It follows from (i) and the Borel-Cantelli lemma that

$$E = \overline{\lim}_{n \rightarrow \infty} E_n = \bigcap_{l=1}^{\infty} \bigcup_{n=l}^{\infty} E_n$$

is a zero set. Observe that for any  $\omega \notin E$  there exists  $\ell$  such that for all  $n \geq \ell$ ,  $\omega \notin E_n$ . It follows that for any  $\epsilon > 0$  and  $t \in [0, \infty)$ , we can find  $n_0 > \max\{\ell, t\}$  and  $1/n_0 < \epsilon/3$ , such that for any  $n > n_0$ , we have (by  $\omega \notin E_n$ ),

$$|B(r, \omega) - B(\frac{k}{2^n}, \omega)| \leq \frac{1}{n}, \quad \forall r \in I_{n,k} \cap D, \quad 0 \leq \frac{k}{2^n} \leq n.$$

This implies that

$$|B(r, \omega) - B(r', \omega)| \leq \varepsilon, \quad \forall r, r' \in [0, t] \cap D, \quad |r - r'| \leq \frac{1}{2^n}.$$

We conclude from this that if  $\omega \notin E$ , then  $B(\cdot, \omega)$  is uniformly continuous on the dyadic rationals, and hence  $B(\cdot, \omega)$  has a continuous extension  $B'(\cdot, \omega)$  on  $[0, \infty)$ ,

$$B'_t(\omega) = B'(t, \omega) = \begin{cases} \lim_{r \rightarrow t} B(r, \omega) & \text{if } \omega \notin E \\ 0 & \text{if } \omega \in E \end{cases}$$

where the  $r$ 's are the dyadic rationals decrease to  $t$ .

(iii) Next we observe that the joint distribution of  $(B_{t_1}, \dots, B_{t_k})$  is the limit of the distributions of  $\{B_{r_1(n)}, \dots, B_{r_k(n)}\}_{n=1}^{\infty}$ , where the  $r_j(n)$ 's are rationals and  $r_j(n) \searrow t_j$ . (check this by the density functions (2.2.3)). Also note that  $(B'_{t_1}, \dots, B'_{t_k})$  also has the same distribution (see the following Lemma 2.2.2). Therefore  $\{B_t\}_t$  and  $\{B'_t\}_t$  have the same finite dimensional distributions in the same probability space. In view of  $P(B_t \neq B'_t) = P(E) = 0$  for each  $t \geq 0$ , we conclude that  $\{B'_t\}_t$  is a continuous version of  $\{B_t\}_t$ .  $\square$

The following simple lemma is needed in (iii).

**Lemma 2.2.2.** *Let  $\{X_n\}_n$  and  $X$  be  $k$ -dimensional r.v. Suppose  $X_n \rightarrow X$  in probability, and  $F_n(x) \rightarrow F(x)$  for all  $x$ , then  $F$  is the distribution function of  $X$ .*

**Proof.** We only prove the 1-dimensional case for simplicity. Let  $F_X$  be the distribution function of  $X$ . Since  $X_n \rightarrow X$  in probability, for  $\varepsilon > 0$ , there exists  $n$  such that for  $k > n$ ,  $P(|X_k - X| \geq \varepsilon) \leq \varepsilon$ . Hence for  $k > n$ ,

$$\begin{aligned} P(X_k \leq x) &\leq P(X \leq x + \varepsilon) + P(|X_k - X| \geq \varepsilon) \\ &\leq P(X \leq x + \varepsilon) + \varepsilon. \end{aligned}$$



It follows that  $\overline{\lim}_n F_n(x) \leq F_X(x)$ . By considering  $P(X_k > x + h)$ , we can use similar technique to show that for  $h > 0$ ,  $F_X(x) \leq \underline{\lim}_n F_n(x + h)$ . Putting the two inequalities together,

$$F(x) = \overline{\lim}_n F_n(x) \leq F_X(x) \leq \underline{\lim}_n F_n(x + h) = F(x + h)$$

Therefore  $F_X(x) = F(x)$  follow by taking  $h \rightarrow 0$ .  $\square$

Finally we prove  $\sum_{n=1}^{\infty} P(B_n) < \infty$  in (i), which will complete the proof of Theorem 2.2.1. We need a technical lemma.

**Lemma 2.2.3.** *Suppose  $X_1, \dots, X_n$  are independent r.v. and are symmetric about 0. Let  $S_n = X_1 + \dots + X_n$ . Then for  $\alpha > 0$  and  $\epsilon > 0$ ,*

$$(i) P(\max_{k \leq n} S_k \geq \alpha) \leq 2P(S_n \geq \alpha);$$

$$(ii) P(\max_{k \leq n} S_k \geq \alpha) \geq 2P(S_n \geq \alpha + 2\epsilon) - \sum_{k=1}^n P(X_k \geq \epsilon).$$

**Proof.** Note that

$$\begin{aligned} P(\max_{k \leq n} S_k \geq \alpha) &= P(\max_{k \leq n} S_k \geq \alpha, S_n \geq \alpha) + P(\max_{k \leq n} S_k \geq \alpha, S_n < \alpha) \\ &= P(S_n \geq \alpha) + P(\max_{k \leq n} S_k \geq \alpha, S_n < \alpha). \end{aligned}$$

(i) We need only show that the last term is  $\leq P(S_n \geq \alpha)$ . Let  $A_k = \{\max_{i < k} S_i < \alpha \leq S_k\}$  ( $k$  is the first time  $S_i \geq \alpha$ ). Then

$$\begin{aligned} P(\max_{k \leq n} S_k \geq \alpha, S_n < \alpha) &= \sum_{k=1}^{n-1} P(A_k \cap \{S_n < \alpha\}) \\ &\leq \sum_{k=1}^{n-1} P(A_k \cap \{S_n - S_k < 0\}) \\ &= \sum_{k=1}^{n-1} P(A_k \cap \{S_n - S_k > 0\}) \\ &\leq \sum_{k=1}^{n-1} P(A_k \cap \{S_n > \alpha\}) \\ &\leq P(S_n \geq \alpha). \end{aligned}$$

(Note that the key step is to switch “ $< 0$ ” to “ $> 0$ ” in the second equality, because  $A_k$  is independent of  $\{S_n - S_k\}$  and that  $S_n - S_k$  is symmetric about 0.) This proves (i).

(ii) We make use of the following two trivial relations

$$(a) \quad S_{k-1} < \alpha, X_k < \varepsilon, S_n - S_k < -\varepsilon \implies S_n < \alpha,$$

$$(b) \quad S_{k-1} < \alpha, X_k < \varepsilon, S_n \geq \alpha + 2\varepsilon \implies S_n - S_k > \varepsilon.$$

Following the same idea as in (i), we have

$$\begin{aligned} & \sum_{k=1}^{n-1} P(A_k \cap \{S_n < \alpha\}) \\ \geq & \sum_{k=1}^{n-1} P(A_k \cap \{X_k < \varepsilon, S_n - S_k < -\varepsilon\}) \quad (\text{by (a)}) \\ \geq & \sum_{k=1}^{n-1} P(A_k \cap \{X_k < \varepsilon, S_n - S_k > \varepsilon\}) \quad (\text{by indep. and symm.}) \\ \geq & \sum_{k=1}^{n-1} P(A_k \cap \{X_k < \varepsilon, S_n \geq \alpha + 2\varepsilon\}) \quad (\text{by (b)}) \\ \geq & \sum_{k=1}^{n-1} P(A_k \cap \{S_n \geq \alpha + 2\varepsilon\}) - P(X_k \geq \varepsilon) \\ \geq & P(S_n \geq \alpha + 2\varepsilon) - \sum_{k=1}^{n-1} P(X_k \geq \varepsilon). \end{aligned}$$

Combining with the previous part, we have (ii).  $\square$

It follows easily from the above that

**Corollary 2.2.4.** *Under the above assumption*

$$P\left(\max_{k \leq n} |S_k| \geq \alpha\right) \leq 2P(|S_n| \geq \alpha).$$

**Proof.** We make use of the symmetry:

$$\begin{aligned} P\left(\max_{k \leq n} |S_k| \geq \alpha\right) &= P\left(\max_{k \leq n} S_k \geq \alpha\right) + P\left(\max_{k \leq n} (-S_k) \geq \alpha\right) \\ &\leq 2(P(S_n \geq \alpha) + P(-S_n \geq \alpha)) = 2P(|S_n| \geq \alpha) \quad \square. \end{aligned}$$

Finally we prove the main lemma for Theorem 2.2.1.

**Lemma 2.2.5.** *With the notations in Theorem 2.2.1, we have  $\sum_n P(B_n) < \infty$ .*

**Proof.** We fix  $\delta$  and  $t$ , then by Lemma 2.2.3

$$\begin{aligned} P\left(\max_{i \leq 2^m} \left|B\left(t + \frac{i}{2^m}\delta\right) - B(t)\right| \geq \alpha\right) &\leq 2P(|B(t + \delta) - B(t)| \geq \alpha) \\ &\leq \frac{2}{\alpha^4} E(|B(t + \delta) - B(t)|^4) \\ &= \frac{6\delta^2}{\alpha^4}. \end{aligned}$$

(We have made use of  $P(|X| \geq \alpha) \leq \alpha^{-4} E(|X|^4)$ , and for  $X$  normal r.v.,  $E(X^4) = 3\sigma^4$ .) Let  $m \rightarrow \infty$ , we have

$$P\left(\sup_{0 < r < 1, r \in D} |B(t + r\delta) - B(t)| > \alpha\right) \leq \frac{6\delta^2}{\alpha^4}.$$

Therefore for  $E_n = \{\omega : \max_{0 \leq t \leq n2^n} (\sup_{r \in I_{n_k} \cap D} |B(r, \omega) - B(k2^{-n}, \omega)| > \frac{1}{n})\}$ ,

$$P(E_n) \leq n2^n (6 \cdot 2^{-2n}) / \left(\frac{1}{n}\right)^4 = 6n^5 2^{-n}.$$

Hence  $\sum P(B_n) < \infty$ .  $\square$

**Remark 1.** By Theorem 2.2.1, we can assume, in addition to (i)-(iii) in the Brownian motion,

(iv) For each  $\omega$ ,  $B(\cdot, \omega)$  is continuous.

**Remark 2.** In view of the estimation in Lemma 2.2.5 and the existence of a separable version for any given stochastic process (Theorem 2.1.3), Theorem 2.2.1 can be extended to the more general case:

**Theorem 2.2.6.** (Kolomogorov's continuity theorem) *Let  $\{X_t\}$  be a stochastic processes. Assume that there exists  $\alpha, \beta > 0$  such that*

$$E(|X(t) - X(s)|^\alpha) \leq K|t - s|^{1+\beta} \quad \forall t, s \geq 0.$$

*Then  $X(t)$  has a continuous version.*

The reader can refer to [3, p.31] for the detail.

**Definition 2.2.7.** A stochastic process  $\{X_t\}_{t \geq 0}$  is called a measurable process on  $(\Omega, \mathcal{F}, P)$  if  $X : T \times \Omega \rightarrow \mathbb{R}$  is  $\mathcal{B} \times \mathcal{F}$  measurable.

**Proposition 2.2.8.** The Brownian motion  $\{B_t\}_{t \geq 0}$  is a measurable process.

**Proof.** Let

$$B^{(n)}(t, \omega) = B\left(\frac{k}{2^n}, \omega\right), \quad \frac{k}{2^n} \leq t < \frac{k+1}{2^n}, \quad k = 0, 1, 2, \dots$$

Then the map  $B^{(n)}(\cdot, \cdot)$  is  $\mathcal{B} \times \mathcal{F}$  measurable, as

$$\{(t, \omega) : B^{(n)}(t, \omega) \geq a\} = \bigcup_{k,n} \left( [k2^{-n}, (k+1)2^{-n}) \times \{B_{k2^{-n}}^{(n)}(\omega) \geq a\} \right)$$

By the continuity of the sample path,  $B^{(n)}(t, \omega) \rightarrow B(t, \omega)$ , hence  $B(\cdot, \cdot)$  is  $\mathcal{B} \times \mathcal{F}$  measurable.  $\square$

As a corollary of the estimation in Lemma 2.2.3, we have

**Theorem 2.2.9.** For the Brownian motion  $\{B_t\}_{t \geq 0}$ , we have

$$P\left(\sup_{s \leq t} B_s \geq \alpha\right) = 2P(B_t \geq \alpha), \quad \forall \alpha \geq 0.$$

**Proof.** From Lemma 2.2.3(i), we have

$$P\left(\max_{k \leq 2^m} B_{k2^{-m}t} \geq \alpha\right) \leq 2P(B_t \geq \alpha).$$

Hence as  $m \rightarrow \infty$ , the continuity of  $B_{(\cdot)}(\omega)$  implies that

$$P\left(\sup_{s \leq t} B_s \geq \alpha\right) \leq 2P(B_t \geq \alpha)$$

On the other hand, Lemma 2.2.3(ii) implies

$$\begin{aligned} P\left(\sup_{s \leq t} B_s \geq \alpha\right) &\geq P\left(\max_{k \leq 2^m} B_{k2^{-m}t} \geq \alpha\right) \\ &\geq 2P\left(B_t \geq \alpha + \frac{2}{m}\right) - 2^m P\left(B_{2^{-m}t} \geq \frac{1}{m}\right). \end{aligned}$$

By the Chebychev inequality ( $P(|X| \geq \epsilon) \leq \epsilon^{-k} E(|X|^k)$ , using  $k = 4$ ), we conclude that the last term  $\leq 2^m(3(t2^{-m})^2)/m^{-4} = 3m^4 t^2 2^{-m}$ . This implies that

$$P\left(\sup_{s \leq t} B_s \geq \alpha\right) \geq 2P\left(B_t \geq \alpha\right)$$

and the theorem follows.  $\square$

We will give a simple proof this theorem again in Section 4 using the strong Markov property and the continuity of the sample paths. In the following we show that the almost all the sample paths are non-differentiable everywhere. First we observe a simple invariant property of the Brownian motion.

**Proposition 2.2.10.** (*Scaling property*) For  $c > 0$ , let

$$B'_t(\omega) = c^{-1} B_{c^2 t}(\omega) .$$

Then  $\{B'_t\}_{t \geq 0}$  is again a Brownian motion.

**Proof.** It is clear that  $\{B'_t\}_{t \geq 0}$  has independent increment, we only need to see the increment has a normal distribution with the correct variance. Recall that if  $X \sim N(0, \sigma^2)$ , then  $cX \sim N(0, c^2 \sigma^2)$ . Hence

$$B'_t(\omega) - B'_s(\omega) = c^{-1}(B_{c^2 t}(\omega) - B_{c^2 s}(\omega)).$$

It is a normal r.v. with variance  $c^{-2}(c^2 t - c^2 s) = t - s$ . This implies that  $\{B'_t\}_{t \geq 0}$  is a Brownian motion.  $\square$

**Theorem 2.2.11.** *Except for a set of zero probability,  $B(\cdot, \omega)$  is nowhere differentiable.*

**Proof.** Let

$$X_{n,k} = \max_{i=0,1,2} \left\{ \left| B\left(\frac{k+(i+1)}{2^n}\right) - B\left(\frac{k+i}{2^n}\right) \right| \right\}$$

be the maximum oscillation of  $B_t$  on three consecutive segments. Then by Proposition 2.2.10,

$$B\left(\frac{k+(i+1)}{2^n}\right) - B\left(\frac{k+i}{2^n}\right) \sim B_{2^{-n}} \sim 2^{-\frac{n}{2}} B_1.$$

Hence for any  $n$ ,  $k$  and  $\epsilon > 0$ , by independence, we have

$$P(X_{n,k} < \epsilon) = P(|B_1| \leq 2^{n/2}\epsilon)^3 = \left(\frac{1}{\sqrt{2\pi}} \int_{|x| < 2^{n/2}\epsilon} e^{-\frac{x^2}{2}} dx\right)^3 \leq (2 \cdot 2^{n/2}\epsilon)^3$$

Define

$$Y_n = \min_{k \leq 2^n} X_{n,k}$$

as the smallest oscillation of the  $\{X_{n,k}\}_k$ , then  $P(Y_n < \epsilon) \leq n2^n(2 \cdot 2^{n/2}\epsilon)^3$ . In particular,

$$P(Y_n < n2^{-n}) \leq n2^n(2 \cdot 2^{n/2} \cdot n2^{-n})^3 \rightarrow 0. \quad (2.2.4)$$

Now consider the upper and lower derivative of  $B(\cdot, \omega)$  from the right ,

$$D^+B(t, \omega) = \limsup_{h \rightarrow 0^+} (B(t+h, \omega) - B(t, \omega))/h$$

$$D_+B(t, \omega) = \liminf_{h \rightarrow 0^+} (B(t+h, \omega) - B(t, \omega))/h$$

Let  $E = \{\omega : \exists t > 0 \ni -\infty < D_+B(t, \omega) \leq D^+B(t, \omega) < \infty\}$ , we claim that  $P(E) = 0$ . Hence for  $\omega \notin E$ ,  $D^+B(t, \omega) = \infty$  or  $D_+B(t, \omega) = -\infty$ , and the theorem follows.

To prove the claim, let  $\omega \in E$ , then there exists  $K > 0$ ,

$$-K < D_+B(t, \omega) \leq D^+B(t, \omega) < K.$$

This implies that there exists  $\delta > 0$  such that for  $t < s < t + \delta$ ,

$$|B(s, \omega) - B(t, \omega)| \leq K|s - t|.$$

Let  $n_0$  be such that  $n_0 > \max\{4K, t\}$  and  $4/2^{n_0} < \delta$ ; for  $n > n_0$ , let  $k$  be such that

$$\left| \frac{k}{2^n} - t \right| < \delta, \quad k = 0, 1, 2, 3,$$

then

$$X_{n,k}(\omega) \leq 4K2^{-n} < n2^{-n}.$$

It follows that  $Y_n(\omega) \leq n2^{-n}$ . Let  $A_n = \{Y_n \leq n2^{-n}\}$ . Note that  $\omega \in E$  implies  $\omega \in A_n$  for  $n \geq n_0$ , i.e.,  $\omega \in \bigcup_{k=1}^{\infty} \bigcap_{n=k}^{\infty} A_n (= \varliminf_{n \rightarrow \infty} A_n)$ . Therefore by (2.2.4),

$$P(E) \leq P(\varliminf_n A_n) = \lim_{n \rightarrow \infty} P\left(\bigcap_{k=n}^{\infty} A_k\right) \leq \lim_{n \rightarrow \infty} P(A_n) \rightarrow 0$$

This proves the claim and the theorem follows.

**Remark.** It is well-known that the the regularity of the sample path can be made precise.

**Theorem** (Law of iterated logarithm). Let  $\{B_t\}_{t \geq 0}$  be a Brownian motion. Then

$$P\left(\varliminf_{s \rightarrow 0} \frac{B_{t+s} - B_s}{\sqrt{2t \log \log \frac{1}{t}}} = -1, \quad \varlimsup_{s \rightarrow 0} \frac{B_{t+s} - B_s}{\sqrt{2t \log \log \frac{1}{t}}} = 1\right) = 1.$$

The proof can be found in standard probability books (e.g., Breiman, Probability). There is a nice proof in “Diffusion Processes and Stochastic Calculus, Baudoin, 2014”, using Doob’s maximal inequality on the exponential martingale  $\{e^{\alpha B_t - \frac{\alpha^2}{2}t}\}_{t \geq 0}$ , and the Borel-Cantelli lemma. The theorem implies that the sample paths are Hölder continuous for order  $\frac{1}{2} - \varepsilon$  for any  $\varepsilon > 0$ . The reader can refer to Falconer, Fractal Geometry for a direct proof, also for the Hausdorff dimension of the paths.

**Exercices**

1. Show that the Poisson Process is a measurable process.
2. Let  $\{B_t\}_{t \geq 0}$  be Brownian motion. For fixed  $t$  and  $s$ , find the distribution of  $B_t + B_s$ .
3. Show that  $\lim_{t \rightarrow 0} tB(1/t) = 0$  almost surely. Define  $B'_t = tB_{1/t}$  for  $t > 0$ . Prove that  $\{B'_t\}_t$  is again a Brownian motion.
4. Show that  $\bigcap_{t > 0} \sigma\{B_s : s \geq t\}$  is a sub- $\sigma$ -field contains only sets of probability 0 and 1. Do the same for  $\bigcap_{\epsilon > 0} \sigma\{B_t : 0 < t < \epsilon\}$ ; give non-trivial examples in the  $\sigma$ -field.
5. Let  $\{W_t\}_{t \geq 0}$  be a stochastic process having independent, stationary increments and satisfies  $E(W_t) = 0$ ,  $E(W_t^2) = t$ . Show that if the finite-dimensional distributions are preserved by the scaling transformation  $W(t) \sim c^{-1}W_{c^2t}$ ,  $c > 0$ , then  $\{W_t\}_{t \geq 0}$  is a Brownian motion (Hint: use the Lindeberg theorem [2, p. 368]).
6. (Fourier expansion of Brownian motion)

- (a) Show that for  $s, t \in [-\pi, \pi]$ ,

$$\min(s, t) = \frac{ts}{\pi} + \frac{2}{\pi} \sum_{n \geq 1} \frac{\sin nt \sin ns}{n^2}.$$

- (b) Let  $\{X_n\}_{n=0}^{\infty}$  be i.i.d standard normal random variables, then

$$W_t = \frac{t}{\sqrt{\pi}} X_0 + \sqrt{\frac{2}{\pi}} \sum_{n \geq 1} \frac{\sin nt}{n} X_n$$

is a Brownian motion on  $[0, \pi]$  (see Breiman, Probability, 1968, P. 259-261).



## 2.3 Some basic properties

Let  $f : [0, t] \rightarrow \mathbb{R}$  be a real-valued function, we say that  $f$  is of *bounded variation* if

$$V(f) = \sup_{\mathcal{P}} \sum_{i=1}^n |f(t_i) - f(t_{i-1})| < \infty .$$

where the supremum is taken over all partition  $\mathcal{P} = \{0 = t_1 < t_2 < \dots < t_n = t\}$  of  $[0, t]$ . It is known that if  $f$  is of bounded variation, then  $f$  is differentiable a.e. A function  $f$  is said to have *quadratic variation* if the limit

$$\lim_{\|\mathcal{P}\| \rightarrow 0} \sum_{i=1}^n |f(t_i) - f(t_{i-1})|^2 \quad \text{exists ,}$$

where  $\|\mathcal{P}\| = \max_i \{t_i - t_{i-1}\}$ . The following shows that bounded variation and bounded quadratic variation are two non-compatible conditions.

**Proposition 2.3.1.** *If  $f : [0, t] \rightarrow \mathbb{R}$  is continuous and is of bounded variation, then  $f$  has zero quadratic variation.*

**Proof.** Observe that for any  $\mathcal{P} = \{0 = t_1 < \dots < t_n = t\}$ ,

$$\sum_{i=1}^n |f(t_i) - f(t_{i-1})|^2 \leq \max_i |f(t_i) - f(t_{i-1})| \cdot V(f)$$

By the uniform continuity of  $f$ , the above expression tends to 0 as  $\|\mathcal{P}\|$  tends to 0.  $\square$

**Theorem 2.3.2.** *Let  $[B](t)$  denote the quadratic variation of  $B_t$ , then  $[B](t) = t$  a.e.*

**Proof** Let  $\delta_n = \|\mathcal{P}_n\|$  and satisfies  $\sum_{n=1}^{\infty} \delta_n < \infty$ . For a partition  $\mathcal{P}_n$  with  $\|\mathcal{P}_n\| \leq \delta_n$ , let

$$T_n = \sum_{i=1}^{n_k} |B(t_i) - B(t_{i-1})|^2.$$

Then the expectation

$$E(T_n) = E\left(\sum_{i=1}^{n_k} |B(t_i) - B(t_{i-1})|^2\right) = \sum_{i=1}^{n_k} (t_i - t_{i-1}) = t. \quad (2.3.1)$$

We claim that

$$\sum_{n=1}^{\infty} E(T_n - E(T_n))^2 = \sum_{n=1}^{\infty} \text{Var}(T_n) < \infty \quad a.e.$$

It follows that  $E\left(\sum_{n=1}^{\infty} (T_n - E(T_n))^2\right) < \infty$ . Hence  $\sum_{n=1}^{\infty} (T_n - E(T_n))^2 < \infty$  a.e., and

$$\lim_{n \rightarrow \infty} (T_n - E(T_n)) = 0.$$

This together with (2.3.1) implies that  $[B](t) = \lim_{n \rightarrow \infty} E(T_n) = t$  a.e.

The claim follows from

$$\begin{aligned} \text{Var}(T_n) &= \text{Var}\left(\sum_{i=1}^{n_k} |B(t_i) - B(t_{i-1})|^2\right) \\ &= \sum_{i=1}^{n_k} \text{Var}(|B(t_i) - B(t_{i-1})|^2) \\ &\leq \sum_{i=1}^{n_k} E((B(t_i) - B(t_{i-1}))^4) \\ &= \sum_{i=1}^{n_k} 3 \cdot (t_i - t_{i-1})^2 \\ &\leq 3\|\mathcal{P}_n\| \cdot \sum_{i=1}^{n_k} (t_i - t_{i-1}) \leq 3t\delta_n \end{aligned}$$

and  $\sum_{n=1}^{\infty} \text{Var}(T_n) \leq 3t \sum_{n=1}^{\infty} \delta_n < \infty$ .

Recall that  $\{X_t\}_{t \geq 0}$  is a martingale if  $E(|X_t|) < \infty$  and for any  $s > 0$

$$E(X_{t+s} \mid \mathcal{F}_t) = X(t) \quad a.e.$$

Here  $\{\mathcal{F}_t\}_t$  is filtration (right continuous sub- $\sigma$ -field) generated by  $\{X_r : 0 \leq r \leq t\}$ .

**Theorem 2.3.3.** *The following processes are martingales:*

(i)  $\{B_t\}_{t \geq 0}$  ; (ii)  $\{B_t^2 - t\}_{t \geq 0}$  ; (iii)  $\{e^{\xi B_t - \frac{\xi^2}{2}t}\}_{t \geq 0}$  .

**Proof.** The proof depends on the independence of  $B_{t+s} - B_t$  and  $B_r$ ,  $0 \leq r \leq t$ , and also

$$E(g(B_{t+s} - B_t) \mid \mathcal{F}_t) = E(g(B_{t+s} - B_t))$$

where  $g$  is a Borel measurable function.

(i) Since  $B_t \sim N(0, t)$ ,  $E(|B_t|) < \infty$ . By independence,

$$\begin{aligned} E(B_{t+s} \mid \mathcal{F}_t) &= E(B_t + (B_{t+s} - B_t) \mid \mathcal{F}_t) \\ &= E(B_t \mid \mathcal{F}_t) + E(B_{t+s} - B_t \mid \mathcal{F}_t) \\ &= B_t + E(B_{t+s} - B_t) = B_t . \end{aligned}$$

(ii) Note that  $E(B_t^2) = t < \infty$  and

$$\begin{aligned} E(B_{t+s}^2) &= (B_t + (B_{t+s} - B_t))^2 \\ &= B_t^2 + 2B_t(B_{t+s} - B_t) + (B_{t+s} - B_t)^2 . \end{aligned}$$

Hence  $E(B_{t+s}^2 \mid \mathcal{F}_t) = B_t^2 + 0 + s$ . It follows that

$$E((B_{t+s}^2 - (t+s)) \mid \mathcal{F}_t) = B_t^2 - t .$$

(iii) It is easy to show by using completing square that

$$E(e^{\xi B_t}) = \int_{\mathcal{R}} e^{\xi x} \cdot \frac{1}{\sqrt{2\pi t}} e^{-x^2/2t} dx = e^{t\xi^2/2} .$$

We then apply the same proof as in (ii).  $\square$

A process  $\{X_t\}_{t \geq 0}$  is a Markov process if for any  $s, t > 0$ ,

$$P(X_{t+s} \in E \mid \mathcal{F}_t) = P(X_{t+s} \in E \mid X_t)$$

where  $\mathcal{F}_t$  generated by  $X_r$ ,  $0 \leq r \leq t$ .

**Theorem 2.3.4.**  $\{B_t\}_{t \geq 0}$  is a Markov process.

**Proof.** Since  $\mathcal{F}_t$  is generated by  $B_{t_1}, B_{t_2}, \dots, B_{t_n}$  for any  $0 < t_1 < \dots < t_n = t$ , to suffice to show that

$$P(B_{t+s} \in E \mid B_{t_1}, \dots, B_{t_n}) = P(B_{t+s} \in E \mid B_{t_n}). \quad (2.3.2)$$

Let  $X_1 = B_{t_1}$ ,  $X_i = B_{t_i} - B_{t_{i-1}}, i = 1, \dots, n$  and  $X_{n+1} = B_{t+s} - B_{t_n}$ . Also let  $S_i = X_1 + \dots + X_i$ , the sum of independent random variables. We have proved in Theorem 1.3.3 that

$$P(S_{n+1} \in E \mid S_1, \dots, S_n) = P(S_{n+1} \in E \mid S_n) = \mu_{n+1}(E - S_n)$$

This verifies (2.3.2) with  $S_i = B_{t_i}$  and  $\mu_{n+1}$  the density function of  $B_{t+s}$ .  $\square$

With the  $\{B_t\}_{t \geq 0}$  as a Markov process, it has a transition probability  $P(y, t; x, s) = P(B_t \leq y \mid B_s = s)$ . It follows that the density function is

$$f(y, t; x, s) = \frac{1}{\sqrt{2\pi(t-s)}} e^{-(y-x)^2/2(t-s)}.$$

The transition probability satisfies the stationary property  $P(y, t; x, s) = P(y, t-s; x, 0)$ .

Analogous to the discrete case, a random variable  $\tau : \Omega \rightarrow [0, \infty)$  is called a stopping time if

$$\{\tau \leq t\} \in \mathcal{F}_t \quad \forall t \geq 0 \quad (2.3.3)$$

where  $\{\mathcal{F}_t\}_{t \geq 0}$  is a filtration (see Section 2.1). It follows that if  $\tau$  is a stopping time, then  $\{\tau < t\} \in \mathcal{F}_t$ ; this follows from

$$\{\tau < t\} = \bigcup_{n=1}^{\infty} \left\{ \tau \leq t - \frac{1}{n} \right\} \in \mathcal{F}_{t-1/n} \subseteq \mathcal{F}_t.$$

Also by the right continuity of  $\{\mathcal{F}_t\}_{t \geq 0}$ , it is easy to show that “ $\{\tau < t\} \in \mathcal{F}_t$  for all  $t \geq 0$ ” actually equivalent to (2.3.3).

The pre- $\tau$ -field  $\mathcal{F}_\tau$  is defined as the family  $M \in \mathcal{F}$  such that

$$M \cap \{\tau \leq t\} \in \mathcal{F}_t, \quad t \geq 0. \quad (2.3.4)$$

The post- $\tau$  field  $\mathcal{F}'_\tau$  is defined as the sub- $\sigma$ -field generated by the process  $\mathcal{F}_{\tau+t}$ .

**Example.** Let  $\mathcal{F}_t = \sigma\{B_s : s \leq t\}$ , then  $\tau = \inf\{t : B_t = 1\}$  is a stopping time. Indeed let  $r$  denote a rational, then

$$\{\tau < t\} = \bigcup_{r < t} \bigcap_{m \geq 0} \{B_r \geq 1 - 1/m\} \in \mathcal{F}_t$$

The event  $M = \{\inf_{s < \alpha} B_s > -1\}$  is the set of paths that hit 1 before hit  $-1$ . It is in  $\mathcal{F}_\tau$  because ( $r, s$  are rationals)

$$M^c \cap \{\tau < t\} = \bigcup_{s < r < t} \bigcap_{m, n \geq 0} \{B_s \leq -(1 - 1/n), B_t \geq 1 - 1/m\} \in \mathcal{F}_t.$$

**Theorem 2.3.5.** *Let  $\tau$  be a stopping time finite a.e., then*

$$B_t^* = B_{\tau+t} - B_\tau$$

*is a Brownian motion. Moreover for  $M \in \mathcal{F}_\tau$ , and for  $E$  any Borel set in  $\mathbb{R}^k$ ,*

$$\begin{aligned} P(((B_{t_1}^* \cdots B_{t_k}^*) \in E) \cap M) &= P((B_{t_1}^*, \dots, B_{t_n}^*) \in E)P(M) \\ &= P((B_{t_1} \cdots B_{t_n}) \in E)P(M). \end{aligned}$$

**Proof.** We will prove the identities, then by taking  $M = \Omega$ ,  $(B_{t_1}^*, \dots, B_{t_k}^*)$  has the same distribution as  $(B_{t_1} \cdots B_{t_k})$ . Hence  $\{B_t^*\}_{t \geq 0}$  is a Brownian motion.

We first prove the case  $\tau$  has a countable range  $D$ . Note that  $B_t^*$  is in the post- $\tau$  field  $\mathcal{F}'_\tau$ , and for any Borel set  $E$  in  $\mathbb{R}$

$$\{B_t^* \in E\} = \bigcup_{s \in D} \{B_{s+t} - B_s \in E, \tau = s\}.$$

Let  $M \in \mathcal{F}_\tau$  and  $E \subseteq \mathbb{R}^k$ , then by the independence,

$$\begin{aligned}
& P(((B_{t_1}^*, \dots, B_{t_n}^*) \in E) \cap M) \\
&= \sum_{s \in D} P((B_{t_1}^*, \dots, B_{t_n}^*) \in E) \cap M \cap \{\tau = s\} \quad (2.3.5) \\
&= \sum_{s \in D} P((B_{t_1}^* \dots B_{t_n}^*) \in E) P(M \cap \{\tau = s\}) \\
&= P((B_{t_1}^*, \dots, B_{t_n}^*) \in E) P(M).
\end{aligned}$$

For the second identity, we take  $M = \Omega$ , then we can replace the  $(B_{t_1}^*, \dots, B_{t_n}^*)$  in (2.3.4) by  $(B_{t_1}, \dots, B_{t_n})$  and follow by the same argument.

For the general  $\tau$ , we let

$$\tau_n = \begin{cases} \frac{k}{2^n} & \text{if } \frac{k-1}{2^n} \leq \tau < \frac{k}{2^n}, \quad k = 1, 2, \dots, \\ 0, & \tau = \infty. \end{cases}$$

It is clear that  $\tau_n \searrow \tau$ . For  $k2^{-n} \leq \tau < (k+1)2^{-n}$ ,

$$\{\tau_n \leq t\} = \{\tau \leq k2^{-n}\} \in F_{k2^{-n}} \subseteq F_t.$$

This implies  $\tau_n$  is a stopping time. Let  $B_t^{(n)} = B_{t+\tau_n} - B_{\tau_n}$  and  $M \in \mathcal{F}_\tau (\subseteq F_{\tau_n})$ .

Then for  $H$  a closed rectangle in  $\mathbb{R}^k$ , by the above, we have

$$P(((B_{t_1}^{(n)}, \dots, B_{t_k}^{(n)}) \in H) \cap M) = P((B_{t_1}, \dots, B_{t_n}) \in H) P(M),$$

Since  $\{\tau_n(\omega)\}$  converges to  $\tau(\omega)$  and the sample paths are continuous, we can take limit so that

$$P(((B_{t_1}^*, \dots, B_{t_k}^*) \in H) \cap M) = P((B_{t_1}, \dots, B_{t_n}) \in H) P(M).$$

The rest of the theorem follows readily.  $\square$

**Theorem 2.3.6.** (The strong Markov property) *Let  $\tau$  be a stopping time finite a.e., then for any Borel set  $E \subseteq \mathbb{R}$ ,*

$$P(B_{t+\tau} \in E \mid \mathcal{F}_\tau) = P(B_{t+\tau} \in E \mid B_\tau).$$

**Proof.** We write  $B_{t+\tau} = (B_{t+\tau} - B_\tau) + B_\tau = B_t^* + B_t$ . It follows from Theorem 2.3.5 that  $B_\tau \in \mathcal{F}_\tau$  and  $B_t^* \in \mathcal{F}'_\tau$ , and they are independent. We can use Theorem 1.3.3 : Let  $X, Y$  be independent and  $Y \in \mathcal{F}$ , then  $P(X + Y \in E | \mathcal{F}) = P(X + Y \in E | Y)$ , to conclude the theorem.  $\square$

**Exercises**

1. Consider a process with three states  $\{a, b, c\}$ , and follows the rule that  $a$  goes to  $b$ ,  $b$  goes to  $c$  and  $c$  goes to  $a$ . This is a Markov chain. Show that

$$P(X_3 = c \mid X_2 = a \text{ or } b, X_1 = c) \neq P(X_3 = c \mid X_2 = a \text{ or } b).$$

Explain this situation in regard to the independence of the future and the past subject to the present.

2. Show that a process  $\{X(t)\}_{t \geq 0}$  with stationary and independent increment and right continuous sample paths has the strong Markov property.

3. Let  $\{N(t)\}_t$  be a Poisson process with rate  $\lambda$ . Prove the following are martingales: a.  $N(t) - \lambda t$ ; b.  $(N(t) - \lambda t)^2 - \lambda t$ ; c.  $e^{\log(1-\xi)N(t) + \xi\lambda t}$ ,  $0 < \xi < 1$ .

4. For  $T < \infty$ , is  $X(t) = B(T - t) - B(T)$  a Brownian motion on  $[0, T]$ ?



## 2.4 The exit time and hitting time

We use  $P_x(\cdot)$  to denote the probability of the Brownian motion starting at  $x$ .

For  $a \in \mathbb{R}$ , let

$$T_a = \inf\{t > 0 : B(t) = a\}$$

be the first time of the Brownian motion hitting  $a$ . Then  $T_a$  is a stopping time.

We first give two propositions to describe the exit and hitting probability.

**Proposition 2.4.1.** *Let  $a < x < b$  and  $\tau = \min\{T_a, T_b\}$  be the exit time, then  $P_x(\tau < \infty) = 1$  and the waiting time for exit is  $E_x(\tau) < \infty$ .*

**Proof.** Note that

$$\{\tau > 1\} = \{B(r) \in (a, b) \ \forall 0 < r < 1\} \subseteq \{B(1) \in (a, b)\}.$$

Then

$$\max_{z \in (a, b)} P_z(B(1) \in (a, b)) \leq \max_{z \in (a, b)} \left( \frac{1}{\sqrt{2\pi}} \int_a^b e^{-(z-y)^2/2} dy \right) := \theta < 1.$$

It follows that

$$\begin{aligned} & P_x(\tau > n) \\ &= P_x(\tau > n-1 \text{ and } B(r) \in (a, b) \ \forall n-1 < r \leq n) \\ &= P_x(\tau > n-1) P_x(B'(r) + B(n-1) \in (a, b) \ \forall 0 < r < 1 \mid \{\tau > n-1\}), \end{aligned}$$

where  $B'(r) = B(r + (n-1)) - B(n-1)$ . The last part can be estimated as follows:

$$\begin{aligned} & \leq P_x(B'(1) + B(n-1) \in (a, b) \mid \{\tau > n-1\}) \\ &= P_x(B'(1) + B(n-1) \in (a, b) \mid \{B_{n-1} \in (a, b)\}) \\ &= (P_x(B_{n-1} \in (a, b)))^{-1} \int_a^b P(B'(1) + y \in (a, b)) d\mu_x(y) \leq \theta, \end{aligned}$$

where  $\mu_x$  is the distribution of  $B_{n-1}$  starts at  $x$ . Hence

$$P_x(\tau > n) \leq P_x(\tau > n-1) \theta \leq \dots \leq \theta^n.$$

This implies that  $P_x(\tau = \infty) = 0$ , i.e.,  $P_x(\tau < \infty) = 1$ . For the second part, we make use of  $E(X) \leq \sum_{n=0}^{\infty} P(X > n)$  for  $X \geq 0$ :

$$E(\tau) \leq \sum_{n=1}^{\infty} \theta^n < \infty. \quad \square$$

**Proposition 2.4.2.** For any  $a, b \in \mathbb{R}$ ,  $P_a(P_b < \infty) = 1$ .

**Remark.** It follows that  $P_a(T_a < \infty) = 1$  at any  $a$ , and the path will return to  $a$  again and again. This property is called the *recurrent* property. We will see in Proposition 2.4.4 that, unlike Proposition 2.4.1, the waiting time for return is  $\infty$ .

**Proof.** We show that  $P_0(T_1 < \infty) = 1$ . The other cases are similar. Observe that for any  $a \neq b$ , the symmetry implies  $P_{(a+b)/2}(T_a < T_b) = \frac{1}{2}$ . Hence

$$P_0(T_{-1} < T_1) = \frac{1}{2}, \quad P_{-1}(T_{-3} < T_1) = \frac{1}{2}, \quad P_{-3}(T_{-7} < T_1) = \frac{1}{2}, \quad \dots$$

By the continuity of the paths, to reach  $-(2^n - 1)$ , they must pass through  $-1, -3, \dots$ . Let  $A_n = \{T_{-(2^n-1)} < T_1\}$ , by the strong Markov property,

$$P_0(A_n) = P_0(T_{-1} < T_1)P_{-1}(T_{-3} < T_1) \cdots P_{-(2^{n-1})}(T_{-(2^n-1)} < T_1).$$

It implies that  $P_0(A_n) = 2^{-n}$ , so that  $P_0(\bigcap_{n=1}^{\infty} A_n) = 0$ . This yields

$$1 = P_0\left(\bigcup_{n=1}^{\infty} A_n^c\right) = \lim_n P_0(T_1 \leq T_{-(2^n-1)}) = P_0(T_1 < \infty)$$

and the proposition follows.  $\square$

Let  $M(t) = \max_{0 \leq s \leq t} B(s)$ . It is clear that

$$\{M(t) \geq a\} = \{T_a \leq t\}. \quad (2.4.1)$$

**Theorem 2.4.3.** For  $a \in \mathbb{R}$ ,  $P(M(t) \geq a) = 2P(B(t) \geq a)$ .

**Remark.** We have proved this in Theorem 2.2.9. Here we give a simple proof by the hitting time, using the paths are continuous.

**Proof.** Note that  $\{B(t) \geq a\} = \{T_a \leq t, B(t) - B(T_a) \geq 0\}$ , and  $\{T_a \leq t\} \in \mathcal{F}_{T_a}$ , which is independent of  $\{B(t) - B(T_a) \geq 0\} \in \mathcal{F}'_{T_a}$ . Hence

$$\begin{aligned} P(\{B(t) \geq a\}) &= P(T_a \leq t, B(t) - B(T_a) \geq 0) \\ &= P(T_a \leq t)P(B(t) - B(T_a) \geq 0) \\ &= P(T_a \leq t)P(B^*(t - T_a) \geq 0) \\ &= P(T_a \leq t) \cdot \frac{1}{2} \quad (\text{by symmetry}) \\ &= \frac{1}{2}P(M(t) \geq a). \quad \square \end{aligned}$$

As an application of Theorem 2.4.3, we have

**Proposition 2.4.4.** The r.v.  $T_a : (\Omega, \mathcal{F}, P) \rightarrow [0, \infty)$  has density

$$f_{T_a} = \frac{|a|}{\sqrt{2\pi}} t^{-3/2} e^{-|a|^2/2t}, \quad t > 0, \quad (2.4.2)$$

and  $E(T_a) = \infty$ .

**Proof.** Let  $a > 0$ , we have by Theorem 2.4.2,

$$\begin{aligned} P(T_a \leq t) &= P(M(t) \geq a) = 2P(B(t) \geq a) \\ &= 2 \int_a^\infty e^{-y^2/2t} dy = \sqrt{\frac{\pi}{2}} \int_{\pi/\sqrt{t}}^\infty e^{-u^2} du. \end{aligned}$$

The density function follows from taking derivative of the above. Since the density of  $T_a$  is  $\approx t^{-3/2}$ , it is clear that  $E(T_a) = \infty$ .  $\square$

**Corollary 2.4.5.** For any  $0 < a < b$ ,  $T_b - T_a$  is independent of  $B(t)$ ,  $t \leq T_a$ ; the distribution function of  $T_b - T_a$  is

$$f_{T_b - T_a}(t) = \frac{b - a}{\sqrt{2\pi}} t^{3/2} e^{-(b-a)^2/2t} .$$

**Proof.** From Theorem 2.3.5,  $B^*(t) = B(t+T_a) - B(T_a)$  is a Brownian motion, and is independent of  $B(s)$ ,  $s \leq T_a$ . Hence the same is for

$$T_b - T_a = \inf\{t > 0 : B^*(t) = b - a\},$$

and the density is given by Proposition 2.4.4.  $\square$

We use the reflection property of the Brownian motion in Theorem 2.4.3. In the following, we formulate it into a theorem.

**Theorem 2.4.6.** (Reflection principle) Let  $\tau$  be a stopping time. Define

$$\widehat{B}(t) = \begin{cases} B(t), & \text{if } t \leq \tau \\ 2B(\tau) - B(t), & \text{if } t > \tau \end{cases}$$

Then  $\widehat{B}_t$  is also a Brownian motion.

**Remark.** Note that for  $t > \tau$ ,  $\widehat{B}_t(\omega) = -(B_t(\omega) - B_\tau(\omega)) + B_\tau(\omega)$  is the reflection along  $a = B_\tau(\omega)$ .

**Proof.** Let

$$C[0, \infty) = \{f \text{ continuous on } [0, \infty), f(0) = 0\}$$

be equipped with the  $\sigma$ -field generated by the cylinder sets. (It contains all the continuous sample paths of the Brownian motion.) Define a map  $\Phi : [0, \infty) \times C[0, \infty) \times C[0, \infty) \rightarrow C[0, \infty)$  by

$$\Phi(T, f, g) = \begin{cases} f(t), & \text{if } 0 \leq t \leq T, \\ f(t) + g(t - T), & \text{if } t \geq T. \end{cases}$$

It is clear that  $\Phi$  is measurable.

Note that  $B_{t \wedge \tau}$ , it is  $\mathcal{F}_\tau$ -measurable. Let  $B_s^* = B_{s+\tau} - B_\tau$ ,  $s > 0$ . Then both  $B_s^*$  and  $-B_s^*$  are Brownian motions with the same distribution, and are independent of  $\mathcal{F}_\tau$ . Hence  $(\tau, B_{t \wedge \tau}, B_s^*)$  and  $(\tau, B_{t \wedge \tau}, -B_s^*)$  have the same distribution (as r.v. on  $[0, \infty) \times C[0, \infty) \times C[0, \infty)$ ). It follows that

$$\Phi(\tau, B_{(\cdot) \wedge \tau}, B_{(\cdot)}^*) \quad \text{and} \quad \Phi(\tau, B_{(\cdot) \wedge \tau}, -B_{(\cdot)}^*)$$

have the same distribution. Note that the first one is just  $B_t$  and the second one is  $\widehat{B}(t)$ . We conclude that  $\widehat{B}(t)$  is also a Brownian motion.  $\square$

As a corollary we have

**Corollary 2.4.7.** *The joint distribution of  $(B(t), M(t))$  has density*

$$f_{B,M}(x, y) = \frac{2}{\sqrt{2\pi}} \frac{2y - x}{t^{3/2}} e^{-(2y-x)^2/2t}, \quad y \geq 0, x \leq 0.$$

**Proof.** Let  $y \geq 0, x \leq 0$ , and let  $\widehat{B}(t)$  be the reflection of  $B(t)$  at  $T_y$ . Then

$$\begin{aligned} P(B(t) \leq x, M(t) \geq y) &= P(B(t) \leq x, T_y \leq t) \\ &= P(\widehat{B}(t) \geq 2y - x, T_y \leq t) \\ &= P(\widehat{B}(t) \geq 2y - x) \\ &= 1 - \frac{1}{\sqrt{2\pi t}} \int_{2y-x}^{\infty} e^{-u^2/2t} du. \end{aligned}$$

The density function is obtained by taking partial derivatives on  $x$  and  $y$ .  $\square$

In the rest of the section, we consider the zeros of  $B(t)$ .

**Lemma 2.4.8.** *Let  $\mathcal{Z}_t = \{B(s) = 0 \text{ for some } s \in (0, t)\}$ . Then for  $a \neq 0$ ,*

$$P_a(\mathcal{Z}_t) = \frac{|a|}{\sqrt{2\pi}} \int_0^t u^{-3/2} e^{-a^2/2u} du.$$

**Proof.** Assume  $a > 0$ . Then

$$\begin{aligned}
 P_a(\mathcal{Z}_t) &= P(\min_{0 \leq s \leq t} (B(s) + a) \leq 0) \\
 &= P(\min_{0 \leq s \leq t} B(s) \leq -a) \\
 &= P(M(t) \geq a) \quad (\text{by symmetry}) \\
 &= P(T_a \leq t) \\
 &= \frac{a}{\sqrt{2\pi}} \int_0^t u^{-3/2} e^{-a^2/2u} du \quad (\text{by Proposition 2.4.4}).
 \end{aligned}$$

**Proposition 2.4.9.** *The probability that  $\{B(t)\}_t$  has at least one zero in time  $(r, s)$  is  $\frac{2}{\pi} \arccos \sqrt{\frac{r}{s}}$ .*

**Proof.** Let  $A_{r,s} = \{B(t) = 0 \text{ for some } t \in (r, s)\}$ , and let

$$h(x) = P(A_{r,s} \mid B_r = x) = P_x(A_{r,s}).$$

Hence by the above lemma,

$$\begin{aligned}
 P(A_{r,s}) &= \int_{-\infty}^{\infty} h(x) \frac{1}{\sqrt{2\pi r}} e^{-x^2/2r} dx \\
 &= \sqrt{\frac{2}{\pi r}} \int_0^{\infty} \left( \frac{x}{\sqrt{2\pi}} \int_0^{s-r} u^{-3/2} e^{x^2/2u} du \right) e^{-x^2/2r} dx \\
 &= \dots \\
 &= \frac{2}{\pi} \arctan \sqrt{\frac{s-r}{r}} = \frac{2}{\pi} \arccos \sqrt{\frac{r}{s}}. \quad \square
 \end{aligned}$$

**Corollary 2.4.10.** *For  $0 < r < s$ , the probability that no zero in  $(r, s)$  is  $\frac{2}{\pi} \arcsin \sqrt{\frac{r}{s}}$ .*

**Proof.** It follows from the above and

$$1 - \frac{2}{\pi} \arccos \sqrt{\frac{r}{s}} = \frac{2}{\pi} \arcsin \sqrt{\frac{r}{s}} \quad \square$$

To conclude, we prove a special property of the zero sets of the sample paths.

**Lemma 2.4.11.**  $P(\bigcap_{0 \leq t \leq 1} B(t) < 0) = P(\bigcap_{0 \leq t \leq 1} B(t) > 0) = 0$ .

**Proof.** We make use of Theorem 2.4.3 :

$$\begin{aligned} P\left(\bigcap_{0 \leq t \leq 1} B(t) \leq 0\right) &= P(\max_{0 \leq t \leq 1} B(t) \leq 0) \\ &= 1 - P(\max_{0 \leq t \leq 1} B(t) > 0) \\ &= 1 - 2P(B(1) > 0) \\ &= 1 - 2 \int_0^\infty \frac{1}{\sqrt{2\pi}} e^{-\frac{x^2}{2}} dx = 0. \quad \square \end{aligned}$$

**Lemma 2.4.12.** Let  $\mathcal{Z}_1(\omega) = \{t : B(t, \omega) = 0, 0 \leq t \leq 1\}$ , then for almost all  $\omega$ ,  $\mathcal{Z}_1(\omega)$  has 0 as a limit point.

**Proof.** From Lemma 2.4.11, we see that

$$P(B(t) \text{ crosses } 0 \text{ for some } 0 \leq t \leq 1) = 1.$$

By the scaling property (Proposition 2.2.10), we conclude that

$$P(B(t) \text{ crosses } 0 \text{ for some } 0 \leq t \leq r) = 1.$$

In particular, we take  $r_n \searrow 0$ . Then

$$P\left(\bigcap_{n=1}^{\infty} \{B(t) \text{ crosses } 0 \text{ for some } 0 \leq t \leq r_n\}\right) = 1.$$

This implies the lemma.  $\square$

**Theorem 2.4.13.** For almost all  $\omega$ ,  $\mathcal{Z}_1(\omega)$  is a perfect set (hence uncountable) and has Lebesgue measure zero .

**Proof.** We use  $|\mathcal{Z}_1|$  to denote the Lebesgue measure of  $\mathcal{Z}_1$ , it is a r.v. and

$$\begin{aligned} E(|\mathcal{Z}_1|) &= E\left(\int_0^1 \chi_{\{B(t)=0\}} dt\right) \\ &= \int_0^1 E(\chi_{\{B(t)=0\}}) dt \\ &= \int_0^1 P(B(t)=0) dt = 0. \end{aligned}$$

Hence  $|\mathcal{Z}_1| = 0$  P-a.e.

Next we note that  $B(\cdot, \omega)$  is continuous, hence  $\mathcal{Z}_1(\omega)$  is closed. We need to show that it has no isolated point, and it is a perfect set.

To this end for any rational  $r \in (0, 1)$ , let  $\tau_r$  be the least  $t \geq r$  such that  $B(t) = 0$ , then  $\tau_r$  is a stopping time. Let

$$A_r = \{\omega : \tau_r(\omega) \text{ is the limit point of } \mathcal{Z}_1(\omega)\}.$$

Then by the strong Markov property and Lemma 2.4.12, we have  $P(A_r) = 1$ . It follows that  $P(\bigcap_r A_r) = 1$  (where the intersection is taken over all rationals  $\geq 0$ ). Now for any  $\omega \in \bigcap_r A_r$  and for  $s \in \mathcal{Z}_1(\omega)$ ,  $s > 0$ , if  $s$  is a left limit point of  $\mathcal{Z}_1(\omega)$ , then it is not an isolated point. If  $s$  is not a left limit point of  $\mathcal{Z}_1(\omega)$ , then  $s = \tau_r(\omega)$  for some rational  $r < s$ . This implies that  $s$  is the right limit point of  $\mathcal{Z}_1(\omega)$  (by the strong Markov property, and use the lemma). In either case,  $s$  is not an isolated point. Hence  $\mathcal{Z}_1(\omega)$  has no isolated point and the proof of theorem is complete.



**Exercises**

1 Show that  $M(t) = \sup_{0 < s \leq t} \{B(s)\}$ ,  $|B(t)|$  and  $M(t) - B(t)$  have the same distribution.

2. For  $a, b > 0$ , let  $\tau = \min\{T_{-a}, T_b\}$  be the first that the Brownian motion hits  $a$  or  $b$ . What is  $P(B(\tau) = -a)$  and  $P(B(\tau) = b)$ ?

What are the hitting probabilities if we change  $a$  and  $b$  to two slant barriers  $-a + rt$  and  $b + rt$  for some  $r > 0$ .

3. Suppose  $X_1, \dots, X_n$  are independent and each has density function

$$h_a(t) = \frac{a}{\sqrt{2\pi}} t^{-3/2} e^{-a^2/2t}, \quad t > 0$$

(the density function for the first time the B.M. hits  $a$ ). Show that

(a)  $(X_1 + \dots + X_n)/n^2$  also has the same distribution. Contrast this with the law of large numbers.

(b)  $P((\max_{k \leq n} X_k)/n^2 \leq x) \rightarrow e^{-a\sqrt{2/(\pi x)}}$  for  $x > 0$ .

4. (a) Show that the probability of the last zero preceding time 1 is distribution over  $(0, 1)$  with density  $\pi^{-1}(t(1-t))^{-1/2}$ .

(b) Similarly calculate the distribution of the position of the first zero following time 1.

(c) Calculate the joint distribution of the two zeros in (a) and (b).



# Chapter 3

## Stochastic Integration and Ito Calculus

### 3.1 Wiener integral

The most basic integration theory is based on the Riemann integral of a real-valued function on an interval  $[0, T]$ . It is easy to extend the integrand to be a vector-valued or a Banach space-valued function. The Riemann Stieltjes integral is an extension such that the integrator is a function of bounded variation. The Lebesgue integral is to allow the integrator to be a measure on a more general measure space. In this section, we will consider the *Wiener integral*  $\int_0^T f(t)dB(t)$ , where  $f(t)$  is a real-valued function, integrating over the Brownian motion.

Suppose  $f$  is a real-valued step function on  $[0, T]$ ,  $f(t) = c_i$   $t_i \leq t < t_{i+1}$ ,  $i = 0, 1, \dots, n-1$ . We write  $\Delta B_i = B(t_{i+1}) - B(t_i)$ , and define

$$I(f) = \int_0^T f(t)dB(t) = \sum_{i=0}^{n-1} c_i \Delta B_i. \quad (3.1.1)$$

**Example 3.1.1.** Let  $f(t)$  be a step function on  $[0, 3]$  that takes values  $-1, 1, 2$  on the intervals  $[0, 1), [1, 2)$  and  $[2, 3]$  respectively. Then

$$\begin{aligned}
 I(f) &= \int_0^3 f(t)dB(t) \\
 &= (-1)(B(1) - B(0)) + 1 \cdot (B(2) - B(1)) + 2 \cdot (B(3) - B(2)) \\
 &= B(1) + (B(2) - B(1)) + 2 \cdot (B(3) - B(1)) \\
 &\sim N(0, 1) + N(0, 1) + N(0, 4) \quad (\text{independent sum}) \\
 &= N(0, 6).
 \end{aligned}$$

**Lemma 3.1.1.** Let  $f$  be a step function on  $[0, T]$ . Then  $I(f) \in L^2(\Omega)$ , it is a normal r.v. with mean 0 and variance

$$\sigma^2 = E((I(f))^2) = \int_0^T |f(t)|^2 dt .$$

**Proof.** Recall that if  $X_1, \dots, X_n$  are independent and  $X_i \sim N(\mu_i, \sigma_i^2)$ , then

$$a_1 X_1 + \dots + a_n X_n \sim N(a_1 \mu_1 + \dots + a_n \mu_n, a_1^2 \sigma_1^2 + \dots + a_n^2 \sigma_n^2).$$

It follows that  $I(f)$  is a normal r.v. with mean 0. For the variance, we have

$$\begin{aligned}
 E(I(f)^2) &= E\left(\sum_{i,j} c_i c_j \Delta B_i \Delta B_j\right) = \sum_i c_i^2 E((\Delta B_i)^2) \\
 &= \sum_i c_i^2 (t_{i+1} - t_i) = \int_0^T |f(t)|^2 dt . \quad \square
 \end{aligned}$$

**Lemma 3.1.2.** Suppose  $f \in L^2[0, T]$ , then there exists a sequence of step function  $\{f_n\}$  converges to  $f$  a.e., and  $\{I(f_n)\}_{n=1}^\infty$  is a Cauchy sequence in  $L^2(\Omega)$ .

**Proof.** The first statement is well known. The second statement follows from

$$E((I(f_n) - I(f_m))^2) = \int_0^T |f_n(t) - f_m(t)|^2 dt . \quad \square$$

**Definition 3.1.3.** For  $f \in L^2[0, T]$ , we define  $I(f) = \lim_{n \rightarrow \infty} I(f_n)$  where  $\{f_n\}$  is as in Lemma 3.1.2, and call  $I(f)$  the Wiener integral of  $f$ .

Clearly  $I(f) \in L^2(\Omega)$  by the completeness of  $L^2(\Omega)$  and the definition is independent of the choice of the subsequence  $\{f_n\}$ .

**Proposition 3.1.4.** For  $f \in L^2[0, T]$ ,  $I(f) \in L^2(\Omega)$  is a normal r.v. with mean 0 and variance  $\sigma^2 = \|f\|^2 = \int_0^T |f(t)|^2 dt$ .

**Proof.** We need to use the fact that if  $X_n \sim N(\mu_n, \sigma_n)$  and if  $\mu = \lim_{n \rightarrow \infty} \mu_n$ ,  $\sigma = \lim_{n \rightarrow \infty} \sigma_n$ , then  $X_n \rightarrow X$  in probability (or in  $L^2(\Omega)$ ) implies  $X \sim N(\mu, \sigma)$  (see Lemma 2.2.2).  $\square$

**Corollary 3.1.5.** If  $f, g \in L^2[0, T]$ , then  $E(I(f)I(g)) = \int_0^T f(t)g(t) dt$ .

**Proof.** This follows from

$$E((I(f) + I(g))^2) = \int_0^T |f + g|^2 = \int_0^T |f|^2 + 2 \int_0^T fg + \int_0^T |g|^2$$

and also

$$\begin{aligned} E((I(f) + I(g))^2) &= E(|I(f)|^2 + 2I(f)I(g) + |I(g)|^2) \\ &= \int_0^T |f|^2 + 2E(I(f)I(g)) + \int_0^T |g|^2. \quad \square \end{aligned}$$

Next we want to consider the relationship of  $(\int_0^T f(t)dB(t))(\omega)$  and  $\int_0^T f(t)dB(t, \omega)$ . Note the  $B(t, \omega)$  has unbounded variation (but has finite quadratic variation),  $\int_0^T f(t)dB(t, \omega)$  is not defined as a Riemann Stieltjes integral literally. On the other hand, we can redefine the integral as follows: For  $[a, b] \subseteq [0, T]$

$$(RS) \int_a^b f(t)dB(t, \omega) := \left[ f(t)B(t, \omega) \right]_a^b - \int_a^b B(t, \omega)df(t)$$

provided the last term is defined.

**Proposition 3.1.6.** *Let  $f$  be continuous and of bounded variation on  $[0, T]$ , then for any  $[a, b] \subseteq [0, T]$ ,*

$$\left( \int_a^b f(t) dB(t) \right)(\omega) = (\text{RS}) \int_a^b f(t) dB(T, \omega).$$

**Proof.** For any partition  $\mathcal{P}_n = \{t_0 < \dots < t_n\}$  of  $[a, b]$ , we consider the step function

$$f_n(t) = \sum_{i=0}^{n-1} f(t_i) \chi_{[t_{i+1}, t_i)}(t).$$

Note that the continuity of  $f$  implies  $f_n \rightarrow f$  in  $L^2[a, b]$  as  $\|\mathcal{P}_n\| \rightarrow 0$ . Hence

$$\int_a^b f(t) dB(t) = \lim_{\|\mathcal{P}_n\| \rightarrow 0} \sum_{i=0}^{n-1} f(t_i) \Delta B_i \quad \text{in } L^2(\Omega).$$

By passing to subsequence and using the same notation for convenience, we have

$$\left( \int_a^b f(t) dB(t) \right)(\omega) = \lim_{\|\mathcal{P}_n\| \rightarrow 0} \sum_{i=0}^{n-1} f(t_i) \Delta B_i(\omega) \quad \text{a.e.}$$

On the other hand, for almost all  $\omega$ ,

$$\begin{aligned} (\text{RS}) \int_a^b f(t) dB(t, \omega) &= \left[ f(t) B(t, \omega) \right]_a^b - \lim_{\|\mathcal{P}_n\| \rightarrow 0} \sum_{i=1}^n B(t_i, \omega) (f(t_i) - f(t_{i-1})) \\ &= \lim_{\|\mathcal{P}_n\| \rightarrow 0} \sum_{i=0}^{n-1} f(t_i) \Delta B_i(\omega). \end{aligned}$$

(The first limit exists as  $f$  is of bounded variation; the second equality follows from the Abel transform of series.) This yields the proposition.  $\square$

**Example 3.1.2.**  $\int_0^1 B(t) dt \sim N(0, 1/3)$ .

By regarding  $B(\cdot) \in C([0, 1], L^2(\Omega))$ , we have by the definition of Reimann

integral,  $(\int_0^1 B(t) dt)(\omega) = \int_0^1 B(t, \omega) dt$  for almost all  $\omega$ , and

$$\begin{aligned} \int_0^1 B(t, \omega) dt &= [tB(t, \omega)]_0^1 - (\text{RS}) \int_0^1 t dB(t, \omega) \\ &= B(1, \omega) - (\text{RS}) \int_0^1 t dB(t, \omega) \\ &= (\text{RS}) \int_0^1 (1-t) dB(t, \omega) \\ &= \left( \int_0^1 (1-t) dB(t) \right)(\omega) \quad (\text{by Proposition 3.1.6}) \end{aligned}$$

Since  $\int_0^1 (1-t)dB(t) \sim N(0, \sigma^2)$  with  $\sigma^2 = \int_0^1 (1-t)^2 dt = 1/3$  (Proposition 3.1.4), we have  $\int_0^1 B(t)dt \sim N(0, 1/3)$ .

**Theorem 3.1.7.** *Let  $f \in L^2[0, \infty)$ , then*

$$Y(t) = \int_0^t f(s)dB(s)$$

*is a Gaussian process with mean 0 and covariance  $\text{cov}(Y(t), Y(t+s)) = \int_0^t |f(u)|^2 du$  for all  $s, t \geq 0$ .*

*Moreover  $\{Y(t)\}_{t \geq 0}$  is a martingale with respect to  $\mathcal{F}_t = \sigma\{B(s) : s \leq t\}$ .*

(Recall that Gaussian process means the joint distribution at  $t_1, \dots, t_k$  is a multivariate normal r.v. which is determined by the mean and the covariance.)

**Proof.** We observe that for  $u, t \geq 0$ ,

$$E\left(\int_t^{t+u} f(s)dB(s) \mid \mathcal{F}_t\right) = 0.$$

Indeed this holds for step functions on  $[t, t+u)$  as

$$\int_t^{t+u} f(s)dB(s) = \sum_{i=0}^{n-1} c_i \Delta B_i$$

and  $E(\Delta B_i \mid \mathcal{F}_t) = E(\Delta B_i) = 0$ . For  $f \in L^2([0, T])$ , we can find step functions  $f_n \rightarrow f$  in  $L^2$ . Hence  $I_n(f) \rightarrow I(f)$  and

$$E\left(\int_t^{t+u} f(s)dB(s) \mid \mathcal{F}_t\right) = \lim_{n \rightarrow \infty} E\left(\int_t^{t+u} f_n(s)dB(s) \mid \mathcal{F}_t\right) = 0.$$

We already see that each  $Y(t)$  is a normal r.v. with mean 0 and variance  $\int_0^t |f(s)|^2 ds$ . The covariance is

$$\begin{aligned} \text{Cov}(Y(t), Y(t+u)) &= E(Y(t)Y(t+u)) \\ &= E\left(E(Y(t)(Y(t) + \int_t^{t+u} f(s)dB(s)) \mid \mathcal{F}_t)\right) \\ &= E(|Y(t)|^2) = \int_0^t |f(s)|^2 ds . \end{aligned}$$

It is direct to see that the joint distribution  $(Y_{t_1}, \dots, Y_{t_n}) \sim N(0, \Sigma)$  with  $\Sigma$  determined by this covariance.

To show that  $\{Y_t\}_{t \geq 0}$  is a martingale, we observe that  $E(|Y(t)|^2) = \int_0^t |f(s)|^2 ds < \infty$  and

$$E(Y(t+u) \mid \mathcal{F}_t) = Y(t) + E\left(\int_t^{t+u} f(s)dB(s) \mid \mathcal{F}_t\right) = Y(t). \quad \square$$



**Exercises**

1. Let  $X(t) = tB(t)$ . Find the quadratic variation of  $X(t)$ .
2. Find all constants  $a$ ,  $b$  and  $c$  such that  $X(t) = \int_0^t (a + bu/t + c(u/t)^2) dB(t)$  is also a Brownian motion.
3. Let  $B(t)$  be a Brownian motion. Show that  $X(t) = \int_0^t (2t - u)dB(u)$  and  $Y(t) = \int_0^t (3t - 4u)dB(u)$  are Gaussian processes with the same mean and covariance functions.
4. Find the distribution of the integral  $\int_0^t B(s) \cos(t - s)ds$ .
5. Given values of  $\alpha$  for which the process  $Y(t) = \int_0^t (t - s)^{-\alpha} dB(s)$  is defined. Find the covariance function of  $Y$ . (This process is called fractional Brownian motion.)
6. Show that if  $\{X_n\}$  is a sequence of normal r.v. and convergent in distribution to  $X$ , the  $X$  is either a normal r.v. or degenerate. Deduce that if  $E(X_n) \rightarrow \mu$  and  $\text{var}(X_n) \rightarrow \sigma^2$ , then  $X \sim N(\mu, \sigma)$ .

### 3.2 Ito integral

In this section we define the Ito integral  $\int_0^T X(t)dB(t)$  where  $X(t)$  is a stochastic process *adapted* to  $\mathcal{F}_t = \sigma\{B(s) : 0 \leq s \leq t\}$ .

First we consider the case  $X(t)$  is a step process on  $[0, T]$ , i.e., there exist  $0 = t_0 < \dots < t_n = T$  such that

$$X(t) = \xi_i, \quad t_i \leq t < t_{i+1}, \quad \xi_i \in \mathcal{F}_{t_i} \quad \text{and} \quad E(|\xi_i|^2) < \infty.$$

We define

$$I(X) = \int_0^T X(t)dB(t) = \sum_{i=0}^{n-1} \xi_i \Delta B_i.$$

**Proposition 3.2.1.** *For the above integral  $I(\cdot)$  on the step processes,*

- (i)  $I(\cdot)$  is linear;
- (ii) for  $[a, b] \subset [0, T]$ ,  $\int_0^T \chi_{[a,b]}(t)dB(t) = B(b) - B(a)$ ;
- (iii)  $E(\int_0^T X(t)dB(t)) = 0$ ;
- (iv)  $E(\int_0^T X(t)dB(t))^2 = \int_0^T E(X(t)^2)dt$ .

(Note that (iii),(iv) imply  $\int_0^T X(t)dB(t) \in L^2(\Omega)$  is a r.v. on  $\Omega$  with mean 0 and  $\sigma^2 = \int_0^T E(X(t)^2)dt$ .)

**Proof.** For (iii), it follows from the independence of  $\xi_i$  and  $\Delta B_i$ ,

$$E(\xi_i \Delta B_i) = E(\xi_i)E(\Delta B_i) = 0.$$

For (iv), we write  $E(|I(f)|^2) = \sum_{i,j} E(\xi_i \xi_j \Delta B_i \Delta B_j)$ . Then using the independence, it is direct to show that only the  $i = j$  terms left, which is of the expression on the right side of (iv).  $\square$

We use  $L_{ad}^2([0, T] \times \Omega)$  to denote the space of measurable process  $X(\cdot, \cdot)$  on  $[0, T] \times \Omega$  satisfying

- (i)  $X(t, \cdot)$  is adapted to  $\{\mathcal{F}_t\}$ , i.e.,  $X(t, \cdot) \in \mathcal{F}_t$ ;
- (ii)  $\int_0^T E(X(t)^2)dt < \infty$ .

For example,  $X(t) = B(t)$  and  $X(t) = \max_{0 \leq s \leq t} B(s)$  are adaptable to  $\{\mathcal{F}_t\}$ ; but  $X(t) = B(t+1)$  is not adaptable to  $\{\mathcal{F}_t\}$ .

**Lemma 3.2.2.** *Suppose  $X \in L_{ad}^2([0, T] \times \Omega)$ , then there exist a sequence of step processes  $\{X^{(n)}\} \subseteq L_{ad}^2([0, T] \times \Omega)$  such that*

$$\lim_{n \rightarrow \infty} \int_0^T E(|X^{(n)}(t) - X(t)|^2)dt = 0. \quad (3.2.1)$$

**Proof.** The idea is similar to the Wiener integral, but a little more complicated. We divide the proof into three steps.

(i) if  $E(X(t)X(s))$  is continuous for any  $s, t \in [0, T]$ , let  $\mathcal{P}_n = \{0 = t_0 < t_1 \cdots < t_n = T\}$  be a partition of  $[0, T]$ , and let

$$X^{(n)}(t) = X(t_i), \quad t_i \leq t < t_{i+1}.$$

From the assumption,  $\lim_{n \rightarrow \infty} E(|X(s) - X(t)|^2) = 0$  implies that  $\lim_{n \rightarrow \infty} E(|X^{(n)}(t) - X(t)|^2) = 0$ . As

$$E(|X^{(n)}(t) - X(t)|^2) \leq 2(E(|X^{(n)}(t)|^2) + E(|X(t)|^2)) \leq 4 \sup_{a \leq s \leq b} E(X(s)|^2).$$

By the dominated convergence theorem,

$$\lim_{n \rightarrow \infty} \int_0^T E(|X^{(n)}(t) - X(t)|^2)dt = 0.$$

(ii) If  $X$  is bounded, then we will construct a sequence of process  $\{Y_n\}_n$  adaptable to  $\{\mathcal{F}_t\}$ , satisfies (i), and  $E(|Y_n(t) - X(t)|^2) \rightarrow 0$ . Then using (i), we

can find a step process  $\{X^{(n)}\}_n$  such that  $E(|X^{(n)}(t) - E(Y_n(t))|^2)$  is sufficiently small, and goes to 0 as  $n \rightarrow \infty$ . Then

$$E(|X^{(n)}(t) - E(X(t))|^2) \rightarrow 0,$$

which yields (3.2.1) as in (i).

To this end, we let  $\varphi_n(t) = ne^{-nt}$ ,  $t \geq 0$ . Note that  $\varphi_n \geq 0$ ,  $\int \varphi_n = 1$  and  $\varphi_n(t)dt \rightarrow \delta_0$ , hence it is an approximate identity. Define

$$Y_n(t) = \varphi_n * X(t) = \int_0^t \varphi_n(s)X(t-s)ds.$$

It is direct to check that

- (a)  $Y_n(t)$  is adaptable to  $\mathcal{F}_t$ ;
- (b)  $\lim_{s \rightarrow t} E(|Y_n(s) - Y_n(y)|^2) = 0$ , hence (i) is satisfied;
- (c)  $\lim_{n \rightarrow \infty} E(|Y_n(t) - X(t)|^2) = 0$ .

(iii) Finally, for the general  $X \in L_{ad}^2([0, T] \times \Omega)$ , we can approximate  $X$  by a sequence of bounded processes:

$$X_n(t, \omega) = \begin{cases} X(t, \omega) & \text{if } |x(t, \omega)| \leq n; \\ 0 & \text{if } |X(t, \omega)| > n. \end{cases}$$

and a routine argument yields (3.2.1).  $\square$

Let  $X \in L_{ad}^2([0, T] \times \Omega)$ , and let  $X^{(n)}$  be as in Lemma 3.2.2. Let  $Y^{(n)} = \int_0^T X^{(n)}(t)dB(t) \in L^2(\Omega)$ . Then by Proposition 3.2.1 (iv)

$$E(|Y^{(n)} - Y^{(m)}|^2) = \int_0^T E(|X^{(n)}(t) - X^{(m)}(t)|^2)dt \longrightarrow 0.$$

Hence  $\{Y^{(n)}\}$  is a Cauchy sequence in  $L^2(\Omega)$ .

**Definition 3.2.3.** For  $X \in L^2_{ad}([0, T] \times \Omega)$ , we define

$$I(X) = \int_0^T X(t)dB(t) = \lim_{n \rightarrow \infty} \int_0^T X^{(n)}(t)dB(t) \in L^2(\Omega),$$

and call  $I(X)$  the Ito integral of  $X$  over  $B(t)$ .

We remark that if  $X \in L^2_{ad}([0, T] \times \Omega)$  and assume that  $E(X(s)Y(t))$  is continuous as in the proof of Lemma 3.2.2, then the integral can be expressed as the Riemann sum

$$\int_0^T X(t)dB(t) = \lim_{\|\mathcal{P}_n\| \rightarrow 0} \sum_{i=0}^{n-1} X(t_i)\Delta B_i.$$

**Example 3.2.1**  $\int_0^T B(t)dB(t) = \frac{1}{2}(B(T)^2 - T)$ .

For  $0 = t_0 < \dots < t_n = T$ , let

$$X^{(n)}(t) = B(t_i) \quad t_i \leq t < t_{i+1}.$$

Then  $I(X^{(n)}) = \sum_{i=0}^{n-1} B(t_i)\Delta B_i$ . Observe that  $a(b-a) = \frac{1}{2}(b^2 - a^2 - (b-a)^2)$ .

Hence

$$\begin{aligned} I(X^{(n)}) &= \frac{1}{2} \sum_{i=0}^{n-1} (B^2(t_{i+1}) - B^2(t_i)) - \frac{1}{2} \sum_{i=0}^{n-1} (B(t_{i+1}) - B(t_i))^2 \\ &= \frac{1}{2}(B(T)^2 - B(0)^2) - \frac{1}{2} \sum_{i=0}^{n-1} (\Delta B_i)^2. \end{aligned}$$

By taking limit, the second part is the quadratic variation of  $B(t)$ , which converges to  $T$  (Theorem 2.3.2). Hence

$$I(B) = \int_0^T B(t)dB(t) = \frac{1}{2}(B(T)^2 - T).$$

**Example 3.2.2.**  $\int_0^T B(t)^2dB(t) = \frac{1}{3}B(T)^3 - \int_0^T B(t)dt$ .

We adopt the same method as last example. By making use of  $a^2(b-a) = \frac{1}{3}(b^3 - a^3 - (b-a)^3 - 3a(b-a)^2)$ , we have

$$\sum_{i=0}^{n-1} B(t_i)^2 \Delta B_i = \frac{1}{3}(B(T)^3 - B(0)^3) - \sum_{i=0}^{n-1} (\Delta B_i)^3 - 3 \sum_{i=0}^{n-1} B(t_i)(\Delta B_i)^2. \quad (3.2.2)$$

Note that

$$E\left(\left|\sum_{i=0}^{n-1} (\Delta B_i)^3\right|^2\right) = \sum_{i=0}^{n-1} E((\Delta B_i)^6) = \sum_{i=0}^{n-1} 15(t_{i+1} - t_i)^3 \leq 15\|\mathcal{P}_n\|^2 \cdot T,$$

which tends to 0 as  $\|\mathcal{P}_n\| \rightarrow 0$ . Hence  $\sum_{i=0}^{n-1} (\Delta B_i)^3 \rightarrow 0$  a.e. as well.

For the last term, we observe that

$$E\left(\left|\sum_{i=0}^{n-1} B(t_i)(\Delta B_i)^2 - \sum_{i=0}^{n-1} B(t_i)\Delta t_i\right|^2\right) = \sum_{i=0}^{n-1} 2t_i \Delta t_i^2 \leq 2\|\mathcal{P}_n\| \cdot T^2 \rightarrow 0.$$

Hence the limit of the last term in (3.2.2) equals  $\int_0^T B(t)dt$ .

**Theorem 3.2.4.** For  $X \in L_{ad}^2([0, T] \times \Omega)$ , the Ito integral

$$Y(t) = \int_0^t X(s)dB(s)$$

has mean zero and variance  $\sigma^2 = E(Y(t))^2 = \int_0^t E(|X(t)|^2)dt$ . Moreover  $\{Y_t\}_t$  is a martingale.

**Proof.** The first part is a consequence of the approximation by adaptable step processes as in Lemma 3.2.2. The second part follows from the same proof as for the Wiener integral (Theorem 3.1.7).  $\square$

In the following we consider the continuity of the sample path of  $Y(t)$ . We need a continuous version of Doob's submartingale inequality (Theorem 1.4.9, Corollary 1.4.10).

**Lemma 3.2.5.** *Let  $\{Y(t)\}_{t \geq 0}$  be a submartingale and assume that almost all sample paths of  $Y(t)$  are continuous. Then*

$$\lambda P\left(\sup_{0 \leq t \leq T} Y(t) \geq \lambda\right) \leq E(Y(T)^+).$$

Furthermore, we have

$$\lambda P\left(\sup_{0 \leq t \leq T} |Y(t)| \geq \lambda\right) \leq E(|Y(T)|).$$

**Proof.** The continuity of the sample paths allows us to write

$$\left\{\sup_{0 \leq t \leq T} Y(t) \geq \lambda\right\} = \bigcap_{n=1}^{\infty} \bigcup_{m=1}^{\infty} \left\{\max_{0 \leq \frac{k}{2^m} \leq T} Y(k/2^m) \geq \lambda - \frac{1}{n}\right\}.$$

By applying Theorem 1.4.9 and Corollary 1.4.10 to the submartingale

$$\{Y(1/2^m), Y(2/2^m), \dots, Y(l/2^m), Y(T)\},$$

and taking limit, the lemma follows.  $\square$

**Theorem 3.2.6.** *Suppose  $X \in L_{ad}^2([0, T] \times \Omega)$ , then the Ito integral  $\{Y(t)\}_t$  is a continuous process on  $[0, T]$ , i.e., almost all its sample paths are continuous.*

*Proof.* We consider the step process first. Let  $X(t) = \sum_{i=0}^{n-1} \xi_i \chi_{[t_i, t_{i+1})}$ ,  $\xi_i \in \mathcal{F}_{t_i}$ .

Then

$$Y(t, \omega) = \sum_{i=0}^{k-2} \xi_i(\omega) \Delta B_i(\omega) + \xi_{k-1}(B(t, \omega) - B(t_{k-1}, \omega)).$$

It is easy to see that  $Y(\cdot, \omega)$  is continuous as  $B(\cdot, \omega)$  is continuous.

Next we consider the general case, let  $\{X^{(n)}\}_n$  be a sequence of step stochastic processes in  $L_{ad}^2([0, T] \times \Omega)$  such that

$$\lim_{n \rightarrow \infty} \int_0^T E(|X(S) - X^{(n)}(s)|^2) ds = 0.$$

By choosing a subsequence if necessary, we may assume that

$$\int_0^T E(|X(s) - X^{(n)}(s)|^2) ds \leq \frac{1}{n^6}.$$

Let  $Y^{(n)}(t) = \int_0^t X^{(n)}(s)dB(s)$ ,  $Y(t) = \int_0^t X(s)dB(s)$ . Then by Lemma 3.2.4,

$$P\left(\sup_{0 \leq t \leq T} |Y^{(m)}(t) - Y^{(n)}(t)| \geq \frac{1}{n}\right) \leq nE(|Y^{(m)}(T) - Y^{(n)}(T)|).$$

Let  $m \rightarrow \infty$ , we have

$$\begin{aligned} P\left(\sup_{0 \leq t \leq T} |Y(t) - Y^{(n)}(t)| \geq \frac{1}{n}\right) &\leq nE(|Y(T) - Y^{(n)}(T)|) \\ &\leq n \left(E(|Y(T) - Y^{(n)}(T)|^2)\right)^{1/2} \\ &\leq n \left(T \int_0^T E(|X(s) - X^{(n)}(s)|^2)ds\right)^{1/2} \\ &\leq n \frac{T^{1/2}}{n^3} = \frac{T^{1/2}}{n^2}. \end{aligned}$$

Since  $\sum n^{-2} < \infty$ . By the Borel-Cantelli lemma, we have

$$P\left(\sup_{0 \leq t \leq T} |Y(t) - Y^n(t)| \geq \frac{1}{n} \text{ i.o.}\right) = 0.$$

Let  $A$  be the complement of the above set. Then  $P(A) = 1$ , and let  $B$  be the set of  $\omega$  such that  $Y^{(n)}(\cdot, \omega)$  is continuous for all  $n$ . Then  $P(A \cap B) = 1$ , and for each  $\omega \in A \cap B$ ,

$$\sup_{0 \leq t \leq T} |Y(t, \omega) - Y^{(n)}(t, \omega)| \leq \frac{1}{n}$$

except for finitely many  $n$ . This implies that  $Y^n(\cdot, \omega) \rightarrow Y(\cdot, \omega)$  uniformly. Hence  $Y(\cdot, \omega)$  is continuous.  $\square$

In the rest of this section, we consider two important extension of the Ito integral. The first one is to extend the integrand to a larger class of processes.

We use  $\mathcal{L}_{ad}(\Omega, L^2[0, T])$  to denote the class of processes  $\{X(t)\}_{t \geq 0}$  such that

- (i)  $X(t)$  is adapted to  $\{\mathcal{F}_t\}_{t \geq 0}$ ;
- (ii)  $\int_0^T |X(t, \omega)|^2 dt < \infty$  a.s.

It is clear that if  $X(\cdot, \omega)$  has continuous sample path, then (ii) is satisfied. Also

$$L^2_{ad}([0, T] \times \Omega) \subset \mathcal{L}_{ad}(\Omega, L^2[0, T])$$



as  $X \in L^2_{ad}([0, T] \times \Omega)$ ,  $\int_0^T E(|X(t)|^2) dt < \infty$ , which implies  $\int_0^T |X(t, \omega)|^2 dt < \infty$  a.e.

**Example 3.2.3.** Consider  $X(t) = e^{B(t)^2}$ . Then a direct calculation shows that

$$E(X(t)) = \begin{cases} (1 - 4t)^{-1/2} & \text{if } 0 \leq t \leq \frac{1}{4}. \\ \infty & \text{otherwise} \end{cases}$$

Hence  $X \notin L^2_{ad}([0, T] \times \Omega)$ . On the other hand  $X(\cdot, \omega)$  is continuous a.e.,  $\int_0^1 |X(t, \omega)|^2 dt < \infty$  a.e., so that  $X \in \mathcal{L}_{ad}(\Omega, L^2[0, T])$ .

We can define the Ito integral as before using the following lemma.

**Lemma 3.2.7.** *Let  $X \in \mathcal{L}_{ab}(\Omega, L^2[0, T])$ . Then there exists a sequence of step process  $X^{(n)} \in L^2_{ab}([0, T] \times \Omega)$  such that*

$$\lim_{n \rightarrow \infty} \int_0^T |X^{(n)}(t) - X(t)|^2 dt = 0$$

*in probability.*

The sequence  $X^{(n)}(t)$  can be taken as the truncation of  $X(t)$  at level  $n$ :

$$X^{(n)}(t, \omega) = \begin{cases} X(t, \omega) & \text{if } \int_0^t |X(s, \omega)|^2 ds \leq n; \\ 0 & \text{otherwise.} \end{cases}$$

The resulting integral  $Y(t) = \int_0^t X(t) dB(t)$  may not have finite expectation. Hence is not necessary integrable, and we cannot consider the martingale property of  $Y(t)$  directly. Nevertheless, we can replace by the following *local martingale*.

Let  $\tau_n$  be the stoping time defined by

$$\tau_n(\omega) = \inf\{t : \int_0^t |X(s, \omega)|^2 ds > n\}$$

and let

$$Y(t \wedge \tau_n) = \int_0^{t \wedge \tau_n} X(t, \omega) dB(t) = \int_0^T X^{(n)} dB(t).$$

It is direct to show that for each  $n$ ,  $Y(t \wedge \tau_n)$  is a martingale with respect to  $\{\mathcal{F}_t\}$ . We say that  $Y(t)$  has the *local martingale property*. By using this we can show that  $Y(t, \omega)$  has continuous sample path for almost all  $\omega \in \Omega$ .

Next we consider extending the "integrator" to more general processes. A motivation of this is the following example. Consider  $X(t) \in L_{ad}^2([0, T] \times \Omega)$  and let

$$Y(t) = \int_0^t X(t) dB(t)$$

be the Ito integral. For convenience we can write it as

$$dY(t) = X(t) dB(t).$$

Then it is reasonable to have a definition so that

$$\int_0^T Z(t) dY(t) = \int_0^T Z(t) X(t) dB(t).$$

Note that  $Y(t)$  is a martingale (Theorem 3.2.4). The Ito integral theory can be extended to  $\int_0^T X(t) dM(t)$  where  $M(t)$  is a martingale with respect to some filtration  $\{\mathcal{F}_t\}$ . In this extension,  $M(t)$  is not necessary continuous, for example it covers the Poisson processes also.

For this the Doob-Meyer decomposition of submartingale plays an important role. For a submartingale  $L(t)$ , the decomposition is

$$L(t) = Z(t) + C(t)$$

where  $Z_t$  is a martingale, and  $C(t)$  is a non-negative increasing process. (Recall we have proved Doob's decomposition for discrete time (Theorem 1.4.3).) A basic application of this is on the submartingale  $M(t)^2$ . The decomposition is  $M(t)^2 = Z(t) + C(t)$  which yields

$$E((\Delta M_i)^2) = E(\Delta C_i)$$

where  $\Delta M_i = M(t_{i+1}) - M(t_i)$  corresponding to some partition. We can define Ito integral  $\int_0^T X(t)dM(t)$  similar to the Brownian motion case; and  $C(t)$  plays the role of  $t$  as the quadratic variation of  $B(t)$  :

$$(\Delta M_i)^2 = (M_{i+1} - M_i)^2 = M_{i+1}^2 - M_i^2 - 2M_i(M_{i+1} - M_i).$$

Hence

$$\begin{aligned} E((\Delta M_i)^2) &= E((M_{i+1} - M_i)^2) \\ &= E\left(E(M_{i+1}^2 - M_i^2 - 2M_i(M_{i+1} - M_i)/\mathcal{F}_{t_i})\right) \\ &= E\left(E(Z(t_{i+1}) + C(t_{i+1}) - (Z(t_i) + C(t_i)))/\mathcal{F}_{t_i}\right) \\ &= E(C(t_{i+1}) - C(t_i)) \\ &= E(\Delta C_i). \end{aligned}$$

The reader can refer to [3, p.75-92] for detail.

**Exercises**

1. Let  $X = \int_a^b |B(t)| dB(t)$ . Find the variance of the random variable  $X$ .
2. Show that  $X_t = e^{B(t)} - 1 - \frac{1}{2} \int_0^t e^{B(s)} ds$  is a martingale.
3. Show that  $X(t) = \int_0^t e^{B(s)^2} dB(s)$  is not a martingale.
4. Let  $M(t) = B^3(t) - 3tB(t)$ . Show that  $M(t)$  is a martingale, first directly then by using Ito integral.
5. Let  $B_1(t), B_2(t)$  be independent Brownian motions. Let  $X(t) = \log(B_1(t)^2 + B_2(t)^2)$ .
  - (a) Show that  $E(|X(t)|) < \infty$  for all  $t > 0$ , and find  $E(|X(t)|)$ .
  - (b) Show that  $X(t)$  is not a martingale, but a local martingale.

(The example shows that a local martingale having integrability does not need to be a martingale.)
6. Let  $N(t)$  be the Poisson process. Show that  $\tilde{N}(t) = N(t) - \lambda t$  is a martingale. Also find the Doob-Meyer decomposition for  $\tilde{N}(t)^2$ .