

Chapter 4

The Divergence Theorem

In this chapter we discuss formulas that connects different integrals. They are

- (a) Green's theorem that relates the line integral of a vector field along a plane curve to a certain double integral in the region it encloses.
- (b) Stokes' theorem that relates the line integral of a vector field along a space curve to a certain surface integral which is bounded by this curve.
- (c) Gauss' theorem that relates the surface integral of a closed surface in space to a triple integral over the region enclosed by this surface.

All these formulas can be unified into a single one called the divergence theorem in terms of differential forms.

4.1 Green's Theorem

Recall that the fundamental theorem of calculus states that

$$\int_a^b f'(x) dx = f(b) - f(a) .$$

It relates the integral of the derivative of a function over an interval $[a, b]$ to the endpoint values of the function. In higher dimension we replace the function by a vector field. A possible two dimensional extension would be a formula relating the double integral of some quantity involving the partial derivatives of a vector field to the line integral of the vector field along its boundary curve. This is the content of Green's theorem.

Theorem 4.1. (Green's Theorem) Let $\mathbf{F} = M\mathbf{i} + N\mathbf{j}$ be a C^1 -vector field in an open set G in the plane. Suppose C is a simple, closed curve in G and the set D it bounds lies completely inside G . Then

$$\iint_D \left(\frac{\partial N}{\partial x} - \frac{\partial M}{\partial y} \right) dA = \oint_C M dx + N dy , \quad (4.1)$$

where C is oriented in the anticlockwise way.

A simple, closed curve divides the plane into two regions, one bounded and the other unbounded. Here C bounds D means D is the bounded region.

Recall that line integral

$$\oint_C M dx + N dy = \oint_C \mathbf{F} \cdot d\mathbf{r} ,$$

is called the *circulation* of \mathbf{F} around the closed curve C . When \mathbf{F} is the velocity vector field of some fluid, its circulation around a curve measures the amount of the fluid flowing around the curve in unit time. When an admissible parametrization $\mathbf{r} : [a, b] \mapsto C$ is chosen, the line integral can be evaluated by the formula

$$\oint_C M dx + N dy = \int_a^b \mathbf{F}(\mathbf{r}(t)) \cdot \mathbf{r}'(t) dt .$$

Proof. We will prove Green's theorem in a special case, namely, D can be expressed simultaneously in the following two ways:

$$D = \{(x, y) : f_1(x) \leq y \leq f_2(x), x \in [a, b]\}$$

and

$$D = \{(x, y) : g_1(y) \leq x \leq g_2(y), y \in [c, d]\} .$$

Typical examples of such regions include ellipses and rectangles.

We shall show that

$$\iint_D \frac{\partial M}{\partial y} dA = - \oint_C M dx , \quad (4.2)$$

and

$$\iint_D \frac{\partial N}{\partial x} dA = \oint_C N dy . \quad (4.3)$$

Green's theorem follows by adding (4.2) and (4.3) together.

The boundary curve C of D consists of the four curves:

$$C_1 : \mathbf{r}_1(x) = (x, f_1(x)), \quad x \in [a, b] ,$$

$$C_2 : \mathbf{r}_2(x) = (x, f_2(x)), \quad x \in [a, b] ,$$

$$\gamma_2 : \gamma_2(y) = (b, y), \quad y \in [f_1(b), f_2(b)] ,$$

$$\gamma_1 : \gamma_1(y) = (a, y), \quad y \in [f_1(a), f_2(b)] ,$$

where γ_1 and γ_2 may degenerate into points. We have $C = C_1 + \gamma_2 - C_2 - \gamma_1$.

By Fubini's theorem

$$\begin{aligned} \iint_D \frac{\partial M}{\partial y} dA &= \int_a^b \int_{f_1(x)}^{f_2(x)} \frac{\partial M}{\partial y}(x, y) dy dx \\ &= \int_a^b M(x, y) \Big|_{f_1(x)}^{f_2(x)} dx \\ &= \int_a^b M(x, f_2(x)) - M(x, f_1(x)) dx \\ &= - \int_{C_1 - C_2} M dx . \end{aligned}$$

On the other hand, $\gamma_1'(y) = (0, 1)$, so

$$\int_{\gamma_1} M dx = \int_{f_1(a)}^{f_2(a)} M(a, y)x'(y) dy = 0,$$

as $x'(y) \equiv 0$. By the same reasoning

$$\int_{\gamma_2} M dx = 0 ,$$

too. Therefore,

$$\iint_D \frac{\partial M}{\partial y} dA = - \int_{C_1 - C_2} M dx = \oint_{C_1 + \gamma_2 - C_2 - \gamma_1} M dx = - \oint_C M dx ,$$

and (4.1) follows. Similarly, we can prove (4.2).

When the region D is of more complicated geometry, one can use horizontal and vertical lines to decompose it into the union of regions of the above types. We will not go into the details. \square

We will discuss four applications of Green's theorem:

- Evaluation of line integrals,
- Study independence of path,
- An area formula,
- Localizing divergence and rotation.

The first application is illustrated in the following example.

Example 4.1. Evaluate

$$\oint_C -y^2 dx + xy dy ,$$

where C is the boundary of the square at $(0, 0)$, $(1, 0)$, $(1, 1)$ and $(0, 1)$ in anticlockwise direction.

A direct evaluation is not difficult, but tedious as it involves evaluating four line integrals. Instead we take advantage of Green's theorem

$$\begin{aligned} \oint_C -y^2 dx + xy dy &= \iint_R \left(\frac{\partial xy}{\partial x} - \frac{\partial -y^2}{\partial y} \right) \\ &= \iint_R 3y dA(x, y) \\ &= \int_0^1 \int_0^1 3y dx dy \\ &= \frac{3}{2} . \end{aligned}$$

Next, we return to the discussion on independence of path of vector fields in Chapter 3. We established Theorem 3.4 which asserts that a vector field in \mathbb{R}^n is conservative if and only if the compatibility condition (3.9) holds (when $n = 2$). Now, by using Green's theorem, we will present a more general result.

A region in \mathbb{R}^n is called *simply connected* if it is connected and every closed curve lying in it can be deformed continuously to a point inside the set itself. The entire plane, a disk, a convex set and more general a star-shaped region are examples of simply connected sets in the plane. On the other hand, a punctured disk (a disk with the center removed) and an annulus are examples of connected but not simply-connected regions. Roughly speaking, simply connected regions are those connected regions which do not enclose any holes.

Theorem 4.2. *Let $\mathbf{F} = M\mathbf{i} + N\mathbf{j}$ be a C^1 -vector field in a simply connected region G in the plane. It is conservative if and only if the compatibility condition holds:*

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x} . \quad (4.4)$$

This generalizes Theorem 3.4 where it is required the vector field to be defined in the entire space.

Proof. In Chapter 3, it was shown that (4.4) (that is (3.9)) holds when the vector field \mathbf{F} is conservative. Conversely, under (4.4) a potential function were constructed under

the assumption that the line integrals along all simple curves connecting two points have the same value. It suffices to verify this property in the present situation. Let γ_1 and γ_2 be two simple curves connecting point A to point B . When these two curves do not intersect, $\gamma \equiv \gamma_1 - \gamma_2$ forms a simple closed curve. Green's theorem implies that

$$\oint_{\gamma} M dx + N dy = 0 ,$$

hence

$$\int_{\gamma_1} M dx + N dy = \int_{\gamma_2} M dx + N dy .$$

When γ_1 and γ_2 intersect, we may add another curve γ_3 connecting A and B so that it does not intersect γ_1 and γ_2 . Thus $\gamma_1 - \gamma_3$ and $\gamma_2 - \gamma_3$ form simple closed curves respectively. Using

$$\int_{\gamma_1} M dx + N dy = \int_{\gamma_3} M dx + N dy = \int_{\gamma_2} M dx + N dy ,$$

we draw the same conclusion. Using this property one can define the potential of \mathbf{F} as in the proof of Theorem 3.4. The existence of a potential Φ shows that

$$\int_A^B \mathbf{F} \cdot \mathbf{t} ds = \Phi(B) - \Phi(A) ,$$

along any path from A to B in G no matter the path is simple or not. We conclude that \mathbf{F} is conservative. \square

Green's theorem is a formula relating the line integral of a curve to a double integral of the region it encloses. This observation leads to a formula expressing the area of the region in terms of a boundary integral.

Let A be the area of the region enclosed by a simple closed curve C in the plane. Applying Green's theorem to the vector field $y\mathbf{i}$ yields

$$A = - \oint_C y dx . \tag{4.5}$$

Similarly, choosing the vector field to be $x\mathbf{j}$ yields

$$A = \oint_C x dy . \tag{4.6}$$

These two formulas together implies a more symmetric formula for the area:

$$A = \frac{1}{2} \oint_C -y dx + x dy . \tag{4.7}$$

Example 4.2. Find the area of one leaf of the four-leaf rose $r = 3 \sin 2\theta$.

The leaf in the first quadrant is ranged over $\theta \in [0, \pi/2]$. In general, when a curve is parametrized by $r(\theta)$ in polar coordinates, it is $r(\theta) \cos \theta \mathbf{i} + r(\theta) \sin \theta \mathbf{j}$ in cartesian coordinates. We have

$$\begin{aligned} & -y dx + x dy \\ &= (-r(\theta) \sin \theta)(r'(\theta) \cos \theta - r(\theta) \sin \theta) + (r(\theta) \cos \theta)(r'(\theta) \sin \theta + r(\theta) \cos \theta) d\theta \\ &= r^2(\theta) d\theta . \end{aligned}$$

Using the area formula, the area of one leaf is

$$\frac{1}{2} \int_0^{\pi/2} 9 \sin^2 2\theta d\theta = \frac{9\pi}{8} .$$

Example 4.3. Find the area of the loop in the folium of Decartes given by the zeros of the equation $x^3 + y^3 = 3xy$.

We first introduce a parametrization of the curve. Letting $y = tx$, we get

$$x = \frac{3t}{1+t^2} , \quad y = \frac{3t^2}{1+t^3} .$$

One can verify that the loop is described by $t \in [0, \infty)$. As t runs from 0 to ∞ , the loop runs in the positive direction. You may look up Wiki for more information concerning this curve.

We have

$$-y dx = \frac{-3t^2}{1+t^3} \frac{3[(1+t^3) - 3t^3]}{(1+t^3)^2} = \frac{-9t^2(1-2t^2)}{(1+t^3)^3} .$$

Therefore, the area of the loop is given by

$$\begin{aligned} \oint_C -y dx &= \int_0^\infty \frac{-9t^2(1-2t^2)}{(1+t^3)^3} dt \\ &= -3 \int_0^\infty \frac{1-2z}{(1+z)^3} dz \\ &= \frac{3}{2} . \end{aligned}$$

These formulas express the area enclosed by a curve in terms of the curve. It has interesting geometric consequence. For instance, together with Fourier series, the last formula can be used to prove the classical isoperimetric inequality, that is, among all regions enclosed by a simple closed curve with the same perimeter, only the disk has the largest area.

Finally, recall that in Chapter 3 we introduce the concept of the circulation and the flux of a vector along a curve. Let $\mathbf{F} = M\mathbf{i} + N\mathbf{j}$ be C^1 -vector field defined on the simple

closed oriented curve C with the chosen tangent \mathbf{t} and normal \mathbf{n} . The circulation and the flux of \mathbf{F} around C is defined to

$$\oint_C M dx + N dy ,$$

and

$$\oint_C M dy - N dx ,$$

respectively. Green's theorem suggests a way to define the circulation and the flux of a vector field at a point. In other words, we can localize circulation and flux.

Let \mathbf{p} be a point in some open set $G \subset \mathbb{R}^2$ where a C^1 -vector field $\mathbf{F} = P\mathbf{i} + Q\mathbf{j}$ is defined. Let C be a simple, closed curve anticlockwise oriented enclosing \mathbf{p} in its interior, and D the region it bounds. The quantity

$$\begin{aligned} \frac{1}{|D|} \oint_C M dx + N dy &= \frac{1}{|D|} \iint_D \left(\frac{\partial N}{\partial x} - \frac{\partial M}{\partial y} \right) dA \\ &\rightarrow \frac{\partial N}{\partial x}(\mathbf{p}) - \frac{\partial M}{\partial y}(\mathbf{p}) . \end{aligned}$$

In view of this, we define the *curl* (or the *rotation*) of \mathbf{F} at \mathbf{p} to be

$$\text{rot } \mathbf{F}(\mathbf{p}) = \left(\frac{\partial N}{\partial x} - \frac{\partial Q}{\partial y} \right) (\mathbf{p}) .$$

Similarly, the flux of \mathbf{F} across C is equal to

$$\oint_C M dy - N dx .$$

By Green's theorem,

$$\begin{aligned} \frac{1}{|D|} \oint_C M dy - N dx &= \frac{1}{|D|} \iint_D \left(\frac{\partial M}{\partial x} + \frac{\partial N}{\partial y} \right) dA \\ &\rightarrow \frac{\partial M}{\partial x}(\mathbf{p}) + \frac{\partial N}{\partial y}(\mathbf{p}) . \end{aligned}$$

Hence, we define the *divergence* (or *flux density*) of \mathbf{F} at \mathbf{p} to be

$$\text{div } \mathbf{F}(\mathbf{p}) = \left(\frac{\partial M}{\partial x} + \frac{\partial N}{\partial y} \right) (\mathbf{p}) .$$

Example 4.4. Evaluate

$$\oint_C \frac{y}{x^2 + y^2} dx + \frac{-x}{x^2 + y^2} dy ,$$

where C is the ellipse $x^2/4 + y^2/9 = 1$ oriented in positive direction.

By a direct computation, the vector field

$$\frac{y}{x^2 + y^2} \mathbf{i} + \frac{-x}{x^2 + y^2} \mathbf{j}$$

satisfies $M_y = N_x$. See the end of Section 3.6 in Chapter 3. However, since it is not defined at the origin, we cannot appeal to Green's theorem to conclude that this line integral vanishes. What we could do is to change it to an easier line integral. In fact, let C_r be a small circle centered at the origin and is contained inside C . We orient C_r in clockwise direction and connect C_r to C by the line segment L which runs from $(r, 0)$ to $(2, 0)$. Then $\Gamma = C - L + C_r + L$ forms a closed curve enclosing a simply-connected domain. Γ is not simple, but we can lift $\pm L$ up a little bit to make it simple. Applying Green's Theorem to this simple, closed curve and then passing to limit, we have

$$\begin{aligned} 0 &= \oint_{\Gamma} M dx + N dy \\ &= \left(\int_C - \int_L + \int_{C_r} + \int_L \right) M dx + N dy \\ &= \left(\int_C + \int_{C_r} \right) M dx + N dy . \end{aligned}$$

Therefore,

$$\begin{aligned} \int_C \frac{y}{x^2 + y^2} dx + \frac{-x}{x^2 + y^2} dy &= - \int_{C_r} \frac{y}{x^2 + y^2} dx + \frac{-x}{x^2 + y^2} dy \\ &= \int_0^{2\pi} (\cos \theta \cos \theta + (-\sin \theta)(-\sin \theta)) dt \\ &= 2\pi . \end{aligned}$$

The trick of adding an artificial line segment to form a simply connected region in this example leads us to the general form of Green's Theorem. Let D be a region bounded by several simple, closed curves C_1, C_2, \dots, C_n where C_1 is the outer one and the rest are sitting inside C_1 .

Theorem 4.3. *Let $\mathbf{F} = P\mathbf{i} + Q\mathbf{j}$ be a C^1 -vector field in D . Then*

$$\iint_D \left(\frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y} \right) dA = \sum_{j=1}^n \oint_{C_j} P dx + Q dy ,$$

where C_1 is oriented in anticlockwise way and $C_j, j \geq 2$, are in clockwise way.

4.2 Stokes' Theorem

Stokes' theorem is a curved version of Green's theorem, where the two dimensional region is replaced by a surface in space. Let $\mathbf{F} = M\mathbf{i} + N\mathbf{j} + P\mathbf{k}$ be a C^1 -vector field in some open set G in space. The curl of the vector field \mathbf{F} is given by

$$\operatorname{curl} \mathbf{F} = \left(\frac{\partial P}{\partial y} - \frac{\partial N}{\partial z} \right) \mathbf{i} + \left(-\frac{\partial P}{\partial x} + \frac{\partial M}{\partial z} \right) \mathbf{j} + \left(\frac{\partial N}{\partial x} - \frac{\partial M}{\partial y} \right) \mathbf{k}.$$

As \mathbf{F} is C^1 , the curl of \mathbf{F} is a continuous vector field in G . When the vector field is a planar one, that is, $M = M(x, y)$, $N = N(x, y)$ and $P \equiv 0$, the curl of \mathbf{F} becomes

$$\left(\frac{\partial N}{\partial x} - \frac{\partial M}{\partial y} \right) \mathbf{k},$$

the term appearing in Green's theorem.

We often use the notation $\nabla \times \mathbf{F}$ to denote the curl of \mathbf{F} as suggested by the formal expression by determinant

$$\begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ \partial_x & \partial_y & \partial_z \\ M & N & P \end{vmatrix}.$$

Let S be an oriented surface in space whose boundary is a curve C . An orientation assigned to C is based on the right hand rule: Let \mathbf{n} be the unit normal of S and \mathbf{b} be the unit vector at C which is tangential and pointing inward to S and perpendicular to C . Then the unit tangent of C \mathbf{t} is chosen such that $\mathbf{t} \times \mathbf{b}$ points to the same direction of \mathbf{n} . When a person is walking along the boundary in the positive direction (\mathbf{t} -direction) with his/her head pointing upward \mathbf{n} , the surface should be lying on his left hand side.

Theorem 4.4. (Stokes' Theorem) *Let S be an oriented surface in space whose boundary is a simple, closed curve C . For a C^1 -vector field \mathbf{F} on S ,*

$$\iint_S \nabla \times \mathbf{F} \cdot \mathbf{n} d\sigma = \oint_C \mathbf{F} \cdot d\mathbf{r},$$

where the orientation of C is described as above.

Proof. In the following it is more convenient to use (x, y, z) and (M, N, P) instead of $x\mathbf{i} + y\mathbf{j} + z\mathbf{k}$ and $M\mathbf{i} + N\mathbf{j} + P\mathbf{k}$.

Let $\mathbf{r} : D \rightarrow S$ be a parametrization of S so that $\mathbf{r}_u \times \mathbf{r}_v$ points to the normal direction of S . It can be shown that the anticlockwise direction of the boundary curve C_1 of D is mapped to C with the same orientation of C . Let

$$\gamma(t) = (u(t), v(t)) \quad t \in [a, b],$$

be a parametrization for C_1 . Letting $\mathbf{r}(u, v) = (x(u, v), y(u, v), z(u, v))$, the map

$$t \mapsto (x(u(t), v(t)), y(u(t), v(t)), z(u(t), v(t))) , \quad t \in [a, b] ,$$

gives a parametrization of the curve C . Using

$$dx = (x_u u' + x_v v') dt, \quad dy = (y_u u' + y_v v') dt, \quad dz = (z_u u' + z_v v') dt ,$$

the left hand side of the Stokes' theorem is

$$\begin{aligned} & \oint_C \mathbf{F} \cdot d\mathbf{r} \\ &= \int_a^b M(x_u u' + x_v v') + N(y_u u' + y_v v') + P(z_u u' + z_v v') dt \\ &= \int_a^b [(Mx_u + Ny_u + Pz_u)u' + (Mx_v + Ny_v + Pz_v)v'] dt \\ &\equiv \int_a^b (Au' + Bv') dt , \end{aligned}$$

where

$$\begin{aligned} & A(u, v) \\ &= M(x(u, v), y(u, v), z(u, v))x_u(u, v) + N(x(u, v), y(u, v), z(u, v))y_u(u, v) + P(x(u, v), y(u, v), z(u, v))z_u(u, v) \end{aligned}$$

and

$$\begin{aligned} & B(u, v) \\ &= M(x(u, v), y(u, v), z(u, v))x_v(u, v) + N(x(u, v), y(u, v), z(u, v))y_v(u, v) + P(x(u, v), y(u, v), z(u, v))z_v(u, v) \end{aligned}$$

Appealing to Green's theorem, the left hand side of the Stokes' formula is equal to

$$\begin{aligned} & \int_a^b (Au' + Bv') dt \\ &= \oint_{C_1} (A, B) \cdot d\gamma \\ &= \iint_D (B_u - A_v) dA(u, v) \\ &= \iint_D [(M_x x_u + M_y y_u + M_z z_u)x_v + Mx_{vu} + (N_x x_u + N_y y_u + N_z z_u)y_v + Ny_{vu} \\ &\quad + (P_x x_u + P_y y_u + P_z z_u)z_v + Pz_{vu} - (M_x x_v + M_y y_v + M_z z_v)x_u - Mx_{uv} \\ &\quad - (N_x x_v + N_y y_v + N_z z_v)y_u - Ny_{uv} - (P_x x_v + P_y y_v + P_z z_v)z_u - Pz_{vu}] dA(u, v) \\ &= \iint_D [(N_x - M_y)(x_u y_v - x_v y_u) + (P_x - M_z)(x_u z_v - x_v z_u) + (P_y - N_z)(y_u z_v - y_v z_u)] dA . \end{aligned}$$

On the other hand, $(u, v) \mapsto \mathbf{r} = (x(u, v), y(u, v), z(u, v))$ parametrizes S by D . We have

$$\mathbf{r}_u \times \mathbf{r}_v = (y_u z_v - y_v z_u, -x_u z_v + x_v z_u, x_u y_v - x_v y_u) .$$

The right hand side of the formula is

$$\begin{aligned} & \iint_S \nabla \times \mathbf{F} \cdot \mathbf{n} \, d\sigma \\ &= \iint_D (P_y - N_z, -P_x + M_z, N_x - M_y) \cdot \mathbf{r}_u \times \mathbf{r}_v \, dA(u, v) \\ &= \iint_D [(P_y - N_z)(y_u z_v - y_v z_u) + (-P_x + M_z)(-x_u z_v + x_v z_u) + (N_x - M_y)(x_u y_v - x_v y_u)] \, dA(u, v) . \end{aligned}$$

By comparing the left and the right hand sides of Stokes' formula, we conclude that the theorem holds.

□

Example 4.5. Evaluate the line integral

$$\oint_C (y + z) \, dx + (z + x) \, dy + x \, dz,$$

where C is the intersection of the cylinder $x^2 + y^2 = 1$ with the plane $x + y + 2z = 10$ oriented in the anticlockwise direction.

The direct way of evaluation is to write down a parametrization for the curve C . Indeed, the projection of this curve onto the xy -plane is the circle $x^2 + y^2 = 1$. Hence an admissible parametrization is

$$\theta \mapsto (\cos \theta, \sin \theta, (10 - \cos \theta - \sin \theta)/2) , \quad \theta \in [0, 2\pi] .$$

But, here we would like to apply Stokes' theorem. Observing that C is the boundary of the oriented surface given by the graph of $z = (10 - x - y)/2$ over the unit disk $D_1 : x^2 + y^2 \leq 1$ with normal pointing upward. For $\mathbf{F} = (y + z)\mathbf{i} + (z + x)\mathbf{j} + x\mathbf{k}$, we have $\nabla \times \mathbf{F} = -\mathbf{i}$. On the other hand, the upward normal is

$$\mathbf{n} = \frac{-\varphi_x \mathbf{i} - \varphi_y \mathbf{j} + \mathbf{k}}{\sqrt{1 + \varphi_x^2 + \varphi_y^2}} = \frac{\sqrt{6}}{6} (\mathbf{i} + \mathbf{j} + 2\mathbf{k}) .$$

By Stokes' theorem,

$$\begin{aligned}
 & \oint_C (y+z) dx + (z+x) dy + x dz \\
 &= \iint_S -\mathbf{i} \cdot \mathbf{n} d\sigma \\
 &= \frac{-\sqrt{6}}{6} \iint_S d\sigma \\
 &= -\iint_{D_1} \frac{1}{2} dA, \quad \left(d\sigma = (1 + |\nabla\varphi|^2) dA = \sqrt{\frac{3}{2}} dA \right) \\
 &= -\frac{\pi}{2}.
 \end{aligned}$$

A very useful consequence of Stokes' theorem is

Corollary 4.5. *Suppose that a simple closed curve C in space bounds two surfaces S_1 and S_2 so that $S = S_1 \cup S_2$ forms a closed surface whose unit outward normal coincides with the normals of S_1 and S_2 . For any C^1 -vector field defined in an open set containing S ,*

$$\iint_{S_2} \nabla \times \mathbf{F} \cdot \mathbf{n} d\sigma = -\iint_{S_1} \nabla \times \mathbf{F} \cdot \mathbf{n} d\sigma.$$

Proof. Let C_1 and C_2 be the respective oriented boundary curves for S_1 and S_2 . It is clear that $C_2 = -C_1$. By Stokes' theorem,

$$\begin{aligned}
 \iint_{S_2} \nabla \times \mathbf{F} \cdot \mathbf{n} d\sigma &= \oint_{C_2} \mathbf{F} \cdot d\mathbf{r} \\
 &= \oint_{-C_1} \mathbf{F} \cdot d\mathbf{r} \\
 &= -\oint_{C_1} \mathbf{F} \cdot d\mathbf{r} \\
 &= -\iint_{S_1} \nabla \times \mathbf{F} \cdot \mathbf{n} d\sigma.
 \end{aligned}$$

□

Example 4.6. Let Σ be the upper hemisphere $x^2 + y^2 + z^2 = 1, z \geq 0$, with normal pointing upward. Evaluate the integral

$$\iint_{\Sigma} \nabla \times \mathbf{F} \cdot \mathbf{n} d\sigma,$$

where $\mathbf{F} = y\mathbf{i} + x^2z\mathbf{j} + e^{xz}\mathbf{k}$.

First of all, we calculate

$$\nabla \times \mathbf{F} = -x^2 \mathbf{i} - ze^{xz} \mathbf{j} + (2xz - 1) \mathbf{k} .$$

Let S be the unit disk in the xy -plane. We regard it as a flat surface in space. With normal pointing down, Σ and S are two oriented surfaces sharing the same boundary. By the above corollary,

$$\begin{aligned} & \iint_{\Sigma} \nabla \times \mathbf{F} \cdot \mathbf{n} \, d\sigma \\ &= - \iint_S \nabla \times (y\mathbf{i} + x^2z\mathbf{j} + e^{xz}\mathbf{k}) \cdot -\mathbf{k} \, d\sigma \\ &= \iint_S (2xz - 1) \, d\sigma . \end{aligned}$$

An obvious parametrization of S is $(x, y) \mapsto (x, y, 0)$. Hence we continue to get

$$\iint_{\Sigma} \nabla \times (y\mathbf{i} + x^2z\mathbf{j} + e^{xz}\mathbf{k}) \cdot \mathbf{n} \, d\sigma = \iint_{D_1} -dA = -\pi .$$

Alternatively, we may use Stokes' theorem to replace it by a line integral. Indeed, the boundary curve is simply the unit circle in the xy -plane which admits the parametrization $C : \theta \mapsto (\cos \theta, \sin \theta)$, $\theta \in [0, 2\pi]$ in positive direction. By Stokes' theorem,

$$\begin{aligned} \iint_{\Sigma} \nabla \times \mathbf{F} \cdot \mathbf{n} \, d\sigma &= \oint_C \mathbf{F} \cdot d\mathbf{r} \\ &= \int_0^{2\pi} (\sin \theta \mathbf{i} + \mathbf{k}) \cdot (-\sin \theta \mathbf{i} + \cos \theta \mathbf{j}) \, d\theta \\ &= - \int_0^{2\pi} \sin^2 \theta \, d\theta \\ &= -\pi . \end{aligned}$$

Next, we examine the meaning of the curl vector $\nabla \times \mathbf{F}$. Let \mathbf{p} be a point in space and introduce a coordinates system using this point as the origin. Let ξ be a unit vector and P be the plane perpendicular to it. Consider a simple closed curve C on P which surrounds the origin. Letting its enclosed region be D , by Stokes' theorem,

$$\begin{aligned} & \frac{1}{|D|} \oint_C M \, dx + N \, dy + P \, dz \\ & \frac{1}{|D|} \iint_D \nabla \times \mathbf{F} \cdot \xi \, d\sigma \\ & \rightarrow \nabla \times \mathbf{F}(\mathbf{p}) \cdot \xi , \end{aligned}$$

as C shrinks to \mathbf{p} . Hence the term $\nabla \times \mathbf{F}(\mathbf{p}) \cdot \xi$ measures the strength of rotation of the vector field \mathbf{F} along the ξ -direction. This gives a meaning to the curl vector. From

$$\nabla \times \mathbf{F} \cdot \xi = |\nabla \times \mathbf{F}| |\xi| \cos \theta ,$$

where $\theta \in [0, \pi]$ is the angle between the curl vector and ξ , we see that the strength of rotation is maximal when ξ is equal to $\nabla \times \mathbf{F}$ or $-\nabla \times \mathbf{F}$.

Vector fields that are gradients of functions constitute a large class of irrotational vector fields.

Theorem 4.6. *Let Φ be C^2 -function in some open set G in space. Then $\mathbf{F} = \nabla\Phi$ is irrotational.*

Proof. When \mathbf{F} is the gradient of Φ , $M = \Phi_x, N = \Phi_y, P = \Phi_z$. Then

$$M_y - N_x = \Phi_{xy} - \Phi_{yx} = 0 .$$

Similarly all other equations in the compatibility conditions hold. □

In Chapter 3, it was shown that a conservative C^1 -vector field $\mathbf{F} = M\mathbf{i} + N\mathbf{j} + P\mathbf{k}$ must satisfy the compatibility condition

$$\frac{\partial N}{\partial x} = \frac{\partial P}{\partial z}, \quad \frac{\partial M}{\partial z} = \frac{\partial P}{\partial x}, \quad \frac{\partial M}{\partial y} = \frac{\partial N}{\partial x} .$$

In other words, a conservative vector field must be irrotational. Now we show that any conservative vector field arises in this way when it lies in a simply connected region. As a consequence of Green's theorem, we have seen this when the vector field is in the plane.

Theorem 4.7. *An irrotational C^1 -vector field \mathbf{F} in some simply connected region G is conservative.*

Proof. It suffices to show that the line integral of \mathbf{F} along any simple closed curve in G vanishes. Let C be a simple closed curve in the region G . Since G is simply connected, C can be deformed into a point inside G . The deformation itself forms an oriented surface S taking C as its boundary. By Stokes' theorem and the compatibility condition,

$$\oint_C \mathbf{F} \cdot d\mathbf{r} = \iint_S \nabla \times \mathbf{F} \cdot \mathbf{n} \, d\sigma = 0 .$$

□

To conclude this section, we consider oriented surfaces which are bounded not by a single but by finitely many nonintersecting simple, closed curves. Just like Theorem 4.3, using the trick of cutting-up one can establish the following general Stokes' theorem.

Theorem 4.8. Let \mathbf{F} be a C^1 -vector field in some open set G in space. Let S be a C^1 -surface in G which is bounded by finitely many nonintersecting simple, closed curves C_1, \dots, C_n . Then

$$\iint_S \nabla \times \mathbf{F} \cdot \mathbf{n} d\sigma = \sum_{j=1}^n \oint_{C_j} \mathbf{F} \cdot d\mathbf{r} .$$

4.3 Divergence Theorem

Recall that there are two different formulations for Green's theorem. Let $\mathbf{F} = M\mathbf{i} + N\mathbf{j}$ be a C^1 -vector field in some open set G and C a simple closed curve whose enclosed region D sits in G . Then

$$\iint_D \left(\frac{\partial N}{\partial x} - \frac{\partial M}{\partial y} \right) dA = \oint_C \mathbf{F} \cdot \mathbf{t} ds ,$$

where \mathbf{t} is the unit tangent along the anticlockwise direction. Alternatively, we can express the formula in the following way:

$$\iint_D \left(\frac{\partial M}{\partial x} + \frac{\partial N}{\partial y} \right) dA = \oint_C \mathbf{F} \cdot \mathbf{n} ds ,$$

where \mathbf{n} is the outward unit normal of C .

Stokes' theorem may be viewed as a generalised version of the tangent form of the Green's theorem where a plane region is replaced by a surface in space. The divergence theorem we are going to discuss may be regarded as a generalised version of the normal form where now a plane region is replaced by a region in space.

Theorem 4.9. Let $\mathbf{F} = M\mathbf{i} + N\mathbf{j} + P\mathbf{k}$ be a C^1 -vector field in an open set G in \mathbb{R}^3 . Let S be a closed C^1 -surface which is the boundary of a bounded region $\Omega \subset G$. Then

$$\iiint_{\Omega} \nabla \cdot \mathbf{F} dV = \iint_S \mathbf{F} \cdot \mathbf{n} d\sigma ,$$

where \mathbf{n} is the unit outward normal at S .

The divergence of a vector field, $\nabla \cdot \mathbf{F}$ is given by

$$\nabla \cdot \mathbf{F} = \frac{\partial M}{\partial x} + \frac{\partial N}{\partial y} + \frac{\partial P}{\partial z} .$$

Proof. We only consider the special case that Ω can be expressed simultaneously in the following three ways:

$$\{(x, y, z) : (x, y) \in D, f_1(x, y) \leq z \leq f_2(x, y)\},$$

$$\{(x, y, z) : (y, z) \in D, g_1(y, z) \leq x \leq g_2(y, z)\},$$

and

$$\{(x, y, z) : (x, z) \in D, h_1(x, z) \leq y \leq h_2(x, z)\},$$

where D is a region in the respective plane. For instance, rectangular boxes and ellipsoids admit such representations.

We will show that in the first case,

$$\iiint_{\Omega} \frac{\partial P}{\partial z} dV = \oiint_S P \mathbf{k} \cdot \mathbf{n} d\sigma. \quad (4.8)$$

A similar argument will establish

$$\iiint_{\Omega} \frac{\partial M}{\partial x} dV = \oiint_S M \mathbf{i} \cdot \mathbf{n} d\sigma,$$

and

$$\iiint_{\Omega} \frac{\partial N}{\partial y} dV + \oiint_S N \mathbf{j} \cdot \mathbf{n} d\sigma.$$

The divergence theorem follows by adding up these three identities.

Let us now prove (4.9). We may further assume that $S = S_1 \cup S_2$ where

$$S_1 = \{(x, y, f_1(x, y)) : (x, y) \in D\}, \quad S_2 = \{(x, y, f_2(x, y)) : (x, y) \in D\}.$$

Moreover, these two surfaces meet a curve whose projection to the xy -plane is a curve C bounding D . We write

$$\oiint_S P \mathbf{k} \cdot \mathbf{n} d\sigma = \iint_{S_1} P \mathbf{k} \cdot \mathbf{n} d\sigma + \iint_{S_2} P \mathbf{k} \cdot \mathbf{n} d\sigma.$$

As

$$\mathbf{n} = \frac{-f_{2x} \mathbf{i} - f_{2y} \mathbf{j} + \mathbf{k}}{\sqrt{1 + |\nabla f_2|^2}}$$

on S_2 ,

$$\iint_{S_2} P \mathbf{k} \cdot \mathbf{n} d\sigma = \iint_{S_2} \frac{P}{\sqrt{1 + |\nabla f_2|^2}} d\sigma = \iint_D P(x, y, f_2(x, y)) dA(x, y).$$

On the other hand,

$$\mathbf{n} = \frac{f_{1x} \mathbf{i} + f_{1y} \mathbf{j} - \mathbf{k}}{\sqrt{1 + |\nabla f_1|^2}}$$

on S_1 (noting that it points downward),

$$\iint_{S_1} P \mathbf{k} \cdot \mathbf{n} d\sigma = - \iint_D P(x, y, f_1(x, y)) dA(x, y).$$

Therefore,

$$\oiint_S P \mathbf{k} \cdot \mathbf{n} \, d\sigma = \iint_D (P(x, y, f_2(x, y)) - P(x, y, f_1(x, y))) \, dA(x, y) .$$

On the other hand,

$$\begin{aligned} & \iiint_\Omega \frac{\partial P}{\partial z} \, dV \\ &= \iiint_D \int_{f_1(x, y)}^{f_2(x, y)} \frac{\partial P}{\partial z} \, dz \, dA \\ &= \iint_D (P(x, y, f_2(x, y)) - P(x, y, f_1(x, y))) \, dA(x, y) , \end{aligned}$$

and the desired result follows. □

Example 4.7. Verify the divergence theorem for the vector field $\mathbf{F} = x\mathbf{i} + y\mathbf{j} + z\mathbf{k}$ and Ω is the ball B , $x^2 + y^2 + z^2 \leq a^2$.

The divergence of \mathbf{F} is

$$\nabla \cdot \mathbf{F} = \frac{\partial}{\partial x}x + \frac{\partial}{\partial y}y + \frac{\partial}{\partial z}z = 3 .$$

Therefore,

$$\iiint_B \nabla \cdot \mathbf{F} \, dV = \iiint_B 3 \, dV = 3 \times \frac{4}{3}\pi a^3 = 4\pi a^3 .$$

On the other hand, it is clear that the outward unit normal at a boundary point (x, y, z) is given by

$$\mathbf{n} = \frac{1}{a}(x\mathbf{i} + y\mathbf{j} + z\mathbf{k}) .$$

We have

$$\begin{aligned} \oiint_{\partial B} \mathbf{F} \cdot \mathbf{n} \, d\sigma &= \oiint_{\partial B} (x\mathbf{i} + y\mathbf{j} + z\mathbf{k}) \cdot \frac{1}{a}(x\mathbf{i} + y\mathbf{j} + z\mathbf{k}) \, d\sigma \\ &= \frac{1}{a} \times a^2 \oiint_{\partial B} d\sigma \\ &= 4\pi a^3 . \end{aligned}$$

Example 4.8. Find the flux of $\mathbf{F} = xy\mathbf{i} + yz\mathbf{j} + xz\mathbf{k}$ outward through the surface of the cube cut from the first octant by the planes $x = 1, y = 1$ and $z = 1$.

Let Ω be the solid cube and C its boundary. By the divergence theorem, the flux is equal to

$$\iint_C \mathbf{F} \cdot \mathbf{n} \, d\sigma = \iiint_\Omega \nabla \cdot \mathbf{F} \, dV .$$

It suffices to calculate the triple integral which is obviously easier. We have $\nabla \cdot \mathbf{F} = x + y + z$. Therefore, the flux is equal to

$$\iiint_{\Omega} (x + y + z) dV = \int_0^1 \int_0^1 \int_0^1 (x + y + z) dx dy dz = \frac{3}{2} .$$

The divergence of a vector field may be regarded as a mapping which maps a vector field to a function. Let \mathbf{p} be a point in some open set where a C^1 -vector field \mathbf{F} is defined. Consider a simple closed surface S which contains \mathbf{p} in its interior D . The outward flux of \mathbf{F} across S is given by

$$\oiint_S \mathbf{F} \cdot \mathbf{n} d\sigma ,$$

where \mathbf{n} is the unit outward normal. By the divergence theorem,

$$\begin{aligned} \frac{1}{|D|} \oiint_S \mathbf{F} \cdot \mathbf{n} d\sigma &= \frac{1}{|D|} \iiint_D \nabla \cdot \mathbf{F} dV \\ &\rightarrow \nabla \cdot \mathbf{F}(\mathbf{p}) , \end{aligned}$$

as S shrinks to \mathbf{p} . Hence $\nabla \cdot \mathbf{F}$ may be viewed as a measurement of the flux of \mathbf{F} at the point \mathbf{p} . When the vector field is two dimensional, its divergence reduces to the one discussed in the last chapter.

The possibility of localizing the divergence is of fundamental important in physics. As an illustration let us derive the equation of continuity. Image the space is occupied with some fluid which is described in terms of its density δ and velocity \mathbf{u} . Both are functions of (x, y, z, t) . We fix a region Ω in space and consider the mass of the fluid inside this region. As the density changes in time, the mass

$$M(t) = \iiint_{\Omega} \delta(x, y, z, t) dV ,$$

is also a function of time. The rate of change in mass is given by

$$\frac{d}{dt} M(t) = \iiint_{\Omega} \frac{\partial}{\partial t} \delta(x, y, z, t) dV . \quad (4.9)$$

On the other hand, the fluid that leaves or enters through a small piece $\Delta\sigma$ on the surface of Ω , S , in time Δt is given by

$$\mathbf{u} \cdot \mathbf{n} \Delta\sigma \Delta t ,$$

where \mathbf{n} is the unit outward normal. Hence the mass leaving or entering the region through $\Delta\sigma$ is equal to

$$\delta \mathbf{u} \cdot \mathbf{n} \Delta\sigma \Delta t .$$

It follows that the mass that leaving/entering S in instant time is equal to

$$\begin{aligned} \frac{d}{dt}M(t) &= \lim_{\Delta t \rightarrow 0} \frac{M(t + \Delta t) - M(t)}{\Delta t} \\ &= - \lim_{\Delta t \rightarrow 0} \frac{1}{\Delta t} \iiint_S \delta \mathbf{u} \cdot \mathbf{n} \Delta \sigma \Delta t \, d\sigma \\ &= - \oiint_S \delta \mathbf{u} \cdot \mathbf{n} \, d\sigma. \end{aligned}$$

Comparing with (4.9), we have

$$\iiint_{\Omega} \frac{d\delta}{dt}(x, y, z, t) \, dV = - \oiint_S \delta \mathbf{u} \cdot \mathbf{n} \, d\sigma .$$

Now, by the divergence theorem,

$$\iiint_{\Omega} \left(\frac{d\delta}{dt} + \nabla \cdot \delta \mathbf{u} \right) \, dV = 0 .$$

Since here Ω could be any region, one conclude that

$$\frac{d\delta}{dt} + \nabla \cdot \delta \mathbf{u} = 0 , \tag{4.10}$$

must hold. This equation, called the equation of continuity, relating the density to the velocity is a form of conservation of mass. Together with other equations such as conservation of energy, etc, one obtain a full system of partial differential equations for the density and the velocity of the fluid.

In the divergence theorem stated above, it is assumed that the boundary of the region is a single closed surface. When the boundary consists of several closed surfaces, we have the following general divergence theorem.

Theorem 4.10. *Let Ω be a region in space whose boundary is composed of finitely many closed C^1 -surfaces S_1, S_2, \dots, S_n . Then for any C^1 -vector field \mathbf{F} in Ω ,*

$$\iiint_{\Omega} \nabla \cdot \mathbf{F} \, dV = \sum_{j=1}^n \oiint_{S_j} \mathbf{F} \cdot \mathbf{n} \, d\sigma ,$$

where \mathbf{n} is the unit outward normal to S_j .

This theorem is the space version of the planar case Theorem 4.3. Its proof is similar by cutting up the region. Since it is inconvenient to draw pictures here, you are referred to the text for details.

A vector field is called *divergence-free* if $\mathbf{F} = 0$. The following result provides us with a large class of divergence vector fields.

Theorem 4.11. *The divergence of a C^1 -curl vector field always vanishes.*

Proof. Let \mathbf{F} be a curl, that is, $\mathbf{F} = \nabla \times \mathbf{H}$ for some vector field $\mathbf{H} = M\mathbf{i} + N\mathbf{j} + P\mathbf{k}$. Then

$$\begin{aligned}\nabla \cdot \mathbf{F} &= \nabla \cdot ((N_z - P_y)\mathbf{i} + (-M_z + P_x)\mathbf{j} + (M_y - N_x)\mathbf{k}) \\ &= (N_z - P_y)_x + (-M_z + P_x)_y + (M_y - N_x)_z \\ &= 0.\end{aligned}$$

□

Recall that the curl of a gradient vector field always vanishes, that is, $\nabla \times \nabla\Phi = \mathbf{0}$. Now, the divergence of a curl vector field always vanishes as well. In the former one asks, when a vector field \mathbf{F} satisfies $\nabla \times \mathbf{F} = \mathbf{0}$, is \mathbf{F} a gradient? The answer is yes when the underlying region is simply connected. Here one may ask a similar question. Namely, when $\nabla \cdot \mathbf{F} = 0$, is there another vector field \mathbf{A} such that $\mathbf{F} = \nabla \times \mathbf{A}$? It turns out the answer is yes when the region is star-shaped.

According to the divergence theorem, a divergence-free vector field \mathbf{F} satisfies

$$\oiint_S \mathbf{F} \cdot \mathbf{n} \, d\sigma = 0,$$

over any simple, closed surface S . In other words, its flux across any closed surface vanishes. A special case is the following theorem of Gauss which appears in the study of electricity.

The electric field generated by a point charge locating at the origin is given by

$$\mathbf{E} = \frac{q}{4\pi\epsilon_0} \frac{1}{r^3} (x\mathbf{i} + y\mathbf{j} + z\mathbf{k}), \quad r = \sqrt{x^2 + y^2 + z^2},$$

where q is the charge and ϵ_0 is some positive constant. The electric field is divergence free:

$$\nabla \cdot \mathbf{E} = \frac{q}{4\pi\epsilon_0} \left(\frac{y^2 + z^2 - 2x^2}{r^5} + \frac{x^2 + z^2 - 2y^2}{r^5} + \frac{y^2 + x^2 - 2z^2}{r^5} \right) = 0.$$

Theorem 4.12. *The flux of the electric field across any closed C^1 -surface enclosing the origin is the same and equal to q/ϵ_0 .*

Proof. Let Ω be the interior of S . Since the electric field is not well-defined at the origin, we cannot apply the divergence directly to Ω . Instead we consider the region Ω_r which is

bounded by S and the small sphere $\Sigma_r : x^2 + y^2 + z^2 = r^2$. Then the electric field is C^1 in Ω_r and is of divergence free. By the divergence theorem,

$$0 = \iiint_{\Omega_r} \nabla \cdot \mathbf{E} \, dV = \iint_S \mathbf{E} \cdot \mathbf{n} \, d\sigma + \iint_{\Sigma_r} \mathbf{E} \cdot \mathbf{n} \, d\sigma ,$$

and the flux across S is equal to

$$- \iint_{\Sigma_r} \mathbf{E} \cdot \mathbf{n} \, d\sigma .$$

When regarded as the inner boundary of Ω , the unit outer normal \mathbf{n} at Σ_r points toward the origin and is given by $-(x\mathbf{i} + y\mathbf{j} + z\mathbf{k})/r$. So,

$$\begin{aligned} - \iint_{\Sigma_r} \mathbf{E} \cdot \mathbf{n} \, d\sigma &= \frac{q}{4\pi\epsilon_0} \iint_{\Sigma_r} \frac{1}{r^3} (x\mathbf{i} + y\mathbf{j} + z\mathbf{k}) \cdot \frac{1}{r} (x\mathbf{i} + y\mathbf{j} + z\mathbf{k}) \, d\sigma \\ &= \frac{q}{4\pi\epsilon_0} \frac{1}{r^4} \iint_{\Sigma_r} (x^2 + y^2 + z^2) \, d\sigma \\ &= \frac{q}{\epsilon_0} . \end{aligned}$$

□