

TA: LEE, Yat Long Luca

Email: ylllee@math.cuhk.edu.hk

Office: Room 505, AB1

Office Hour: Send me an email first, then we will arrange a meeting (if you need it).

Tutorial Arrangement:

- (1330 - 1355/ 15:30 - 15:55): Problems.
- (1355 - 1415/ 15:55 - 16:15): Class exercises.
- (1415 - 1430/ 16:15 - 16:30): Submission of Class Exercise via Gradescope.
- (1430 - 1530/ 16:30 - 17:30): Late submission period.

1 Line Integrals of Vector Fields

1.1 Vector Fields

Vector fields: $\mathbf{F} : \mathbb{R}^n \longrightarrow \mathbb{R}^m$ given by $\mathbf{F}(x_1, \dots, x_n) = (F_1(x_1, \dots, x_n), \dots, F_m(x_1, \dots, x_n))$. In this course you will mostly see vector fields defined in $\mathbb{R}^2$ or $\mathbb{R}^3$, then you can write it as

$$\mathbf{F} = F_1\mathbf{i} + F_2\mathbf{j} + F_3\mathbf{k}$$

etc.

Example:

Gradient of a scalar function: Let $f : \mathbb{R}^3 \longrightarrow \mathbb{R}$ be a C^1 function, then

$$\nabla f = \left(\frac{\partial f}{\partial x}, \frac{\partial f}{\partial y}, \frac{\partial f}{\partial z} \right) = \frac{\partial f}{\partial x}\mathbf{i} + \frac{\partial f}{\partial y}\mathbf{j} + \frac{\partial f}{\partial z}\mathbf{k}$$

is a vector field, and is called a gradient field.

1.2 Line Integrals

Let C be a curve with parametrization $\mathbf{r}(t)$ that gives C the counterclockwise direction. Then the line integral of $\mathbf{F}$ over C is given by

$$\int_C \mathbf{F} \cdot d\mathbf{r}$$

Steps for calculation:

- (i) Substitute $\mathbf{r}(t)$ into $\mathbf{F}$, i.e., find $\mathbf{F}(\mathbf{r}(t))$.
- (ii) Differentiate $\mathbf{r}(t)$ and get $\mathbf{r}'(t)$.
- (iii) Evaluate the integral with respect to t , where $t \in [a, b]$, i.e.,

$$\int_C \mathbf{F} \cdot d\mathbf{r} = \int_a^b \mathbf{F}(\mathbf{r}(t)) \cdot \mathbf{r}'(t) dt$$

Exercise**Q1**

Evaluate $\int_C \mathbf{F} \cdot d\mathbf{r}$, where

$$\mathbf{F}(x, y, z) = z\mathbf{i} + xy\mathbf{j} - y^2\mathbf{k}$$

along the curve C given by

$$\mathbf{r}(t) = t^2\mathbf{i} + t\mathbf{j} + \sqrt{t}\mathbf{k},$$

for $t \in [0, 1]$

1.2.1 Line Integral with respect to dx, dy, dz

Let $\mathbf{F} : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ be the following vector field

$$\mathbf{F}(x, y, z) = F_1(x, y, z)\mathbf{i} + F_2(x, y, z)\mathbf{j} + F_3(x, y, z)\mathbf{k}$$

and C be the curve

$$\mathbf{r}(t) = r_1(t)\mathbf{i} + r_2(t)\mathbf{j} + r_3(t)\mathbf{k}$$

then

$$\int_C F_1(x, y, z) dx = \int_a^b F_1(\mathbf{r}(t))r'_1(t) dt,$$

$$\int_C F_2(x, y, z) dy = \int_a^b F_2(\mathbf{r}(t))r'_2(t) dt,$$

$$\int_C F_3(x, y, z) dz = \int_a^b F_3(\mathbf{r}(t))r'_3(t) dt.$$

To summarize,

$$\begin{aligned} \int_C \mathbf{F} \cdot d\mathbf{r} &= \int_C (F_1, F_2, F_3) \cdot (dx, dy, dz) \\ &= \int_C F_1 dx + \int_C F_2 dy + \int_C F_3 dz \end{aligned}$$

Exercise

Q2

Evaluate the line integral

$$\int_C -y dx + z dy + 2x dz,$$

where C is the helix $\mathbf{r}(t) = (\cos t, \sin t, t)$ for $0 \leq t \leq 2\pi$.

1.3 Work Done

Exercise

Q3

Find the work done by the force field

$$\mathbf{F} = x\mathbf{i} + y\mathbf{j} + z\mathbf{k}$$

in moving an object along the curve C parametrized by

$$\mathbf{r}(t) = \cos(\pi t)\mathbf{i} + t^2\mathbf{j} + \sin(\pi t)\mathbf{k}$$

for $0 \leq t \leq 1$.

1.4 Conservative Vector Fields

Important properties:

Let $\mathbf{F}$ be a continuous vector field in a connected, open set G in $\mathbb{R}^n$. The following statements are equivalent:

- (i) $\mathbf{F}$ is independent of path;
- (ii) For any closed curve C in G ,

$$\oint_C \mathbf{F} \cdot d\mathbf{r} = 0,$$

- (iii) $\mathbf{F}$ is conservative.

If $\mathbf{F}$ is conservative, then it is the gradient of a potential function Φ , i.e., $\mathbf{F} = \nabla\Phi$. Then

$$\begin{aligned} \int_C \mathbf{F} \cdot d\mathbf{r} &= \int_C \nabla\Phi \cdot d\mathbf{r} \\ &= \int_a^b \nabla\Phi(\mathbf{r}(t)) \cdot \mathbf{r}'(t) dt \\ &= \Phi(\mathbf{r}(b)) - \Phi(\mathbf{r}(a)) \end{aligned}$$

The proof above is shown in class, the idea is to use Chain rule in a "reverse" way.

Important property:

A vector field $\mathbf{F}$ is conservative if and only if

$$\frac{\partial F_k}{\partial x_j} = \frac{\partial F_j}{\partial x_k}$$

for all j, k . For example, if $\mathbf{F}(x_1, x_2, x_3) = (F_1(x_1, x_2, x_3), F_2(x_1, x_2, x_3), F_3(x_1, x_2, x_3))$, then $\mathbf{F}$ is conservative if and only if

$$\frac{\partial F_1}{\partial x_2} = \frac{\partial F_2}{\partial x_1}, \quad \frac{\partial F_1}{\partial x_3} = \frac{\partial F_3}{\partial x_1}, \quad \frac{\partial F_2}{\partial x_3} = \frac{\partial F_3}{\partial x_2}$$

Exercises

Q4

Find the potential function (if exists) for the vector field

$$\mathbf{F}(x, y, z) = (y^2 + 4zx)\mathbf{i} + y(2x + 2z)\mathbf{j} + (y^2 + 2x^2)\mathbf{k}.$$

