

# MATH 2050C Lecture 10 (Feb 18)

[Problem set 5 posted, due on Feb 26.]

Last week: "Limit Thm" ASSUME  $\lim(x_n), \lim(y_n)$  exist.

$$\begin{cases} \lim(x_n \pm y_n) = \lim(x_n) \pm \lim(y_n) \\ \lim(x_n y_n) = \lim(x_n) \cdot \lim(y_n) \\ \lim\left(\frac{x_n}{y_n}\right) = \frac{\lim(x_n)}{\lim(y_n) \neq 0} \end{cases}$$

If  $x_n \leq y_n \quad \forall n \in \mathbb{N}$   
then  $\lim(x_n) \leq \lim(y_n)$  also OK.  
 $\forall n \geq K$  for some  $K \in \mathbb{N}$

Q: How to prove that  $\lim(x_n)$  exist?


Thm: ("Squeeze / Sandwich Theorem")

Let  $(x_n), (y_n), (z_n)$  be seq. of real numbers s.t.

(1)  $x_n \leq y_n \leq z_n \quad \forall n \in \mathbb{N}$  ( $\forall n \geq K$  for some  $K$ )

(2)  $\lim(x_n) = W = \lim(z_n)$

THEN,  $\lim(y_n) = W$ .



Remark: We do NOT need to assume  $\lim(y_n)$  exists, it follows from the theorem.

E.g.)  $\lim\left(\frac{\sin n}{n}\right) = 0$  because  $-\frac{1}{n} \leq \frac{\sin n}{n} \leq \frac{1}{n}$

Proof: Let  $\epsilon > 0$  be fixed but arbitrary.

$\lim(x_n) = W \Rightarrow \exists K_1 \in \mathbb{N}$  s.t.  $|x_n - W| < \epsilon \quad \forall n \geq K_1$  (\*)

$\lim(z_n) = W \Rightarrow \exists K_2 \in \mathbb{N}$  s.t.  $|z_n - W| < \epsilon \quad \forall n \geq K_2$  (\*\*)

Choose  $K := \max \{k_1, k_2\}$ , then  $\forall n \geq K$

$$-\varepsilon < \overset{(*)}{x_n} - \underset{(1)}{w} \overset{(1)}{\leq} \underset{(1)}{y_n} - \underset{(1)}{w} \overset{(1)}{\leq} \underset{(**)}{z_n} - \underset{(**)}{w} < \varepsilon$$

i.e.  $|y_n - w| < \varepsilon$

---

### Thm: ("Ratio Test")

Let  $(x_n)$  be a seq. st.

(1)  $x_n > 0 \quad \forall n \in \mathbb{N}$

(2)  $\lim \left( \frac{x_{n+1}}{x_n} \right) = L < 1$   
*crucial!*

THEN,  $\lim(x_n) = 0$ .

Motivation  
Geometric seq.  
 $(ar^n) \rightarrow 0$   
provided  $|r| < 1$

Ex: Prove this!

E.g.) Consider  $(x_n) = \left( \frac{n}{2^n} \right)$ , then

$$\left( \frac{x_{n+1}}{x_n} \right) = \left( \frac{\frac{n+1}{2^{n+1}}}{\frac{n}{2^n}} \right) = \left( \frac{n+1}{n} \cdot \frac{1}{2} \right) \rightarrow \frac{1}{2} < 1$$

By Ratio Test.  $\lim \left( \frac{n}{2^n} \right) = 0$ .

Proof: Idea: Compare  $(x_n)$  with a geometric seq.  $(b^n)$ , where  $0 < b < 1$  and apply Squeeze Thm!

Since  $L < 1$ ,  $\exists r \in \mathbb{R}$  st.  $L < r < 1$ .

Take  $\varepsilon = r - L > 0$ , by (2),  $\exists K \in \mathbb{N}$  st.

$$\left| \frac{x_{n+1}}{x_n} - L \right| < \varepsilon = r - L \quad \forall n \geq K$$

$$\Rightarrow \overset{(4)}{0} < \frac{x_{n+1}}{x_n} < L + (r - L) = r < 1$$

$\frac{x_{n+1}}{x_n} \approx L < 1$   
 $x_{n+1} \approx L x_n$   
 $x_{n+2} \approx L^2 x_n$   
 $x_{n+k} \approx L^k x_n$

Thus,  $x_{n+1} < r x_n \quad \forall n \geq k$ .

$$\text{i.e. } 0 < x_n < r x_{n-1} < r^2 x_{n-2} < \dots < r^{n-k} x_k$$

Note:  $\lim_{n \rightarrow \infty} (r^{n-k} x_k) = 0$  since  $r < 1$ .  
k fixed

By Sandwich Thm,  $\lim (x_n) = 0$

Remark: Ratio Test fails if  $L = 1$ .

Consider the seq.  $(x_n) = (n)$ , which is divergent

$$\text{But } \left( \frac{x_{n+1}}{x_n} \right) = \left( \frac{n+1}{n} \right) \rightarrow 1 = 1$$

Ex: Construct an example that  $\left( \frac{x_{n+1}}{x_n} \right) \rightarrow 1$  from below.