

MATH 2050C Lecture 4 (Jan 21)

[Problem set 2 will be posted today, due next Friday.]

Goal: \mathbb{R} is a complete ordered field.

Completeness Property: Every $\emptyset \neq S \subseteq \mathbb{R}$ which is bounded above must have a supremum in \mathbb{R} . [Note: \mathbb{Q} fails this!]

Last time, we proved:

Prop: $u = \sup S$ iff

① $s \leq u, \forall s \in S$ (i.e. u is an upper bd.)

② $\forall \varepsilon > 0, \exists s' \in S$ s.t. $u - \varepsilon < s'$ (i.e. u is the smallest upper bd.)

Similarly, for infimum, we have:

Prop: $u = \inf S$ iff

① $s \geq u, \forall s \in S$ (i.e. u is a lower bd.)

② $\forall \varepsilon > 0, \exists s' \in S$ s.t. $u + \varepsilon > s'$ (i.e. u is the greatest lower bd.)

Q: What about the existence of infimum?

A: It follows from the completeness property.

Prop: Every $\emptyset \neq S \subseteq \mathbb{R}$ that is bounded below must have an infimum in \mathbb{R} .

Proof: Given $\emptyset \neq S \subseteq \mathbb{R}$, consider the subset:

$$\emptyset \neq \bar{S} := \{-s \mid s \in S\} \in \mathbb{R}$$

Claim: \bar{S} is bdd above.

Pf of Claim:


Since S is bdd below, i.e.

\exists some lower bd. u of S

$$\Leftrightarrow u \leq s \quad \forall s \in S$$

$$\Rightarrow -u \geq -s \quad \forall s \in S$$

$\Rightarrow -u$ is an upper bd for \bar{S} i.e. \bar{S} is bdd above.



By Completeness Property, $\sup \bar{S}$ exists in \mathbb{R} .

Claim: $\inf S$ exists. $\inf S = -\sup \bar{S}$.

Pf of Claim:

Check: $-\sup \bar{S}$ is a lower bd for S

(Ex:)

This is the same by reversing the arguments of the claim above.

Check: $-\sup \bar{S}$ is the greatest lower bd. for S

Let $\varepsilon > 0$ be fixed but arbitrary.

(Want to show: $\exists s' \in S$ s.t. $-\sup \bar{S} + \varepsilon > s'$) — (*)

By ② of supremum for \bar{S} ,

$$\sup \bar{S} - \varepsilon < \bar{s}' \quad \text{for some } \bar{s}' \in \bar{S}$$

By defⁿ, we write $\bar{s}' = -s'$ for some $s' \in S$

So, $\sup \bar{S} - \varepsilon < -s' \Rightarrow -\sup \bar{S} + \varepsilon > s'$ for some $s' \in S$
which is (*)

Archimedean Property: \mathbb{N} is NOT bdd above.

Pf: Suppose NOT, i.e. \mathbb{N} is bdd above.

By Completeness Property, $\sup \mathbb{N} =: u \in \mathbb{R}$ exists.

So, $u - 1 < n'$ for some $n' \in \mathbb{N}$.

$$\Rightarrow u < n' + 1 \in \mathbb{N}$$


$\Rightarrow u$ is NOT an upper bd for \mathbb{N} \leftarrow **Contradiction!**

Corollaries:

(i) $\inf \left\{ \frac{1}{n} : n \in \mathbb{N} \right\} = 0$

(ii) $\forall \varepsilon > 0, \exists n \in \mathbb{N}$ s.t. $0 < \frac{1}{n} < \varepsilon$

(iii) $\forall \gamma > 0, \exists!$ $n \in \mathbb{N}$ s.t. $n - 1 < \gamma < n$
 \uparrow
unique



Ex: Prove these!