

1. Recall the definition for the notion of *orthogonality* from the handout *Inner product, norm, and orthogonality*:

Let  $\mathbf{u}, \mathbf{v} \in \mathbb{R}^n$ .

We say  $\mathbf{u}$  is orthogonal to  $\mathbf{v}$ , and write  $\mathbf{u} \perp \mathbf{v}$ , if and only if  $\langle \mathbf{u}, \mathbf{v} \rangle = 0$ .

Also recall these basic properties of orthogonality:

(a) Suppose  $\mathbf{u}, \mathbf{v} \in \mathbb{R}^n$ .

Then  $\mathbf{u} \perp \mathbf{v}$  if and only if  $\mathbf{v} \perp \mathbf{u}$ .

(b) Suppose  $\mathbf{u} \in \mathbb{R}^n$ .

Then  $\mathbf{u} \perp \mathbf{u}$  if and only if  $\mathbf{u} = \mathbf{0}_n$ .

(c) Suppose  $\mathbf{u} \in \mathbb{R}^n$ .

Then  $(\mathbf{u} \perp \mathbf{v} \text{ for any } \mathbf{v} \in \mathbb{R}^n)$  if and only if  $\mathbf{u} = \mathbf{0}_n$ .

(d) Suppose  $\mathbf{u}, \mathbf{v} \in \mathbb{R}^n$ .

Then  $\|\mathbf{u} + \mathbf{v}\|^2 = \|\mathbf{u}\|^2 + \|\mathbf{v}\|^2$  if and only if  $\mathbf{u} \perp \mathbf{v}$ .

## 2. Theorem (A).

Let  $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_k$  be non-zero vectors in  $\mathbb{R}^n$ .

Suppose  $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_k$  are pairwise orthogonal (in the sense that  $\mathbf{u}_i \perp \mathbf{u}_j$  whenever  $i \neq j$ .)

Then the statements below hold:

(a)  $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_k$  are linearly independent.

(b) For any  $\mathbf{v} \in \mathbb{R}^n$ , if  $\mathbf{v}$  is a linear combination of  $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_k$  then

$$\mathbf{v} = \frac{\langle \mathbf{v}, \mathbf{u}_1 \rangle}{\|\mathbf{u}_1\|^2} \mathbf{u}_1 + \frac{\langle \mathbf{v}, \mathbf{u}_2 \rangle}{\|\mathbf{u}_2\|^2} \mathbf{u}_2 + \dots + \frac{\langle \mathbf{v}, \mathbf{u}_k \rangle}{\|\mathbf{u}_k\|^2} \mathbf{u}_k.$$

### 3. Proof of Theorem (A).

Let  $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_k$  be non-zero vectors in  $\mathbb{R}^n$ . Suppose  $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_k$  are pairwise orthogonal.

(a) Pick any  $\alpha_1, \alpha_2, \dots, \alpha_k \in \mathbb{R}$ . Suppose  $\alpha_1 \mathbf{u}_1 + \alpha_2 \mathbf{u}_2 + \dots + \alpha_k \mathbf{u}_k = \mathbf{0}$ .

For each  $j = 1, 2, \dots, k$ , we have

$$\begin{aligned}\alpha_j \|\mathbf{u}_j\|^2 &= \alpha_1 \langle \mathbf{u}_1, \mathbf{u}_j \rangle + \alpha_2 \langle \mathbf{u}_2, \mathbf{u}_j \rangle + \dots + \alpha_k \langle \mathbf{u}_k, \mathbf{u}_j \rangle \\ &= \langle \alpha_1 \mathbf{u}_1 + \alpha_2 \mathbf{u}_2 + \dots + \alpha_k \mathbf{u}_k, \mathbf{u}_j \rangle \\ &= \langle \mathbf{0}, \mathbf{u}_j \rangle = 0\end{aligned}$$

Since  $\mathbf{u}_j$  is not the zero vector,  $\|\mathbf{u}_j\| \neq 0$ . Then  $\alpha_j = 0$ .

It follows that  $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_k$  are linearly independent.

(b) Exercise. (Imitate what has been done above.)

#### 4. Definition. (Orthonormal set and orthonormal basis.)

Let  $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_k \in \mathbb{R}^n$ .

(a) We say that  $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_k$  constitute an orthonormal set in  $\mathbb{R}^n$  if and only if  $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_k$  are pairwise orthogonal and  $\|\mathbf{u}_j\| = 1$  for each  $j = 1, 2, \dots, k$ .

(b) Suppose  $V$  is a subspace of  $\mathbb{R}^n$ .

Then we say that  $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_k$  constitute an orthonormal basis for  $V$  if and only if  $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_k$  constitute a basis for  $V$  and constitute an orthonormal set.

#### Remark.

When  $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_k$  constitute an orthonormal set in  $\mathbb{R}^n$ , they constitute an orthonormal basis for  $\text{Span}(\{\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_k\})$ .

## 5. Theorem (B).

Let  $W$  be a subspace of  $\mathbb{R}^n$ .

Suppose  $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_k$  constitute an orthonormal basis for  $W$ .

Suppose  $\mathbf{s}, \mathbf{t} \in W$ .

Define  $\beta_j = \langle \mathbf{s}, \mathbf{u}_j \rangle$ ,  $\gamma_j = \langle \mathbf{t}, \mathbf{u}_j \rangle$  for each  $j = 1, 2, \dots, k$ .

Then the statements below hold:

(a)  $\mathbf{s} = \beta_1 \mathbf{u}_1 + \beta_2 \mathbf{u}_2 + \dots + \beta_k \mathbf{u}_k$ .

(b)  $\|\mathbf{s}\|^2 = \beta_1^2 + \beta_2^2 + \dots + \beta_k^2$ .

(c)  $\langle \mathbf{s}, \mathbf{t} \rangle = \beta_1 \gamma_1 + \beta_2 \gamma_2 + \dots + \beta_k \gamma_k$ .

## 6. Proof of Theorem (B).

Let  $W$  be a subspace of  $\mathbb{R}^n$ .

Suppose  $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_k$  constitute an orthonormal basis for  $W$ .

Suppose  $\mathbf{s}, \mathbf{t} \in W$ .

Define  $\beta_j = \langle \mathbf{s}, \mathbf{u}_j \rangle$ ,  $\gamma_j = \langle \mathbf{t}, \mathbf{u}_j \rangle$  for each  $j = 1, 2, \dots, k$ .

(a) Since  $\mathbf{s} \in W$  and  $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_k$  constitute a basis for  $W$ ,  $\mathbf{s}$  is a linear combination of  $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_k$ .

Then, by Theorem (A),

$$\mathbf{s} = \beta_1 \mathbf{u}_1 + \beta_2 \mathbf{u}_2 + \dots + \beta_k \mathbf{u}_k.$$

(b) We have

$$\begin{aligned} \|\mathbf{s}\|^2 &= \langle \mathbf{s}, \mathbf{s} \rangle \\ &= \langle \mathbf{s}, \beta_1 \mathbf{u}_1 + \beta_2 \mathbf{u}_2 + \dots + \beta_k \mathbf{u}_k \rangle \\ &= \beta_1 \langle \mathbf{s}, \mathbf{u}_1 \rangle + \beta_2 \langle \mathbf{s}, \mathbf{u}_2 \rangle + \dots + \beta_k \langle \mathbf{s}, \mathbf{u}_k \rangle \\ &= \beta_1^2 + \beta_2^2 + \dots + \beta_k^2. \end{aligned}$$

(c) Exercise.

## 7. Theorem (C).

Let  $W$  be a subspace of  $\mathbb{R}^n$ .

Suppose  $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_k$  constitute an orthonormal basis for  $W$ .

Suppose  $\mathbf{z} \in \mathbb{R}^n$ . Define  $\alpha_1 = \langle \mathbf{z}, \mathbf{u}_1 \rangle$ ,  $\alpha_2 = \langle \mathbf{z}, \mathbf{u}_2 \rangle$ , ...,  $\alpha_k = \langle \mathbf{z}, \mathbf{u}_k \rangle$ .

Define  $\mathbf{v} \in W$  by  $\mathbf{v} = \alpha_1 \mathbf{u}_1 + \alpha_2 \mathbf{u}_2 + \dots + \alpha_k \mathbf{u}_k$ . Define  $\mathbf{y} \in \mathbb{R}^n$  by  $\mathbf{y} = \mathbf{z} - \mathbf{v}$ .

Then the statements below hold:

(a) i.  $\mathbf{z} = \mathbf{v} + \mathbf{y}$ .

ii.  $\mathbf{y} \perp \mathbf{s}$  for any  $\mathbf{s} \in W$ . (In particular,  $\mathbf{y} \perp \mathbf{v}$ .)

(b) Suppose  $\mathbf{s} \in W$ .

Then  $\|\mathbf{z} - \mathbf{s}\| \geq \|\mathbf{z} - \mathbf{v}\|$ . Equality holds if and only if  $\mathbf{s} = \mathbf{v}$ .

(c) The inequality  $\|\mathbf{z}\|^2 \geq \alpha_1^2 + \alpha_2^2 + \dots + \alpha_k^2$  holds.


Moreover, the statements below are logically equivalent:

i.  $\mathbf{z} \in W$ .


ii.  $\mathbf{z} = \alpha_1 \mathbf{u}_1 + \alpha_2 \mathbf{u}_2 + \dots + \alpha_k \mathbf{u}_k$ .

iii.  $\|\mathbf{z}\|^2 = \alpha_1^2 + \alpha_2^2 + \dots + \alpha_k^2$ .

iv. For any  $\mathbf{x} \in \mathbb{R}^n$ ,  $\langle \mathbf{z}, \mathbf{x} \rangle = \alpha_1 \langle \mathbf{u}_1, \mathbf{x} \rangle + \alpha_2 \langle \mathbf{u}_2, \mathbf{x} \rangle + \dots + \alpha_k \langle \mathbf{u}_k, \mathbf{x} \rangle$ .







## 8. Illustrations of the construction described in Theorem (C).

(a) Let  $\mathbf{u}_1 = \begin{bmatrix} \sqrt{3}/2 \\ 1/2 \end{bmatrix}$ , and  $W = \text{Span}(\{\mathbf{u}_1\})$

Note that  $\|\mathbf{u}_1\| = 1$ .

Then  $\mathbf{u}_1$  constitute an orthonormal basis for  $W$ .

• Suppose  $\mathbf{z} = \begin{bmatrix} z_1 \\ z_2 \end{bmatrix}$ .

Define  $\alpha_1 = \langle \mathbf{z}, \mathbf{u}_1 \rangle$ .

Define  $\mathbf{v} = \alpha_1 \mathbf{u}_1$ .

$$\text{Then } \mathbf{v} = \left(\frac{\sqrt{3}}{2}z_1 + \frac{1}{2}z_2\right) \begin{bmatrix} \sqrt{3}/2 \\ 1/2 \end{bmatrix} = \begin{bmatrix} 3z_1/4 + \sqrt{3}z_2/4 \\ \sqrt{3}z_1/4 + z_2/4 \end{bmatrix} = \begin{bmatrix} 3/4 & \sqrt{3}/4 \\ \sqrt{3}/4 & 1/4 \end{bmatrix} \mathbf{z}.$$

Define  $\mathbf{y} = \mathbf{z} - \mathbf{v}$ .

$$\text{Then } \mathbf{y} = \begin{bmatrix} z_1/4 - \sqrt{3}z_2/4 \\ -\sqrt{3}z_1/4 \end{bmatrix} + 3z_2/4 = \begin{bmatrix} 1/4 & -\sqrt{3}/4 \\ -\sqrt{3}/4 & 3/4 \end{bmatrix} \mathbf{z}.$$

$\mathbf{z}$  is ‘decomposed’ into the sum of  $\mathbf{v}, \mathbf{y}$  which form a pair of vectors orthogonal to each other, and in which the vector  $\mathbf{y}$  is orthogonal to every vector in  $W$ .

(b) Let  $\mathbf{u}_1 = \mathbf{e}_1^{(3)}$ ,  $\mathbf{u}_2 = \mathbf{e}_2^{(3)}$ , and  $W = \text{Span}(\{\mathbf{u}_1, \mathbf{u}_2\})$ .

Note that  $\|\mathbf{u}_1\| = \|\mathbf{u}_2\| = 1$  and  $\langle \mathbf{u}_1, \mathbf{u}_2 \rangle = 0$ .

Then  $\mathbf{u}_1, \mathbf{u}_2$  constitute an orthonormal basis for  $W$ .

• Suppose  $\mathbf{z} = \begin{bmatrix} z_1 \\ z_2 \\ z_3 \end{bmatrix}$ .

Define  $\alpha_1 = \langle \mathbf{z}, \mathbf{u}_1 \rangle$ ,  $\alpha_2 = \langle \mathbf{z}, \mathbf{u}_2 \rangle$ .

Define  $\mathbf{v} = \alpha_1 \mathbf{u}_1 + \alpha_2 \mathbf{u}_2$ .

$$\text{Then } \mathbf{v} = z_1 \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} + z_2 \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} = \begin{bmatrix} z_1 \\ z_2 \\ 0 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{bmatrix} \mathbf{z}.$$

Define  $\mathbf{y} = \mathbf{z} - \mathbf{v}$ .

$$\text{Then } \mathbf{y} = \begin{bmatrix} 0 \\ 0 \\ z_3 \end{bmatrix} = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix} \mathbf{z}.$$

$\mathbf{z}$  is 'decomposed' into the sum of  $\mathbf{v}, \mathbf{y}$  which form a pair of vectors orthogonal to each other, and in which the vector  $\mathbf{y}$  is orthogonal to every vector in  $W$ .

(c) Let  $\mathbf{u}_1 = \begin{bmatrix} 1/3 \\ 2/3 \\ 2/3 \end{bmatrix}$ ,  $\mathbf{u}_2 = \begin{bmatrix} -2/3 \\ -1/3 \\ 2/3 \end{bmatrix}$ , and  $W = \text{Span}(\{\mathbf{u}_1, \mathbf{u}_2\})$ .

Note that  $\|\mathbf{u}_1\| = \|\mathbf{u}_2\| = 1$  and  $\langle \mathbf{u}_1, \mathbf{u}_2 \rangle = 0$ .

Then  $\mathbf{u}_1, \mathbf{u}_2$  constitute an orthonormal basis for  $W$ .

• Suppose  $\mathbf{z} = \begin{bmatrix} z_1 \\ z_2 \\ z_3 \end{bmatrix}$ .

Define  $\alpha_1 = \langle \mathbf{z}, \mathbf{u}_1 \rangle$ ,  $\alpha_2 = \langle \mathbf{z}, \mathbf{u}_2 \rangle$ .

Define  $\mathbf{v} = \alpha_1 \mathbf{u}_1 + \alpha_2 \mathbf{u}_2$ .

$$\text{Then } \mathbf{v} = \left(\frac{z_1}{3} + \frac{2z_2}{3} + \frac{2z_3}{3}\right) \begin{bmatrix} 1/3 \\ 2/3 \\ 2/3 \end{bmatrix} + \left(-\frac{2z_1}{3} - \frac{z_2}{3} + \frac{2z_3}{3}\right) \begin{bmatrix} -2/3 \\ -1/3 \\ 2/3 \end{bmatrix} = \cdots = \begin{bmatrix} 5/9 & 4/9 & -2/9 \\ 4/9 & 5/9 & 2/9 \\ -2/9 & 2/9 & 8/9 \end{bmatrix} \mathbf{z}.$$

Define  $\mathbf{y} = \mathbf{z} - \mathbf{v}$ .

$$\text{Then } \mathbf{y} = \cdots = \begin{bmatrix} 4/9 & -4/9 & 2/9 \\ -4/9 & 4/9 & -2/9 \\ 2/9 & -2/9 & 1/9 \end{bmatrix} \mathbf{z}.$$

$\mathbf{z}$  is ‘decomposed’ into the sum of  $\mathbf{v}, \mathbf{y}$  which form a pair of vectors orthogonal to each other, and in which the vector  $\mathbf{y}$  is orthogonal to every vector in  $W$ .

(d) Let  $\mathbf{u}_1 = \begin{bmatrix} 1/2 \\ 1/2 \\ 1/2 \\ 1/2 \end{bmatrix}$ ,  $\mathbf{u}_2 = \begin{bmatrix} -1/2 \\ -1/2 \\ 1/2 \\ 1/2 \end{bmatrix}$ , and  $W = \text{Span}(\{\mathbf{u}_1, \mathbf{u}_2\})$ .

Note that  $\|\mathbf{u}_1\| = \|\mathbf{u}_2\| = 1$  and  $\langle \mathbf{u}_1, \mathbf{u}_2 \rangle = 0$ .

Then  $\mathbf{u}_1, \mathbf{u}_2$  constitute an orthonormal basis for  $W$ .

• Suppose  $\mathbf{z} = \begin{bmatrix} z_1 \\ z_2 \\ z_3 \\ z_4 \end{bmatrix}$ . Define  $\alpha_1 = \langle \mathbf{z}, \mathbf{u}_1 \rangle$ ,  $\alpha_2 = \langle \mathbf{z}, \mathbf{u}_2 \rangle$ . Define  $\mathbf{v} = \alpha_1 \mathbf{u}_1 + \alpha_2 \mathbf{u}_2$ .

$$\text{Then } \mathbf{v} = \left(\frac{z_1}{2} + \frac{z_2}{2} + \frac{z_3}{2} + \frac{z_4}{2}\right) \begin{bmatrix} 1/2 \\ 1/2 \\ 1/2 \\ 1/2 \end{bmatrix} + \left(-\frac{z_1}{2} - \frac{z_2}{2} + \frac{z_3}{2} + \frac{z_4}{2}\right) \begin{bmatrix} -1/2 \\ -1/2 \\ 1/2 \\ 1/2 \end{bmatrix} = \dots = \begin{bmatrix} 1/2 & 1/2 & 0 & 0 \\ 1/2 & 1/2 & 0 & 0 \\ 0 & 0 & 1/2 & 1/2 \\ 0 & 0 & 1/2 & 1/2 \end{bmatrix} \mathbf{z}.$$

$$\text{Define } \mathbf{y} = \mathbf{z} - \mathbf{v}. \text{ Then } \mathbf{y} = \dots = \begin{bmatrix} 1/2 & -1/2 & 0 & 0 \\ -1/2 & 1/2 & 0 & 0 \\ 0 & 0 & 1/2 & -1/2 \\ 0 & 0 & -1/2 & 1/2 \end{bmatrix} \mathbf{z}.$$

$\mathbf{z}$  is ‘decomposed’ into the sum of  $\mathbf{v}, \mathbf{y}$  which form a pair of vectors orthogonal to each other, and in which the vector  $\mathbf{y}$  is orthogonal to every vector in  $W$ .

(e) Let  $\mathbf{u}_1 = \begin{bmatrix} 1/3 \\ 1/3 \\ 0 \\ 2/3 \end{bmatrix}$ ,  $\mathbf{u}_2 = \begin{bmatrix} 2/3 \\ -1/3 \\ 2/3 \\ 0 \end{bmatrix}$ ,  $\mathbf{u}_3 = \begin{bmatrix} 0 \\ -2/3 \\ 1/3 \\ 2/3 \end{bmatrix}$ , and  $W = \text{Span}(\{\mathbf{u}_1, \mathbf{u}_2, \mathbf{u}_3\})$ .

Note that  $\|\mathbf{u}_1\| = \|\mathbf{u}_2\| = \|\mathbf{u}_3\| = 1$  and  $\langle \mathbf{u}_1, \mathbf{u}_2 \rangle = \langle \mathbf{u}_1, \mathbf{u}_3 \rangle = \langle \mathbf{u}_2, \mathbf{u}_3 \rangle = 0$ .

Then  $\mathbf{u}_1, \mathbf{u}_2, \mathbf{u}_3$  constitute an orthonormal basis for  $W$ .

• Suppose  $\mathbf{z} = \begin{bmatrix} z_1 \\ z_2 \\ z_3 \\ z_4 \end{bmatrix}$ . Define  $\alpha_1 = \langle \mathbf{z}, \mathbf{u}_1 \rangle$ ,  $\alpha_2 = \langle \mathbf{z}, \mathbf{u}_2 \rangle$ ,  $\alpha_3 = \langle \mathbf{z}, \mathbf{u}_3 \rangle$ . Define  $\mathbf{v} = \alpha_1 \mathbf{u}_1 + \alpha_2 \mathbf{u}_2 + \alpha_3 \mathbf{u}_3$ .

Then

$$\mathbf{v} = \left(\frac{z_1}{3} + \frac{2z_2}{3} + \frac{2z_4}{3}\right) \begin{bmatrix} 1/3 \\ 2/3 \\ 0 \\ 2/3 \end{bmatrix} + \left(\frac{2z_1}{3} - \frac{z_2}{3} + \frac{2z_3}{3}\right) \begin{bmatrix} 2/3 \\ -1/3 \\ 2/3 \\ 0 \end{bmatrix} + \left(-\frac{2z_2}{3} - \frac{z_3}{3} + \frac{2z_4}{3}\right) \begin{bmatrix} 0 \\ -2/3 \\ -1/3 \\ 2/3 \end{bmatrix} = \begin{bmatrix} 5/9 & 0 & 4/9 & 2/9 \\ 0 & 1 & 0 & 0 \\ 4/9 & 0 & 5/9 & -2/9 \\ 2/9 & 0 & -2/9 & 8/9 \end{bmatrix} \mathbf{z}.$$

Define  $\mathbf{y} = \mathbf{z} - \mathbf{v}$ . Then  $\mathbf{y} = \dots = \begin{bmatrix} 4/9 & 0 & -4/9 & -2/9 \\ 0 & 0 & 0 & 0 \\ -4/9 & 0 & 4/9 & 2/9 \\ -2/9 & 0 & 2/9 & 1/9 \end{bmatrix} \mathbf{z}$ .

$\mathbf{z}$  is ‘decomposed’ into the sum of  $\mathbf{v}, \mathbf{y}$  which form a pair of vectors orthogonal to each other, and in which the vector  $\mathbf{y}$  is orthogonal to every vector in  $W$ .

## 9. Proof of Theorem (C).

Let  $W$  be a subspace of  $\mathbb{R}^n$ .

Suppose  $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_k$  constitute an orthonormal basis for  $W$ .

Suppose  $\mathbf{z} \in \mathbb{R}^n$ .

Define  $\alpha_1 = \langle \mathbf{z}, \mathbf{u}_1 \rangle$ ,  $\alpha_2 = \langle \mathbf{z}, \mathbf{u}_2 \rangle$ , ...,  $\alpha_k = \langle \mathbf{z}, \mathbf{u}_k \rangle$ .

Define  $\mathbf{v} = \alpha_1 \mathbf{u}_1 + \alpha_2 \mathbf{u}_2 + \dots + \alpha_k \mathbf{u}_k$ , and  $\mathbf{y} = \mathbf{z} - \mathbf{v}$ .

(a) i. By definition,  $\mathbf{z} = \mathbf{v} + \mathbf{y}$ .

ii. Pick any  $\mathbf{s} \in W$ . Define  $\beta_1 = \langle \mathbf{s}, \mathbf{u}_1 \rangle$ ,  $\beta_2 = \langle \mathbf{s}, \mathbf{u}_2 \rangle$ , ...,  $\beta_k = \langle \mathbf{s}, \mathbf{u}_k \rangle$ .

Then  $\mathbf{s} = \beta_1 \mathbf{u}_1 + \beta_2 \mathbf{u}_2 + \dots + \beta_k \mathbf{u}_k$ .

Note that  $\langle \mathbf{v}, \mathbf{s} \rangle = \alpha_1 \beta_1 + \alpha_2 \beta_2 + \dots + \alpha_k \beta_k$ .

Also note that

$$\begin{aligned} \langle \mathbf{z}, \mathbf{s} \rangle &= \langle \mathbf{z}, \beta_1 \mathbf{u}_1 + \beta_2 \mathbf{u}_2 + \dots + \beta_k \mathbf{u}_k \rangle \\ &= \beta_1 \langle \mathbf{z}, \mathbf{u}_1 \rangle + \beta_2 \langle \mathbf{z}, \mathbf{u}_2 \rangle + \dots + \beta_k \langle \mathbf{z}, \mathbf{u}_k \rangle \\ &= \alpha_1 \beta_1 + \alpha_2 \beta_2 + \dots + \alpha_k \beta_k. \end{aligned}$$

Then  $\langle \mathbf{y}, \mathbf{s} \rangle = \langle \mathbf{z} - \mathbf{v}, \mathbf{s} \rangle = \langle \mathbf{z}, \mathbf{s} \rangle - \langle \mathbf{v}, \mathbf{s} \rangle = 0$ .

Therefore  $\mathbf{y} \perp \mathbf{s}$ .

(b) Suppose  $\mathbf{s} \in W$ .

Note that  $\mathbf{v} \in W$ . Then  $\mathbf{v} - \mathbf{s} \in W$ . Then  $\mathbf{z} - \mathbf{v} \perp \mathbf{v} - \mathbf{s} \in W$ .

• We have  $\|\mathbf{z} - \mathbf{s}\|^2 = \|(\mathbf{z} - \mathbf{v}) + (\mathbf{v} - \mathbf{s})\|^2 = \|\mathbf{z} - \mathbf{v}\|^2 + \|\mathbf{v} - \mathbf{s}\|^2$ .

Since  $\|\mathbf{v} - \mathbf{s}\|^2 \geq 0$ , we have  $\|\mathbf{z} - \mathbf{s}\|^2 \geq \|\mathbf{z} - \mathbf{v}\|^2$ .

Then  $\|\mathbf{z} - \mathbf{s}\| \geq \|\mathbf{z} - \mathbf{v}\|$ .

• Suppose  $\mathbf{s} = \mathbf{v}$ . Then  $\|\mathbf{z} - \mathbf{s}\| = \|\mathbf{z} - \mathbf{v}\|$ .

• Suppose  $\|\mathbf{z} - \mathbf{s}\| = \|\mathbf{z} - \mathbf{v}\|$ . Then  $\|\mathbf{v} - \mathbf{s}\|^2 = 0$ .

Therefore  $\mathbf{s} - \mathbf{v} = \mathbf{0}$ . Hence  $\mathbf{s} = \mathbf{v}$ .

(c) Exercise. (Apply the definition of  $\mathbf{v}$  and  $\mathbf{y}$ .

The inequality concerned is simply ' $\|\mathbf{z}\| \geq \|\mathbf{v}\|$ ' in disguise.

Equality holds if and only if  $\mathbf{y} = \mathbf{0}$ .)



10. Recall the definition for the notion of *orthogonal complement of a subspace of  $\mathbb{R}^n$*  from the handout *Orthogonal complement*.

*Suppose  $W$  is a subspace of  $\mathbb{R}^n$ .*

*The perp of  $W$ , which as a set is given by  $W^\perp = \{\mathbf{x} \in \mathbb{R}^n : \mathbf{x} \perp \mathbf{u} \text{ for any } \mathbf{u} \in W\}$ , is called the orthogonal complement of  $W$  in  $\mathbb{R}^n$ .*

Also recall the result  $(\star)$  from the same handout:

*Suppose  $W$  is a subspace of  $\mathbb{R}^n$ . Then for any  $\mathbf{z} \in \mathbb{R}^n$ , there exist some unique  $\mathbf{s} \in W$ ,  $\mathbf{t} \in W^\perp$  such that  $\mathbf{z} = \mathbf{s} + \mathbf{t}$ .*

With the help of the result  $(\star)$ , we can enrich the content of part (a) in Theorem (C) by appending a ‘uniqueness part’.

## 11. Theorem (D).

Let  $W$  be a subspace of  $\mathbb{R}^n$ .

Suppose  $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_k$  constitute an orthonormal basis for  $W$ .

Suppose  $\mathbf{z} \in \mathbb{R}^n$ .

Define  $\alpha_1 = \langle \mathbf{z}, \mathbf{u}_1 \rangle$ ,  $\alpha_2 = \langle \mathbf{z}, \mathbf{u}_2 \rangle$ , ...,  $\alpha_k = \langle \mathbf{z}, \mathbf{u}_k \rangle$ .

Define  $\mathbf{v} \in W$  by  $\mathbf{v} = \alpha_1 \mathbf{u}_1 + \alpha_2 \mathbf{u}_2 + \dots + \alpha_k \mathbf{u}_k$ .

Define  $\mathbf{y} \in \mathbb{R}^n$  by  $\mathbf{y} = \mathbf{z} - \mathbf{v}$ .

Then the statements below hold:

(a) i.  $\mathbf{z} = \mathbf{v} + \mathbf{y}$ .

ii.  $\mathbf{y} \perp \mathbf{s}$  for any  $\mathbf{s} \in W$ . (In particular,  $\mathbf{y} \perp \mathbf{v}$ .)

(b) Suppose  $\mathbf{v}', \mathbf{y}' \in \mathbb{R}^n$ .

Suppose  $\mathbf{v}' \in W$ ,  $\mathbf{z} = \mathbf{v}' + \mathbf{y}'$ , and  $\mathbf{y}' \perp \mathbf{s}$  for any  $\mathbf{s} \in W$ . Then  $\mathbf{v}' = \mathbf{v}$  and  $\mathbf{y}' = \mathbf{y}$ .

## Remarks.

- In plain words, statement (b) is saying that  $\mathbf{z}$  is decomposed in a unique way as a sum of two vectors, one in  $W$  and the other in  $W'$ . The two vectors are  $\mathbf{v}$  and  $\mathbf{y}$  respectively.

The vector  $\mathbf{v}$  is determined independent of the choice of orthonormal bases for  $W$ :

Suppose that  $\mathbf{u}'_1, \mathbf{u}'_2, \dots, \mathbf{u}'_k$  also constitute an orthonormal basis for  $W$ , and  $\alpha'_1 = \langle \mathbf{z}, \mathbf{u}'_1 \rangle, \alpha'_2 = \langle \mathbf{z}, \mathbf{u}'_2 \rangle, \dots, \alpha'_k = \langle \mathbf{z}, \mathbf{u}'_k \rangle$ .

Further suppose that  $\mathbf{v}' = \alpha'_1 \mathbf{u}'_1 + \alpha'_2 \mathbf{u}'_2 + \dots + \alpha'_k \mathbf{u}'_k$  and  $\mathbf{y}' = \mathbf{z} - \mathbf{v}'$ .

Then it happens that  $\mathbf{v}' = \mathbf{v}$  and  $\mathbf{y}' = \mathbf{y}$ .

- *Terminology.*

This uniqueness makes sense of naming the vectors  $\mathbf{v}, \mathbf{y}$  with reference to  $\mathbf{z}$  and  $W$ .

The vector  $\mathbf{v}$  is called the orthogonal projection of the vector  $\mathbf{z}$  onto  $W$ . It is denoted by  $\text{pr}_W(\mathbf{z})$ .

The vector  $\mathbf{y}$  is called the orthogonal complement of  $\mathbf{z}$  with respect to  $W$ .

The other parts of Theorem (C) can be re-stated in terms of orthogonal projections.

## 12. Theorem (E).

Let  $W$  be a subspace of  $\mathbb{R}^n$ , and  $\mathbf{z} \in \mathbb{R}^n$ .

(a) Suppose  $\mathbf{s} \in W$ .

Then  $\|\mathbf{z} - \mathbf{s}\| \geq \|\mathbf{z} - \mathbf{pr}_W(\mathbf{z})\|$ .

Equality holds if and only if  $\mathbf{s} = \mathbf{pr}_W(\mathbf{z})$ .

(b) The inequality  $\|\mathbf{z}\| \geq \|\mathbf{pr}_W(\mathbf{z})\|$  holds.

Equality holds if and only if  $\mathbf{z} \in W$ .

### Remarks.

- Statement (a) says that amongst all vectors in  $W$ , it is  $\mathbf{pr}_W(\mathbf{z})$  whose distance with  $\mathbf{z}$  is the smallest.

In plain words,  $\mathbf{pr}_W(\mathbf{z})$  is the ‘closest (or best) approximation’ to  $\mathbf{z}$  amongst all vectors in  $W$ .

This result is the corner stone of the ‘least square method’ for finding approximations.

- Statement (b) says that the ‘length’ of the vector  $\mathbf{v}$  is no less than that of its projection onto  $W$ , which is  $\mathbf{pr}_W(\mathbf{z})$ .

This inequality is known as Bessel’s Inequality.

### 13. Theorem (F).

Let  $W$  be a subspace of  $\mathbb{R}^n$ .

Suppose  $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_k$  constitute an orthonormal basis for  $W$ .

Define the  $(n \times k)$ -matrix  $U$  by  $U = \left[ \mathbf{u}_1 \mid \mathbf{u}_2 \mid \dots \mid \mathbf{u}_k \right]$ .

Then the statements below hold:

- (a) For any  $\mathbf{z} \in \mathbb{R}^n$ ,  $\text{pr}_W(\mathbf{z}) = UU^t\mathbf{z}$ .                      (c)  $\mathcal{C}(UU^t) = W$ .  
(b)  $UU^t$  is symmetric and idempotent.                      (d)  $\mathcal{N}(UU^t) = W^\perp$ .

#### Remarks.

- When  $\mathbf{s}_1, \mathbf{s}_2, \dots, \mathbf{s}_k$  constitute an orthonormal basis for  $W$  and  $S = \left[ \mathbf{s}_1 \mid \mathbf{s}_2 \mid \dots \mid \mathbf{s}_k \right]$ , we have  $\text{pr}_W(\mathbf{z}) = SS^t\mathbf{z}$  for any  $\mathbf{z} \in \mathbb{R}^n$ . It follows that  $UU^t = SS^t$ .

This  $(n \times n)$ -square matrix is independent of the choice of orthonormal bases for  $W$ .

- *Terminology.*

This uniqueness makes sense of naming the matrix  $UU^t$  with reference to  $W$ .

The matrix  $UU^t$  is called the projection matrix from  $\mathbb{R}^n$  onto  $W$ . Multiplication by this matrix from the left to a vector in  $\mathbb{R}^n$  results in the orthogonal projection of that vector onto  $W$ .

## 14. Proof of Theorem (F).

Let  $W$  be a subspace of  $\mathbb{R}^n$ .

Suppose  $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_k$  constitute an orthonormal basis for  $W$ .

Define the  $(n \times k)$ -matrix  $U$  by  $U = \left[ \mathbf{u}_1 \mid \mathbf{u}_2 \mid \dots \mid \mathbf{u}_k \right]$ .

(a) Pick any  $\mathbf{z} \in \mathbb{R}^n$ . We have

$$\begin{aligned} UU^t \mathbf{z} &= U \begin{bmatrix} \mathbf{u}_1^t \\ \mathbf{u}_2^t \\ \vdots \\ \mathbf{u}_k^t \end{bmatrix} \mathbf{z} = U \begin{bmatrix} \mathbf{u}_1^t \mathbf{z} \\ \mathbf{u}_2^t \mathbf{z} \\ \vdots \\ \mathbf{u}_k^t \mathbf{z} \end{bmatrix} = \left[ \mathbf{u}_1 \mid \mathbf{u}_2 \mid \dots \mid \mathbf{u}_k \right] \begin{bmatrix} \langle \mathbf{z}, \mathbf{u}_1 \rangle \\ \langle \mathbf{z}, \mathbf{u}_2 \rangle \\ \vdots \\ \langle \mathbf{z}, \mathbf{u}_k \rangle \end{bmatrix} \\ &= \langle \mathbf{z}, \mathbf{u}_1 \rangle \mathbf{u}_1 + \langle \mathbf{z}, \mathbf{u}_2 \rangle \mathbf{u}_2 + \dots + \langle \mathbf{z}, \mathbf{u}_k \rangle \mathbf{u}_k = \text{pr}_W(\mathbf{z}) \end{aligned}$$

(b) We have  $(UU^t)^t = (U^t)^t U^t = UU^t$ . Then  $UU^t$  is symmetric.

We have  $(UU^t)^2 = (UU^t)(UU^t) = U(U^t U)U^t = U I_k U^t = UU^t$ . Then  $UU^t$  is idempotent.

(c) We verify that  $W = \mathcal{C}(UU^t)$ :

- [We verify that for any  $\mathbf{x} \in \mathbb{R}^n$ , if  $\mathbf{x} \in W$  then  $\mathbf{x} \in \mathcal{C}(UU^t)$ .]

Pick any  $\mathbf{x} \in \mathbb{R}^n$ . Suppose  $\mathbf{x} \in W$ .

Since  $\mathbf{x} \in W$ , We have  $\mathbf{x} = \text{pr}_W(\mathbf{x})$ .

By the result in part (a), we have  $\text{pr}_W(\mathbf{x}) = UU^t\mathbf{x}$ .

Then  $\mathbf{x} = UU^t\mathbf{x}$ . Therefore, by definition,  $\mathbf{x} \in \mathcal{C}(UU^t)$ .

- [We verify that for any  $\mathbf{x} \in \mathbb{R}^n$ , if  $\mathbf{x} \in \mathcal{C}(UU^t)$  then  $\mathbf{x} \in W$ .]

Pick any  $\mathbf{x} \in \mathbb{R}^n$ . Suppose  $\mathbf{x} \in \mathcal{C}(UU^t)$ .

Then there exists some  $\mathbf{s} \in \mathbb{R}^k$  such that  $\mathbf{x} = UU^t\mathbf{s}$ .

Define  $\mathbf{p} \in \mathbb{R}^k$  by  $\mathbf{p} = U^t\mathbf{s}$ .

Then  $\mathbf{x} = U\mathbf{p}$ .

Therefore, by definition,  $\mathbf{x} \in \mathcal{C}(U)$ .

By definition,  $W = \text{Span}(\{\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_k\}) = \mathcal{C}(U)$ . Hence  $\mathbf{x} \in W$ .

(d) We have verified that  $\mathcal{C}(UU^t) = W$ .

By part (b),  $UU^t$  is symmetric.

Then  $\mathcal{N}((UU^t)) = \mathcal{N}((UU^t)^t) = (\mathcal{C}(UU^t))^\perp = W^\perp$ .

## 15. Illustrations of the content of Theorem (F).

(a) Let  $\mathbf{u}_1 = \begin{bmatrix} \sqrt{3}/2 \\ 1/2 \end{bmatrix}$ , and  $W = \text{Span}(\{\mathbf{u}_1\})$

$\mathbf{u}_1$  constitute an orthonormal basis for  $W$ .

Define  $U = \mathbf{u}_1$ .

$$\text{We have } UU^t = \begin{bmatrix} 3/4 & \sqrt{3}/4 \\ \sqrt{3}/4 & 1/4 \end{bmatrix}.$$

$UU^t$  is the projection matrix from  $\mathbb{R}^2$  onto  $W$ : for any  $\mathbf{z} \in \mathbb{R}^2$ ,  $\text{pr}_W(\mathbf{z}) = UU^t\mathbf{z}$ .

(b) Let  $\mathbf{u}_1 = \mathbf{e}_1^{(3)}$ ,  $\mathbf{u}_2 = \mathbf{e}_2^{(3)}$ , and  $W = \text{Span}(\{\mathbf{u}_1, \mathbf{u}_2\})$ .

$\mathbf{u}_1, \mathbf{u}_2$  constitute an orthonormal basis for  $W$ .

Define  $U = \begin{bmatrix} \mathbf{u}_1 & | & \mathbf{u}_2 \end{bmatrix}$ .

$$\text{We have } UU^t = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{bmatrix}.$$

$UU^t$  is the projection matrix from  $\mathbb{R}^3$  onto  $W$ : for any  $\mathbf{z} \in \mathbb{R}^3$ ,  $\text{pr}_W(\mathbf{z}) = UU^t\mathbf{z}$ .



(c) Let  $\mathbf{u}_1 = \begin{bmatrix} 1/3 \\ 2/3 \\ 2/3 \end{bmatrix}$ ,  $\mathbf{u}_2 = \begin{bmatrix} -2/3 \\ -1/3 \\ 2/3 \end{bmatrix}$ , and  $W = \text{Span}(\{\mathbf{u}_1, \mathbf{u}_2\})$ .

$\mathbf{u}_1, \mathbf{u}_2$  constitute an orthonormal basis for  $W$ .

Define  $U = \left[ \mathbf{u}_1 \mid \mathbf{u}_2 \right]$ .

We have  $UU^t = \begin{bmatrix} 5/9 & 4/9 & -2/9 \\ 4/9 & 5/9 & 2/9 \\ -2/9 & 2/9 & 8/9 \end{bmatrix}$ .

$UU^t$  is the projection matrix from  $\mathbb{R}^3$  onto  $W$ : for any  $\mathbf{z} \in \mathbb{R}^3$ ,  $\text{pr}_W(\mathbf{z}) = UU^t\mathbf{z}$ .

(d) Let  $\mathbf{u}_1 = \begin{bmatrix} 1/2 \\ 1/2 \\ 1/2 \\ 1/2 \end{bmatrix}$ ,  $\mathbf{u}_2 = \begin{bmatrix} -1/2 \\ -1/2 \\ 1/2 \\ 1/2 \end{bmatrix}$ , and  $W = \text{Span}(\{\mathbf{u}_1, \mathbf{u}_2\})$ .

$\mathbf{u}_1, \mathbf{u}_2$  constitute an orthonormal basis for  $W$ .

Define  $U = \left[ \mathbf{u}_1 \mid \mathbf{u}_2 \right]$ .

We have  $UU^t = \begin{bmatrix} 1/2 & 1/2 & 0 & 0 \\ 1/2 & 1/2 & 0 & 0 \\ 0 & 0 & 1/2 & 1/2 \\ 0 & 0 & 1/2 & 1/2 \end{bmatrix}$ .

$UU^t$  is the projection matrix from  $\mathbb{R}^4$  onto  $W$ : for any  $\mathbf{z} \in \mathbb{R}^4$ ,  $\text{pr}_W(\mathbf{z}) = UU^t\mathbf{z}$ .

(e) Let  $\mathbf{u}_1 = \begin{bmatrix} 1/3 \\ 1/3 \\ 0 \\ 2/3 \end{bmatrix}$ ,  $\mathbf{u}_2 = \begin{bmatrix} 2/3 \\ -1/3 \\ 2/3 \\ 0 \end{bmatrix}$ ,  $\mathbf{u}_3 = \begin{bmatrix} 0 \\ -2/3 \\ 1/3 \\ 2/3 \end{bmatrix}$ , and  $W = \text{Span}(\{\mathbf{u}_1, \mathbf{u}_2, \mathbf{u}_3\})$ .

$\mathbf{u}_1, \mathbf{u}_2, \mathbf{u}_3$  constitute an orthonormal basis for  $W$ .

Define  $U = \left[ \mathbf{u}_1 \mid \mathbf{u}_2 \mid \mathbf{u}_3 \right]$ .

We have  $UU^t = \begin{bmatrix} 5/9 & 0 & 4/9 & 2/9 \\ 0 & 1 & 0 & 0 \\ 4/9 & 0 & 5/9 & -2/9 \\ 2/9 & 0 & -2/9 & 8/9 \end{bmatrix}$ .

$UU^t$  is the projection matrix from  $\mathbb{R}^4$  onto  $W$ : for any  $\mathbf{z} \in \mathbb{R}^4$ ,  $\text{pr}_W(\mathbf{z}) = UU^t\mathbf{z}$ .