

1. (a) Suppose ζ is a complex number. Verify that $|\zeta|^2 = \zeta\bar{\zeta}$.
 (b) Let z, w be complex numbers. Suppose $w \neq 0$. Verify that $\frac{z}{w} = \frac{z\bar{w}}{|w|^2}$.
Remark. This looks easy. And it is indeed easy. But it suggests division involving complex numbers can be done in a much more comfortable way. (How?)
2. Let z, w be complex numbers.
 - (a) Verify that $|zw| = |z||w|$.
 - (b) Suppose $z \neq 0$ and $w \neq 0$, and θ, φ are respective arguments of z, w . Verify that $zw = |zw|(\cos(\theta + \varphi) + i \sin(\theta + \varphi))$.

3. Let $\omega = \frac{\sqrt{3} + i}{2}$.
 - (a) Write down the respective values of $\omega^2, \omega^3, \omega^{11}, \omega^{12}$.
 - (b) Hence, or otherwise, find the value of $\sum_{k=0}^{2230} \omega^{k+1}$.

4. Let a, b, c be real numbers. Suppose $a^2 + b^2 + c^2 = 1$ and $c \neq 1$. Define $z = \frac{a + bi}{1 - c}$.
 - (a) Express $|z|^2$ in terms of c alone.
 - (b) Express each of a, b, c in terms of z, \bar{z} alone.
5. (a) Prove the statement below, known as the **Parallelogram Identity**:
 - Suppose z, w are complex numbers. Then $|z + w|^2 + |z - w|^2 = 2|z|^2 + 2|w|^2$.
 (b) Prove the statements below:
 - i. Suppose r, s, t are complex numbers. Then $|2r - s - t|^2 + |2s - t - r|^2 + |2t - r - s|^2 = 3(|s - t|^2 + |t - r|^2 + |r - s|^2)$.
 - ii. Let ζ, α, β be complex numbers. Suppose $\zeta^2 = \alpha^2 + \beta^2$. Then $|\zeta + \alpha| + |\zeta - \alpha| = |\zeta + \beta| + |\zeta - \beta|$.

6. Let $z \in \mathbb{C}$. Suppose $4|z + 1| = |z + 16|$. Verify that z lies on the circle with centre at 0 and with radius 4 (on the Argand plane).

7. \diamond Write $\mu = 1 + i, \nu = 3 - i$.
 - (a) Let $\eta \in \mathbb{C}$. Prove that the point η lies on the ('infinite') straight line joining μ, ν iff $\operatorname{Re}(\eta) + \operatorname{Im}(\eta) - 2 = 0$.
 - (b) Consider the curve C on the Argand plane defined by the equation $|z - \mu| = |z - \nu|$.
 - i. Prove that the curve C is also described by the equation $\operatorname{Re}(z) - \operatorname{Im}(z) - 2 = 0$.
 - ii. Hence, or otherwise, prove that the curve C is the perpendicular bisector for the line segment joining μ, ν (on the Argand plane).

Remark. Given the same straight line, there are various ways to write down an equation (with unknown z in the complex numbers) whose solution set is the straight line concerned on the Argand plane:

- $a\operatorname{Re}(z) + b\operatorname{Im}(z) + c = 0$, in which a, b, c are given real number with a, b being not both zero.
- $\alpha\bar{z} + \bar{\alpha}z + d = 0$, in which α is a given non-zero complex number and d is a given real number.
- $|z - \beta| = |z - \gamma|$, in which β, γ are given distinct complex numbers.

8. *There is no need to give any justifications for your answers in this question.*

Find all the solutions of the system of equations $\begin{cases} |z - 2 - 2i| = 2 \\ |z - 4 + 2i| = |z - 2i| \end{cases}$ with unknown z in \mathbb{C} .

Remark. What are the curves described by the respective equations in the Argand plane?

9. *There is no need to give any justifications for your answers in this question.*

Consider the system of equations $(S_{\alpha, r}) : \begin{cases} |z - 2i| = 2 \\ |z - 4 - 4i| = |z| \\ |z - \alpha| = r \end{cases}$ with unknown z in \mathbb{C} . Here α is some complex number and r is a non-negative real number.

Suppose that $(S_{\alpha, r})$ has two distinct solutions.

- (a) Write down all solutions of $(S_{\alpha, r})$.
- (b) What is the smallest possible value of r ?
- (c) What is the value of α if $|\operatorname{Re}(\alpha)| = |\operatorname{Im}(\alpha)|$?

Remark. What are the curves described by the respective equations in the Argand plane?

10. Let ω be a complex number. Suppose $|\omega| = 1$ and $\text{Im}(\omega) \geq 0$. Further suppose $\omega^2 + \frac{5}{\omega} - 2$ is purely imaginary. Find the value of ω .

Remark. Use the polar form of ω .

11. Let α, β, γ be real numbers. Suppose $\alpha + \beta + \gamma = 2\pi$.

Define λ, μ, ν by $\lambda = \cos(\alpha) + i \sin(\alpha)$, $\mu = \cos(\beta) + i \sin(\beta)$, $\nu = \cos(\gamma) + i \sin(\gamma)$ respectively.

(a) Find the value of $\lambda\mu\nu$.

(b) Prove that $\cos(\alpha) = \frac{1}{2}(\lambda + \frac{1}{\lambda})$ and $\cos(2\alpha) = \frac{1}{2}(\lambda^2 + \frac{1}{\lambda^2})$.

(c) \diamond Hence, or otherwise, prove that $\cos(2\alpha) + \cos(2\beta) + \cos(2\gamma) = 4 \cos(\alpha) \cos(\beta) \cos(\gamma) - 1$.

12. (a) \diamond Prove the statements below:

i. Suppose z is a complex number. Then $|\text{Re}(z)| \leq |z|$. Moreover, equality holds iff z is real.

ii. Suppose z be a complex number. Then $|\text{Im}(z)| \leq |z|$. Moreover, equality holds iff z is purely imaginary.

(b) \diamond Prove the statement below:

• Suppose u, v are complex numbers. Then $|\text{Re}(u\bar{v})| \leq |u||v|$. Moreover, equality holds iff there exist some real numbers p, q such that $pu + qv = 0$ and p, q are not both zero.

(c) \clubsuit Prove the statements below (known collectively as the **Triangle Inequality for complex numbers**):

i. Suppose z, w are complex numbers. Then $|z + w| \leq |z| + |w|$. Moreover, equality holds iff there exist some non-negative real numbers s, t such that $sz = tw$ and s, t are not both zero.

ii. Suppose u, v are complex numbers. Then $||u| - |v|| \leq |u - v|$. Moreover, equality holds iff there exist some non-negative real numbers h, k such that $hu = kv$ and h, k are not both zero.

13. We introduce the definitions below:

- Let $z \in \mathbb{C}$. The number z is said to be a **Gaussian integer** if both of $\text{Re}(z)$, $\text{Im}(z)$ are integers.
- The set of all Gaussian integers is denoted by \mathbb{G} .

(a) Prove the statements below:

i. Suppose $s \in \mathbb{Z}$. Then $s \in \mathbb{G}$.

ii. Let s is a Gaussian integer. Suppose $s \neq 0$. Then $|s| \geq 1$.

iii. Suppose $s, t \in \mathbb{G}$. Then $\bar{s} \in \mathbb{G}$, $s + t \in \mathbb{G}$ and $st \in \mathbb{G}$.

iv. Suppose $s, t \in \mathbb{G}$. Further suppose $|st| = 1$. Then $|s| = |t| = 1$.

(b) We introduce the definition below:

- Let $u, v \in \mathbb{G}$. The number u is said to be **\mathbb{G} -divisible** by v if there exists some $s \in \mathbb{G}$ such that $u = sv$.

Prove the statements below:

i. Suppose $u \in \mathbb{G}$. Then u is \mathbb{G} -divisible by u .

ii. \diamond Let $u, v \in \mathbb{G}$. Suppose (u is \mathbb{G} -divisible by v and v is \mathbb{G} -divisible by u). Then $|u| = |v|$.

iii. Let $u, v, w \in \mathbb{G}$. Suppose (u is \mathbb{G} -divisible by v and v is \mathbb{G} -divisible by w). Then u is \mathbb{G} -divisible by w .

iv. Let $u, v, t \in \mathbb{G}$. Suppose (u is \mathbb{G} -divisible by t and v is \mathbb{G} -divisible by t). Then $u + v$ is \mathbb{G} -divisible by t .

v. Let $u, v, t \in \mathbb{G}$. Suppose (u is \mathbb{G} -divisible by t or v is \mathbb{G} -divisible by t). Then uv is \mathbb{G} -divisible by t .

vi. Let $u, v \in \mathbb{G}$. Suppose $u \neq 0$ and u is \mathbb{G} -divisible by v . Then $|v| \leq |u|$.

(c) \diamond We introduce the definition below:

- Let $z \in \mathbb{C}$. The number z is said to be a **Gaussian rational** if there exist some Gaussian integer u, v such that $v \neq 0$ and $u = vz$.

Prove the statements below:

i. Suppose z is a rational number. Then z is a Gaussian rational.

ii. Suppose z is a Gaussian integer. Then z is a Gaussian rational.

iii. Suppose $z \in \mathbb{C}$. Then z is a Gaussian rational iff (there exist some $s, t \in \mathbb{Q}$ such that $z = s + ti$.)

iv. Suppose z, w are Gaussian rationals. Then $z + w$ is a Gaussian rational.

v. Suppose z, w are Gaussian rationals. Then $z - w$ is a Gaussian rational.

vi. Suppose z, w are Gaussian rationals. Then zw is a Gaussian rational.

vii. Suppose z, w are Gaussian rationals. Further suppose $w \neq 0$. Then $\frac{z}{w}$ is a Gaussian rational.

Remark. Preview on your algebra course. The set \mathbb{G} is a subset of \mathbb{C} and contains \mathbb{Z} as a subset. In your algebra course, the set \mathbb{G} is likely denoted by $\mathbb{Z}[i]$. The set \mathbb{G} , together with addition and multiplication for complex numbers, constitutes an *integral domain*; further together with the modulus for complex numbers, it constitutes an *Euclidean domain*. It behaves in many ways similar to integers, and similar to polynomials with real coefficients. Many results on the notion of ‘*divisibility*’ for integers and polynomials with real coefficients in school maths have their analogues for Gaussian integers. The set of all Gaussian rationals is a subset of \mathbb{C} and contains each of \mathbb{Q} and \mathbb{G} as a subset. In your algebra course, it is likely denoted by $\mathbb{Q}[i]$ in some contexts, and by $\mathbb{Q}(i)$ in some other contexts. This set, together with addition and multiplication for complex numbers, constitutes a *field*.