Assignment 10

Coverage: 16.4 and 16.5(part) in Text.

Exercises: 16.4 no 7, 11, 14, 23, 26, 28, 35, 37, 39. 16.5 no. 4, 8, 10, 13, 33. Hand in 16.4 no 14, 28, 35; 16.5 no 8, 33 November 23.

Supplementary Problems

1. Let D be the parallelogram formed by the lines x + y = 1, x + y = 3, y = 2x - 3, y = 2x + 2. Evaluate the line integral

$$\oint_C dx + 3xy \, dy$$

where C is the boundary of D oriented in anticlockwise direction. Suggestion: Try Green's theorem and then apply change of variables formula.

2. Let $F = M\mathbf{i} + N\mathbf{j}$ be a smooth vector field which is defined in \mathbb{R}^2 except at the origin. Suppose that it satisfies the component test $M_y = N_x$. Show that for any simple closed curve γ enclosing the origin and oriented in positive direction, one has

$$\oint_{\gamma} M dx + N dy = \varepsilon \int_{0}^{2\pi} \left[-M(\varepsilon \cos \theta, \varepsilon \sin \theta) \sin \theta + N(\varepsilon \cos \theta, \varepsilon \sin \theta) \cos \theta \right] d\theta ,$$

for all sufficiently small ε . What happens when γ does not enclose the origin?

3. Let

$$\mathbf{H} = \frac{-y}{x^2 + y^2}\mathbf{i} + \frac{x}{x^2 + y^2}\mathbf{j},$$

which is defined in the plane except at the origin.

- (a) Explain why **H** is conservative in the half plane $\{(x, y) : x > 0\}$.
- (b) Find a potential function for **H** in this half plane.