

Recall

Reflexive spaces

Let X be a Banach space and $Q: X \rightarrow X^{**}$ be the *canonical map (natural embedding)*, i.e.,

$$(Qx)(x^*) := x^*(x) \text{ or symmetrically, } \langle x^*, Qx \rangle := \langle x, x^* \rangle.$$

If $QX = X^{**}$, then X is called *reflexive*.


Let M be a closed subspace of X .

- X reflexive $\iff X^*$ reflexive.
- X reflexive $\iff M$ & X/M reflexive. The \Leftarrow direction is called *Three space property* and the proof relies on the isometric isomorphism $\tilde{r}: X^*/M^\perp \rightarrow M^*$ where \tilde{r} is the split of the restriction map $r: X^* \rightarrow M^*$ along the natural projection $\pi: X^* \rightarrow X^*/M^\perp$.

If X is a **separable** Banach space, then:

- (Helley's selection) bounded sequence in X^* has w^* -convergent subsequence.
- (In X^* , a sequence is w^* -convergent \implies norm convergent.) $\iff \dim X < \infty$.
- X is reflexive \implies bounded sequence in X has weakly convergent subsequence.

Suppose M is a nonzero proper closed subspace of a Banach space X . In the same notation of Lecture Notes, we may have the following diagram



where the blue arrows denote the isometries by [LN, Prop. 4.12, Prop. 5.1, & Lem. 5.8]. We denote the dashed lines to show the relationship between a Banach space and its dual. Note that $M^* = X^*/M^\perp$ by \tilde{r} and $M^\perp = (X/M)^*$ by π^* .

$C[0, 1]$ is not reflexive

Example 1. $C[0, 1]$ is not reflexive.

In the following, we consider the spaces to be Banach spaces. We prove [Example 1](#) by the necessary conditions or properties of reflexive spaces, i.e., by checking that $C[0, 1]$ does not have some property that belongs to a reflexive space. Note that $C[0, 1]$ is separable.

Proof by closed subspaces of a reflexive space are reflexive. It suffices to construct an embedding $T: c_0 \rightarrow C[0, 1]$. For $n \in \mathbb{N}$, let $d_n = \frac{1}{n} - \frac{1}{n+1}$ and define a ‘triangle’ shaped function

$$f_n(t) = \begin{cases} \frac{4[t-(1/(n+1)+d_n/4)]}{d_n} & , t \in [(1/(n+1) + d_n/4), (1/(n+1) + d_n/2)) \\ -\frac{4[t-(1/(n+1)+d_n/2)]}{d_n} + 1 & , t \in [(1/(n+1) + d_n/2), (1/(n+1) + 3d_n/4)] \\ 0 & \text{otherwise.} \end{cases}$$

Then $\text{supp}(f_n) \subset (1/(n+1), 1/n)$ and $\|f_n\|_\infty = 1$. For $x = (x_n)_{n=1}^\infty \in c_0$, define Tx by

$$Tx(t) = \sum_{n=1}^{\infty} x_n f_n(t) \quad \text{for } t \in [0, 1].$$

Since the supports of $f_n(t)$ are disjoint, for every $t \in [0, 1]$, $|Tx(t)| \leq \|x\|_\infty$. Since for each x_n there exists t_n such that $Tx(t_n) = x_n$, we have $\|x\|_\infty = \|Tx\|_\infty$. The injection and linearity is easily checked.

Hence T embeds c_0 into $C[0, 1]$. Tc_0 is a closed subspace by the completeness of c_0 and not reflexive since c_0 is not reflexive, thus $C[0, 1]$ is not reflexive. \square

Proof by the dual of a reflexive separable space is separable. Recall from Tutorial 3 that $(C[0, 1])^* = BV_0^+[0, 1]$. We will show that $BV_0^+[0, 1]$ is not separable. For any $x \in (0, 1)$, define

$$f_x(t) = \begin{cases} 0 & , t \in [0, x) \\ 1 & , t \in [x, 1]. \end{cases}$$

Then $f_x \in BV_0^+[0, 1]$ and for any $x \neq y$, $V(f_x - f_y) = 2$. However, the cardinality of $\{f_x : x \in (0, 1)\} \subset BV_0^+[0, 1]$ is uncountable. Hence $(C[0, 1])^* = BV_0^+[0, 1]$ is not separable. \square

Proof by the weakly sequentially compactness of closed unit ball in a reflexive separable space.

Consider the sequence of functions $f_n(x) = x^n \in C[0, 1]$. Then $\|f_n\|_\infty = 1$ and every subsequence of f_n will converge pointwisely to $f = \begin{cases} 1 & x = 1 \\ 0 & x \neq 1 \end{cases} \notin C[0, 1]$.

For every $x \in [0, 1]$, it follows (from Homework 3) that the evaluation functional $\delta_x(f) = f(x)$ for $f \in C[0, 1]$ is bounded, thus $\delta_x \in (C[0, 1])^*$. Suppose otherwise that $C[0, 1]$ is reflexive. Then by [LN, Coro. 6.12] there exists a subsequence of f_n weakly convergent in $C[0, 1]$, and hence pointwisely convergent in $C[0, 1]$, which is a contradiction. \square