

Math 2010 E Advanced Calculus I

Differential Calculus of Functions of Several variables.

$f(x, y)$, $f(x, y, z)$, $\underbrace{f(x_1, \dots, x_n)}_{\text{mainly}}$ $n=2, 3$ (most of the times)

Vectors in \mathbb{R}^n = $\vec{x} = (x_1, \dots, x_n) \in \mathbb{R}^n$

$\mathbb{R}^n = \mathbb{R} \times \dots \times \mathbb{R}$ (n copies of \mathbb{R})

$= \{(x_1, \dots, x_n) : x_i \in \mathbb{R} \text{ for } i=1, \dots, n\}$



Remarks : (i) In Textbook, "bold face" \mathbf{x} is used to denote a vector.

(ii) In high school, \vec{AB} is a vector with initial point A & terminal point B.

For $\vec{x} = (x_1, \dots, x_n)$ is a vector with initial point $A = O = (0, \dots, 0)$ & terminal point $B = (x_1, \dots, x_n)$.

(iii) \mathbb{R}^n is called Cartesian product of n -copies of \mathbb{R} .

(x_1, \dots, x_n) Cartesian/rectangular coordinates of the point.



Basic Operations of Vectors

Let $\vec{a} = (a_1, \dots, a_n) \in \mathbb{R}^n$

$\vec{b} = (b_1, \dots, b_n) \in \mathbb{R}^n$

$r \in \mathbb{R}$

Equality

$$\vec{a} = \vec{b} \Leftrightarrow a_i = b_i \text{ for } i=1, \dots, n.$$

Addition

$$\vec{a} + \vec{b} = (a_1 + b_1, \dots, a_n + b_n)$$


Scalar Multiplication

$$r\vec{a} = (ra_1, \dots, ra_n)$$

Subtraction

$$\begin{aligned}\vec{a} - \vec{b} &= \vec{a} + (-1)\vec{b} \\ &= (a_1 - b_1, \dots, a_n - b_n)\end{aligned}$$

(Parallelogram law)



(Ex: What is $\vec{a} - \vec{b}$?)

length, Dot (Inner) Product in \mathbb{R}^n

Defn: Let $\vec{a} = (a_1, \dots, a_n) \in \mathbb{R}^n$
 $\vec{b} = (b_1, \dots, b_n) \in \mathbb{R}^n$

Then dot product (or inner product)

$$\vec{a} \cdot \vec{b} = a_1 b_1 + \dots + a_n b_n$$

↑
 (Have to write the dot !)

In particular,

$$\vec{a} \cdot \vec{a} = a_1^2 + \dots + a_n^2 \stackrel{\text{def}}{=} \|\vec{a}\|^2$$

$\|\vec{a}\| = \sqrt{a_1^2 + \dots + a_n^2}$ is called the length/magnitude of the vector \vec{a} .

(In Textbook, it's denoted by $|\vec{a}|$.)

Remark: Another notation for dot (inner) product

$$\langle \vec{a}, \vec{b} \rangle = \vec{a} \cdot \vec{b}$$

Properties of Dot Product

Let $\vec{a}, \vec{b}, \vec{c} \in \mathbb{R}^n$, $r \in \mathbb{R}$. Then

$$(1) \quad (\vec{a} + \vec{b}) \cdot \vec{c} = \vec{a} \cdot \vec{c} + \vec{b} \cdot \vec{c}$$

$$\vec{a} \cdot (\vec{b} + \vec{c}) = \vec{a} \cdot \vec{b} + \vec{a} \cdot \vec{c}$$

$$(2) \quad (r\vec{a}) \cdot \vec{b} = \vec{a} \cdot (r\vec{b}) = r(\vec{a} \cdot \vec{b})$$

$$(3) \quad \vec{a} \cdot \vec{b} = \vec{b} \cdot \vec{a}$$


$$(4) \quad \vec{a} \cdot \vec{a} \geq 0 \text{ and "equality holds" } \Leftrightarrow \vec{a} = \vec{0} = (0, \dots, 0)$$

$$(5) \quad \vec{a} \cdot \vec{a} = \|\vec{a}\|^2$$

$$(6) \quad \vec{a} \cdot \vec{b} = \|\vec{a}\| \|\vec{b}\| \cos \theta$$

where $\theta = \text{angle between } \vec{a} \text{ & } \vec{b}$

Hence, if $\vec{a}, \vec{b} \neq \vec{0}$,



$$\vec{a} \cdot \vec{b} = 0 \Leftrightarrow \cos \theta = 0 \Leftrightarrow \vec{a} \perp \vec{b} \text{ (perpendicular)}$$

Remarks (i) For $n=2, 3$, (5) & (6) are geometric properties of \mathbb{R}^n .


(ii) For $n \geq 4$, $\|\vec{a}\|$ in (5) & (6) is defined before and the θ in (6) is the definition

$$\theta \stackrel{\text{def}}{=} \cos^{-1} \left(\frac{\vec{a} \cdot \vec{b}}{\|\vec{a}\| \|\vec{b}\|} \right) \quad \text{for high dimensional vectors}$$

(iii) If $\|\vec{u}\|=1$, then \vec{u} is called a unit vector.

$$\begin{aligned} \text{eg: } \hat{i} &= (1, 0, 0) \\ \hat{j} &= (0, 1, 0) \\ \hat{k} &= (0, 0, 1) \end{aligned} \quad \left. \begin{array}{l} \\ \\ \end{array} \right\} \text{are unit vectors in } \mathbb{R}^3.$$

Pf of (6) (for $n \leq 3$) (Assume $\theta \leq \frac{\pi}{2}$)



Note $\|\vec{b} - \vec{a}\|^2 = (\vec{b} - \vec{a}) \cdot (\vec{b} - \vec{a})$ (by (5))

$$\begin{aligned}
 &= \vec{b} \cdot (\vec{b} - \vec{a}) - \vec{a} \cdot (\vec{b} - \vec{a}) \\
 &= (\vec{b} \cdot \vec{b} - \vec{b} \cdot \vec{a}) - (\vec{a} \cdot \vec{b} - \vec{a} \cdot \vec{a}) \\
 &= \|\vec{b}\|^2 - 2\vec{a} \cdot \vec{b} + \|\vec{a}\|^2
 \end{aligned} \quad (\text{by (3) \& (5)})$$

From the figure

$$\begin{aligned}
 \|\vec{b} - \vec{a}\|^2 &= (\|\vec{a}\| - \|\vec{b}\| \cos \theta)^2 + (\|\vec{b}\| \sin \theta)^2 \\
 &= \|\vec{a}\|^2 - 2\|\vec{a}\| \|\vec{b}\| \cos \theta + \|\vec{b}\|^2 \cos^2 \theta + \|\vec{b}\|^2 \sin^2 \theta \\
 &= \|\vec{a}\|^2 - 2\|\vec{a}\| \|\vec{b}\| \cos \theta + \|\vec{b}\|^2 \\
 \therefore \|\vec{a}\| \|\vec{b}\| \cos \theta &= \vec{a} \cdot \vec{b} . \quad \times
 \end{aligned}$$


e.g. : Suppose \vec{v} & \vec{w} are vectors of the same length.

Show that $(\vec{v} + \vec{w}) \cdot (\vec{v} - \vec{w}) = 0$.

Soln : $(\vec{v} + \vec{w}) \cdot (\vec{v} - \vec{w}) = \vec{v} \cdot \vec{v} + \vec{w} \cdot \vec{v} - \vec{v} \cdot \vec{w} - \vec{w} \cdot \vec{w}$

$$\begin{aligned}
 &= \|\vec{v}\|^2 - \|\vec{w}\|^2 \\
 &= 0 \quad \text{by assumption that } \vec{v} \text{ & } \vec{w} \text{ have same length}
 \end{aligned} \quad \times$$


Remark : $(\vec{v} + \vec{w}) \perp (\vec{v} - \vec{w})$ provided $\|\vec{v}\| = \|\vec{w}\|$,



Geometric meaning :

Diagonals of a rhombus are perpendicular.

e.g:



AB is diameter, C on the circle

Show that $\angle ACB = 90^\circ = \frac{\pi}{2}$

Solu: $\vec{AC} = \vec{AO} + \vec{OC}$ (check)

$$\vec{BC} = \vec{BO} + \vec{OC}$$

$$AB = \text{diameter} \Rightarrow \vec{BO} = -\vec{AO}$$

$$\begin{aligned}
 \text{Then } \vec{AC} \cdot \vec{BC} &= (\vec{AO} + \vec{OC}) \cdot (\vec{BO} + \vec{OC}) \\
 &= (\vec{AO} + \vec{OC}) \cdot (-\vec{AO} + \vec{OC}) \\
 &= -\|\vec{AO}\|^2 - \vec{OC} \cdot \vec{AO} + \vec{AO} \cdot \vec{OC} + \|\vec{OC}\|^2 \\
 &= -\|\vec{AO}\|^2 + \|\vec{OC}\|^2 \\
 &= 0 \quad \text{since } \|\vec{OC}\| = \|\vec{AO}\| = \text{radius}.
 \end{aligned}$$

$$\therefore \angle ACB = \frac{\pi}{2}$$

※

Two inequalities (Au's Book §1.2)

Cauchy-Schwarz Inequality

Let $\vec{a}, \vec{b} \in \mathbb{R}^n$. Then $|\vec{a} \cdot \vec{b}| \leq \|\vec{a}\| \|\vec{b}\|$

$$\text{i.e. } \left| \sum_{i=1}^n a_i b_i \right| \leq \sqrt{\sum_{i=1}^n a_i^2} \sqrt{\sum_{i=1}^n b_i^2}$$

Equality holds $\Leftrightarrow \vec{a} = r\vec{b}$ or $\vec{b} = r\vec{a}$

$$|\vec{a} \cdot \vec{b}| = \|\vec{a}\| \|\vec{b}\| \text{ for some } r \in \mathbb{R}$$


Pf: Case 1 If $\vec{a} = \vec{0}$ or $\vec{b} = \vec{0}$, then both sides are 0
equality holds $\vec{a} = 0\vec{b}$ or $\vec{b} = 0\vec{a}$.

Case 2 $\vec{a} \neq \vec{0}$ & $\vec{b} \neq \vec{0}$.

$$\text{Let } f(t) = \|\vec{t}\vec{a} - \vec{b}\|^2 \text{ for } t \in \mathbb{R}.$$

$$\begin{aligned} \text{Then } f(t) &= (\vec{t}\vec{a} - \vec{b}) \cdot (\vec{t}\vec{a} - \vec{b}) \\ &= t^2 \|\vec{a}\|^2 - 2\vec{t}\vec{a} \cdot \vec{b} + \|\vec{b}\|^2 \quad (\geq 0) \end{aligned}$$

$$\text{Note } f(t) = \|\vec{t}\vec{a} - \vec{b}\|^2 \geq 0, \forall t \in \mathbb{R}$$



$$\therefore t^2 \|\vec{a}\|^2 - 2\vec{t}\vec{a} \cdot \vec{b} + \|\vec{b}\|^2 \geq 0$$

$$\Rightarrow \Delta \leq 0$$

$$\text{i.e. } (-2\vec{a} \cdot \vec{b})^2 - 4(\|\vec{a}\|^2)(\|\vec{b}\|^2) \leq 0$$

$$\Rightarrow |\vec{a} \cdot \vec{b}| \leq \|\vec{a}\| \|\vec{b}\|$$

Equality holds $\Leftrightarrow \Delta = 0$
 $\Leftrightarrow f(t)$ has a repeated root $r \in \mathbb{R}$
 $\Leftrightarrow \|r\vec{a} - \vec{b}\|^2 = 0$
 $\Leftrightarrow r\vec{a} = \vec{b}$. ~~✓~~