

## Inverse function

Thm 6.1.8 Let •  $I \subseteq \mathbb{R}$  be an interval

- $f: I \rightarrow \mathbb{R}$  be strictly monotone and continuous.
- $J = f(I)$  and  $g: J \rightarrow \mathbb{R}$  be the strictly monotone & continuous function inverse to  $f$ .

If  $f$  is differentiable at  $c \in I$  and  $f'(c) \neq 0$ , then  $g$  is differentiable at  $d = f(c)$  and

$$g'(d) = \frac{1}{f'(c)} = \frac{1}{f'(g(d))}$$

(Note  $f'(c) \neq 0$  doesn't follow from  $f$  being strictly monotone:  
eg.  $f(x) = x^3$  is strictly monotone, but  $f'(0) = 0$ .  
In this case, the inverse  $g(x) = x^{1/3}$  is not differentiable at  $x=0$ .)

Pf: Since  $f$  is differentiable at  $x=c$ , Carathéodory's Thm 6.1.5

$\Rightarrow \exists \varphi: I \rightarrow \mathbb{R}$  with  $\varphi$  continuous at  $c$  such that

$$\begin{cases} f(x) - f(c) = \varphi(x)(x-c), \quad \forall x \in I, \text{ and} \\ \varphi(c) = f'(c) \end{cases}$$

Since  $f'(c) \neq 0$  and  $\varphi$  is continuous at  $c$ ,  $\exists \delta > 0$  such that

$$\varphi(x) \neq 0, \quad \forall x \in (c-\delta, c+\delta) \cap I.$$

$$\text{Let } U = f((c-\delta, c+\delta) \cap I) \subset J$$

Then the inverse function  $g$  satisfies  $f(g(y)) = y, \forall y \in U$ .

$$\text{Hence } y - d = f(g(y)) - f(c) = \varphi(g(y))(g(y) - c)$$

$$= \varphi(g(y))(g(y) - g(d)) \quad \left( \begin{array}{l} d = f(c) \in U \\ \Rightarrow c = g(d) \end{array} \right)$$

Since  $g(y) \in (c-\delta, c+\delta) \cap I, \forall y \in U$ ,

we have  $\varphi(g(y)) \neq 0$ .

$$\text{Hence } g(y) - g(d) = \frac{1}{\varphi(g(y))} (y - d).$$

Since  $g$  is continuous on  $J$  and  $\varphi$  is continuous at  $c = g(d) \neq 0$ ,

$\frac{1}{\varphi} \circ g$  is continuous at  $d$ .

Then by Carathéodory's Thm 6.1.5,  $g$  is differentiable at  $d = f(c)$

$$\text{and } g'(d) = \frac{1}{\varphi(g(d))} = \frac{1}{\varphi(c)} = \frac{1}{f'(c)} \quad \text{. } \times$$

Thm 6.1.9 (Same notations as in Thm 6.1.8)

Let  $f: I \rightarrow \mathbb{R}$  be strict monotone (no need to assume continuity).

If  $f$  is differentiable on  $I$  and  $f'(x) \neq 0, \forall x \in I$ . Then the

inverse function  $g$  is differentiable on  $J = f(I)$  and

$$g' = \frac{1}{f' \circ g}$$

Pf:  $f$  diff. on  $I \Rightarrow f$  is continuous. then apply Thm 6.1.8 to all  $x \in I$ . ~~✗~~

Remark on simplified notations:

Usually, we write  $y=f(x)$  and  $x=g(y)$  for functions inverse to each other. Then the formula in Thm 6.1.9 can be written as

$$g'(y) = \frac{1}{(f \circ g)'(y)} \quad \forall y \in J$$

$$\sim (g \circ f)'(x) = \frac{1}{f'(x)}, \quad \forall x \in I$$

In this notation, one often simply write

$$g'(y) = \frac{1}{f'(x)}$$

without explicitly stated that  $y=f(x)$  &  $x=g(y)$ !

eg 6.1.10

(a)  $f(x) = x^5 + 4x + 3$  gives a strictly increasing (why?) and continuous function on  $\mathbb{R}$  (and  $f(\mathbb{R}) = \mathbb{R}$  why?)

$$f'(x) = 5x^4 + 4 \geq 4 > 0.$$

Therefore, Thm 6.1.8  $\Rightarrow g = f^{-1}$  is differentiable  $\forall y \in \mathbb{R}$ .

And for example, at  $x=1$ ,  $g'(8) = g'(f(1)) = \frac{1}{f'(1)} = \frac{1}{9}$

(b)  $f: [0, \infty) \rightarrow [0, \infty)$  given by  $f(x) = x^n$  where  $n = 2, 4, 6, \dots$

Then  $f$  is strictly increasing continuous on  $[0, \infty)$

Note that  $f([0, \infty)) = [0, \infty)$ . The inverse function  $g$  defines on  $[0, \infty)$  and is strictly increasing and continuous.

Since  $f'(x) = nx^{n-1} > 0$ ,  $\forall x > 0$ , &  $f((0, \infty)) = (0, \infty)$ ,

$g$  is differentiable  $\forall y > 0$  and

$$g'(y) = \frac{1}{f'(g(y))} = \frac{1}{n(g(y))^{n-1}} = \frac{1}{n(y^{\frac{1}{n}})^{n-1}} = \frac{1}{n} y^{\frac{1}{n}-1}$$

(The inverse is denoted by  $g(y) = y^{\frac{1}{n}}$ ,  $\forall y \in [0, \infty)$ .)

Note:  $g$  is not differentiable at  $y=0$  (one side derivative doesn't exist). Omitted! . But the argument is the same as in the next example.)

(c)  $n = 3, 5, 7, \dots$ .  $F(x) = x^n$ ,  $\forall x \in \mathbb{R}$ , is strictly increasing & continuous.

Inverse is  $G(y) = y^{\frac{1}{n}}$ ,  $\forall y \in \mathbb{R}$ .

As in example (b) above,  $G$  is differentiable  $\forall y \neq 0$

and  $G'(y) = \frac{1}{n} y^{\frac{1}{n}-1}$  (check!)

And again,  $G$  is not differentiable at  $y=0$

Pf Suppose that  $G$  is differentiable at  $y=0$ .

Then consider the composite function  $y = F(G(y))$ .

Since  $G(0)=0$  and  $F'(0)=0$  exists.

$$\text{Chain rule implies } 1 = \frac{dy}{dy} = \underbrace{F'(G(0))}_0 \underbrace{G'(0)}_{\text{exists}} = 0$$

which is a contradiction.  $\therefore G'(0)$  doesn't exist ~~✗~~

(d) Recall if  $r = \frac{m}{n} > 0$ ,  $m, n \in \{1, 3, 5, \dots\}$ , then

$$x^r = x^{\frac{m}{n}} \text{ is defined as } (x^{\frac{1}{n}})^m, \quad \forall x \geq 0.$$

Therefore, the function  $R: [0, \infty) \rightarrow [0, \infty)$  defined by

$$R(x) = x^r, \quad \forall x \geq 0$$

is a composite function  $R = f \circ g$  where

$$g(x) = x^{\frac{1}{n}}, \quad x \geq 0 \quad (\text{the inverse discussed in eg(b)})$$

$$\text{and } f(x) = x^m, \quad x \geq 0$$


$$(\text{i.e. } R(x) = x^r = (x^{\frac{1}{n}})^m = f(g(x)), \quad \forall x \in [0, \infty))$$

Then Chain rule  $\Rightarrow \quad \forall x \in [0, \infty)$

$$\begin{aligned} R'(x) &= f'(g(x)) g'(x) = m (x^{\frac{1}{n}})^{m-1} \cdot \frac{1}{n} x^{\frac{1}{n}-1} \\ &= \left(\frac{m}{n}\right) x^{\left(\frac{m}{n}\right)-1} \end{aligned}$$

$\therefore (x^r)' = r x^{r-1}, \quad \forall x \geq 0$ , true for all rational  $r > 0$ .

(e)  $\sin x$  is strictly increasing on  $I = [-\frac{\pi}{2}, \frac{\pi}{2}]$   
and maps  $I$  to  $J = [-1, 1]$ .



$\Rightarrow$  inverse exists, and we denote it by

$$\text{Arcsin} : [-1, 1] \rightarrow [-\frac{\pi}{2}, \frac{\pi}{2}]$$

i.e. If  $x \in [-\frac{\pi}{2}, \frac{\pi}{2}]$  &  $y \in [-1, 1]$ , then

$$y = \sin x \Leftrightarrow x = \text{Arcsin } y.$$

Note that  $D \sin x = \cos x \neq 0$  for  $x \in (-\frac{\pi}{2}, \frac{\pi}{2})$  (no end pts.)

Thm 6.1.8  $\Rightarrow$

$$\begin{aligned} D \text{Arcsin } y &= \frac{1}{D \sin x} = \frac{1}{\cos x} = \frac{1}{\sqrt{1 - \sin^2 x}} \\ &= \frac{1}{\sqrt{1 - y^2}}, \quad \forall y \in (-1, 1) \end{aligned}$$

(Note:  $D \text{Arcsin } y$  does not exist for  $y = \pm 1$ . Check!)