

Properties of Integral

Thm 7.1.5 Suppose $f, g \in R[a, b]$. Then

(a) $kf \in R[a, b]$, $\forall k \in \mathbb{R}$ and

$$\int_a^b kf = k \int_a^b f$$

(b) $f+g \in R[a, b]$ and

$$\int_a^b (f+g) = \int_a^b f + \int_a^b g$$

(c) $f(x) \leq g(x) \quad \forall x \in [a, b] \Rightarrow \int_a^b f \leq \int_a^b g$.

Pf: (a) Ex. (Similar to the proof of (b) & easier)

(b) $f, g \in R[a, b] \Rightarrow$

$\forall \epsilon > 0$, $\exists \delta_1 > 0$ st. $|S(f, \dot{\mathcal{P}}) - \int_a^b f| < \epsilon$, $\forall \dot{\mathcal{P}}$ with $\|\dot{\mathcal{P}}\| < \delta_1$

& $\exists \delta_2 > 0$ st. $|S(g, \dot{\mathcal{P}}) - \int_a^b g| < \epsilon$, $\forall \dot{\mathcal{P}}$ with $\|\dot{\mathcal{P}}\| < \delta_2$.

Also note that for any $\dot{\mathcal{P}} = \{\overline{[x_{i-1}, x_i]}, t_i\}_{i=1}^n$

$$\begin{aligned} S(f+g; \dot{\mathcal{P}}) &= \sum_{i=1}^n (f+g)(t_i) (x_i - x_{i-1}) \\ &= \sum_{i=1}^n f(t_i) (x_i - x_{i-1}) + \sum_{i=1}^n g(t_i) (x_i - x_{i-1}) \\ &= S(f; \dot{\mathcal{P}}) + S(g; \dot{\mathcal{P}}) \end{aligned}$$

Then $\forall \dot{\sigma}$ with $\|\dot{\sigma}\| < \delta = \min\{\delta_1, \delta_2\}$, we have

$$\begin{aligned} & |S(f+g; \dot{\sigma}) - (\int_a^b f + \int_a^b g)| \\ & \leq |S(f; \dot{\sigma}) - \int_a^b f| + |S(g; \dot{\sigma}) - \int_a^b g| \\ & < \varepsilon + \varepsilon = 2\varepsilon \end{aligned}$$

Since $\varepsilon > 0$ is arbitrary, we've proved that

$$f+g \in R[a, b] \text{ and } \int_a^b (f+g) = \int_a^b f + \int_a^b g.$$

(c) As in (b), we conclude, $\forall \varepsilon > 0$, $\exists \delta > 0$ s.t,

for $\dot{\sigma}$ with $\|\dot{\sigma}\| < \delta$,

$$|S(f; \dot{\sigma}) - \int_a^b f| < \varepsilon \text{ and } |S(g; \dot{\sigma}) - \int_a^b g| < \varepsilon$$

$$\Rightarrow \int_a^b f - \varepsilon < S(f; \dot{\sigma}) \quad \& \quad S(g; \dot{\sigma}) < \int_a^b g + \varepsilon.$$

Now $f(x) \leq g(x)$, $\forall x \in [a, b] \Rightarrow$

$$S(f; \dot{\sigma}) = \sum_{i=1}^n f(t_i)(x_i - x_{i-1}) \leq \sum_{i=1}^n g(t_i)(x_i - x_{i-1}) = S(g; \dot{\sigma})$$

$$\therefore \int_a^b f - \varepsilon < S(f; \dot{\sigma}) \leq S(g; \dot{\sigma}) < \int_a^b g + \varepsilon$$

$$\text{or } \int_a^b f < \int_a^b g + 2\varepsilon.$$

Since $\varepsilon > 0$ is arbitrary, $\int_a^b f \leq \int_a^b g$ ~~**~~

Boundedness Theorem

Thm 7.1.6 $f \in \mathcal{R}[a, b] \Rightarrow f$ is bounded on $[a, b]$.

(of course, $f \in \mathcal{R}[a, b] \nLeftrightarrow f$ is bounded on $[a, b]$, see later section)

Pf Let $f \in \mathcal{R}[a, b]$ and $\int_a^b f = L$.

And suppose on the contrary that f is unbounded on $[a, b]$

$f \in \mathcal{R}[a, b]$ with $\int_a^b f = L$ (Take $\varepsilon = 1$ in the def)

$\Rightarrow \exists \delta > 0$ such that

$\forall \tilde{\mathcal{P}}$ with $\|\tilde{\mathcal{P}}\| < \delta$,

$$|\int(f; \tilde{\mathcal{P}}) - L| < 1$$

$$\Rightarrow |\int(f; \tilde{\mathcal{P}})| < |L| + 1 \quad (\times),$$

If $\mathcal{P} = \{[x_{i-1}, x_i]\}_{i=1}^n$ be a partition of $[a, b]$.

Then f unbounded

$\Rightarrow \exists$ a subinterval $[x_{i_0-1}, x_{i_0}]$ s.t.

f is unbounded on $[x_{i_0-1}, x_{i_0}]$.

Therefore, we can find x_{i_0} such that

$$|f(x_{i_0})(x_{i_0} - x_{i_0-1})| > |L| + 1 + \left| \sum_{i \neq i_0} f(x_i)(x_i - x_{i-1}) \right| \quad (\times)_2$$

Then the corresponding tagged partition $\overset{\circ}{\mathcal{P}} = \{[x_{i-1}, x_i]; t_i\}_{i=1}^n$
with tags $\begin{cases} t_i = x_i & \text{for } i \neq i_0 \\ t_{i_0} \end{cases}$

gives Riemann sum

$$S(f; \overset{\circ}{\mathcal{P}}) = f(t_{i_0})(x_{i_0} - x_{i_0-1}) + \sum_{i \neq i_0} f(x_i)(x_i - x_{i-1})$$

$$\Rightarrow f(t_{i_0})(x_{i_0} - x_{i_0-1}) = S(f; \overset{\circ}{\mathcal{P}}) - \sum_{i \neq i_0} f(x_i)(x_i - x_{i-1})$$

$$\Rightarrow |f(t_{i_0})(x_{i_0} - x_{i_0-1})| \leq |S(f; \overset{\circ}{\mathcal{P}})| + \left| \sum_{i \neq i_0} f(x_i)(x_i - x_{i-1}) \right|$$

$$\left(\text{by } (\ast)_1 \right) \leq (L) + 1 + \left| \sum_{i \neq i_0} f(x_i)(x_i - x_{i-1}) \right|$$

which contradicts $(\ast)_2$

$\therefore f$ must be bounded. ~~XX~~

Ex 7.1.7 Thomae's function

$$f(x) = \begin{cases} \frac{1}{n}, & \text{if } x = \frac{m}{n} \in [0, 1] \text{ & } m, n \text{ have no common factors} \\ 0, & \text{if } x = 0 \\ 0, & \text{if } x \text{ is irrational} \text{ & } x \in [0, 1]. \end{cases}$$

$N = 11, 23, \dots$

Then $f \in R[0, 1]$ & $\int_a^b f = 0$

Note: h is discontinuous at every rational number in $[0,1]$
& continuous at every irrational number in $[0,1]$.
(see eg 5.1.6(h) of the Textbook)

Pf: (Similar to eg 7.1.4(d))

$\forall \varepsilon > 0$, the set $E_\varepsilon = \{x \in [0,1] : h(x) \geq \frac{\varepsilon}{2}\}$ is a finite set

(For instance $\frac{\varepsilon}{2} = \frac{1}{5}$, then $E_\varepsilon = \{0, 1, \frac{1}{2}, \frac{1}{3}, \frac{2}{3}, \frac{1}{4}, \frac{3}{4}, \frac{1}{5}, \frac{2}{5}, \frac{3}{5}, \frac{4}{5}\}$)

Let $N_\varepsilon = \#$ of elements in E_ε .

Define $\delta_\varepsilon = \frac{\varepsilon}{4N_\varepsilon} > 0$.

Then $\forall \dot{\mathcal{P}} = \{[x_{i-1}, x_i]; t_i\}_{i=1}^n$ with $\|\dot{\mathcal{P}}\| < \delta_\varepsilon$,

$$S(h; \dot{\mathcal{P}}) = \sum_{\substack{i=1 \\ t_i \notin E_\varepsilon}}^n R(t_i)(x_i - x_{i-1}) + \sum_{\substack{i=1 \\ t_i \in E_\varepsilon}}^n R(t_i)(x_i - x_{i-1})$$

$\leftarrow t_i \in \text{at most } 2 \text{ subintervals} \Rightarrow \text{at most } 2N_\varepsilon \text{ terms}$

$$< \sum_{i=1}^n \frac{\varepsilon}{2}(x_i - x_{i-1}) + 2N_\varepsilon \|\dot{\mathcal{P}}\| \quad (h(x) \leq 1)$$

$$< \frac{\varepsilon}{2} + 2N_\varepsilon \frac{\varepsilon}{4N_\varepsilon} = \frac{\varepsilon}{2} + \frac{\varepsilon}{2} = \varepsilon$$

Since clearly $S(h; \dot{\mathcal{P}}) \geq 0$ & $\varepsilon > 0$ is arbitrary,

we have $h \in \mathcal{R}[a, b]$ & $\int_a^b h = 0$.

