

HW 5.

1* Let $m(E) < +\infty$, $m^{\#}(E) \ni f: E \rightarrow [0, \infty]$, measurable.

For each $n \in \mathbb{N}$, let

$$A_{n,k} := \left\{ x \in E : \frac{k-1}{2^n} \leq f(x) < \frac{k}{2^n} \right\}, \quad k=1, 2, \dots, 2^n, \dots, 2 \cdot 2^n, \dots, 3 \cdot 2^n, \dots, n \cdot 2^n.$$

$$B_n := \{ x \in E : n \leq f(x) \} \quad (\text{all in } m \text{ of } E = \left(\bigcup_{k=1}^{n \cdot 2^n} A_{n,k} \right) \cup B_n)$$

and let $\varphi_n := n \chi_{B_n} + \sum_{k=1}^{n \cdot 2^n} \frac{k-1}{2^n} \chi_{A_{n,k}}$, i.e.

$$\varphi_n(x) = \begin{cases} n & \text{if } n \leq f(x) \\ \frac{k-1}{2^n} & \text{if } \frac{k-1}{2^n} \leq f(x) < \frac{k}{2^n} \text{ with } k=1, 2, \dots, n \cdot 2^n \end{cases}$$

Show that $\mathcal{S}^+(E) \ni \varphi_n \uparrow f$ (pointwisely on E).

2. Let $F = \bigcup_{n=1}^{\infty} F_n$, disjoint closed sets F_1, \dots, F_n .

Let $f: F \rightarrow \mathbb{R}$ be such that $f|_{F_n}$ is cts, $\forall n$.

Show that f is cts.

3* Let $F_n \subseteq (n, n+1]$ be closed ($\mathbb{R} \setminus F_n$ open) $\forall n \in \mathbb{N}$, and let $F = \bigcup_{n \in \mathbb{N}} F_n$.

Show that $f: F \rightarrow \mathbb{R}$ is continuous if each $f|_{F_n}$ is cts. (The condition $F_n \subseteq (n, n+1]$

cannot be dropped, e.g. F_n is a singleton, F is countable - closed or not for F)

4. Let $\emptyset \neq F \subseteq \mathbb{R}$ be closed and $f: F \rightarrow \mathbb{R}$ be cts. Show (the Tietze Extension Th): f can be continuously extended to be on the whole of \mathbb{R} , via the following elementary method: let $G := \mathbb{R} \setminus F$ ($\neq \emptyset$, wlog) $= \bigcup_{n=1}^{\infty} I_n$ disjoint open intervals, by ...

Then $\bar{I}_n \setminus I_n \subseteq F \forall n$ (\bar{I}_n is the closed interval, the closure of I_n), and

f can be extended to \bar{f} so as $\bar{f}|_F = f$, & \bar{f} is "linear" on each \bar{I}_n (& its); hence \bar{f} is

cts: $\forall x_0 \in \mathbb{R}$, \bar{f} is cts at x_0 , that is

\bar{f} is right-cts (& left-cts, similarly)

For the right-cts of x_0 we assume w.l.g.
(trivially true otherwise that $x_0 \in F$).


and that $\forall \delta > 0$,

$\bigvee_{\delta}^+ (x_0) = (x_0, x_0 + \delta)$ intersects both F and G , (#)

Let $\varepsilon > 0$. By the given continuity of f on F
 $\exists \delta_0 > 0$ s.t.

$$(*) \quad |f(x) - f(x_0)| < \varepsilon \quad \forall x \in F \cap [x_0 - \delta_0, x_0 + \delta_0]$$

By (#), we take $x'_0 \in F \cap (x_0, x_0 + \delta_0)$,



and it follows that

$$(**) \quad |\bar{f}(x) - \bar{f}(x_0)| < \varepsilon \quad \forall x \in (x_0, x'_0) \quad (\text{so "right-cts"})$$

Since this is already true if $x \in F$, we only need

to consider the case $x \in G \cap (x_0, x'_0)$ with

$x \in I_n$ for some n . Now, as $x_0, x'_0 \in F$,

this entails that $x \in I_n \subseteq (x_0, x'_0)$

The end-ptz ^(\bar{x} say) of I_n are not in G so

$\bar{x} \in F$ and it follows from (*) that

$$|f(\bar{x}) - f(x_0)| < \varepsilon$$

$$\| \bar{f}(\bar{x}) - \bar{f}(x_0) \| < \varepsilon$$

and, by the def of \bar{f} on \bar{I}_n , it follows that

$$|\bar{f}(\cdot) - \bar{f}(x_0)| < \varepsilon \text{ on } I_n \ni x$$

so (**) is checked for $x \in G \cap (x_0, x'_0)$, and

hence $\forall x \in (F \cup G) \cap (x_0, x'_0) = (x_0, x'_0)$,

the whole interval.

5* Check (similarly to Q4) the left-continuity of \bar{f} at x_0 .