

MATH2060AB Homework 2

Reference Solutions

6.2.15. Since f' is bounded on I , then there exists some constant $C > 0$ such that

$$\sup_{x \in I} |f'(x)| < C.$$

Then for any $x, y \in I$, by the Mean Value Theorem, there exists some $c \in I$ such that

$$|f(x) - f(y)| = |f'(c)(x - y)| \leq C|x - y|.$$

Hence, f satisfies a Lipschitz condition on I .

6.2.20. (a) By the Mean Value Theorem, there exists some $c_1 \in (0, 1)$ such that

$$f'(c_1) = \frac{f(1) - f(0)}{1 - 0} = 1.$$

(b) By the Mean Value Theorem, there exists some $c_2 \in (1, 2)$ such that

$$f'(c_2) = \frac{f(2) - f(1)}{2 - 1} = 0.$$

(c) Using the conclusions in (a) and (b), since $f'(c_2) < 1/3 < f'(c_1)$, by Darboux's Theorem, there exists some $c_3 \in (c_1, c_2) \subset (0, 2)$ such that

$$f'(c_3) = 1/3.$$

6.3.3. For any $\epsilon > 0$, define $\delta := \epsilon$ such that for $|x - 0| < \epsilon$,

$$\left| \frac{f(x) - f(0)}{x - 0} \right| \leq |x| < \epsilon.$$

Then $f'(0) = 0$. Similarly we have $g'(0) = 0 = f'(0)$. Consider $f(x)/g(x) = \sin 1/x$. Define $x_n = 1/(2n\pi)$ and $y_n = 1/(\pi/2 + 2n\pi)$ for $n = 1, 2, \dots$. Then $\lim_{n \rightarrow \infty} f(x_n)/g(x_n) = 0 \neq 1 = \lim_{n \rightarrow \infty} f(y_n)/g(y_n)$. Hence, $\lim_{x \rightarrow 0} f(x)/g(x)$ does not exist.

6.3.5. Since $\lim_{x \rightarrow 0} |x| = 0$ and $\lim_{x \rightarrow 0} |x/\sin x| = 1$, we have $\lim_{x \rightarrow 0} |x^2/\sin x| = 0$. By $0 \leq \left| \frac{x^2 \sin \frac{1}{x}}{\sin x} \right| \leq \left| \frac{x^2}{\sin x} \right|$ and Squeezing Theorem, it holds that

$$\lim_{x \rightarrow 0} |f(x)/g(x)| = \lim_{x \rightarrow 0} \left| \frac{x^2 \sin \frac{1}{x}}{\sin x} \right| = 0,$$

which yields $\lim_{x \rightarrow 0} f(x)/g(x) = 0$. Direct calculation gives that

$$f'(x)/g'(x) = \frac{2x \sin \frac{1}{x} - \cos \frac{1}{x}}{\cos x}.$$

Let $x_n = 1/(2n\pi)$ and $y_n = 1/(\pi/2 + 2n\pi)$, then

$$\lim_{n \rightarrow \infty} f'(x_n)/g'(x_n) = -1 \neq 0 = \lim_{n \rightarrow \infty} f'(y_n)/g'(y_n).$$

Hence, $\lim_{x \rightarrow 0} f(x)/g(x)$ does not exist.

6.3.6. (a) By $\lim_{x \rightarrow 0} e^x + e^{-x} - 2 = \lim_{x \rightarrow 0} 1 - \cos x = \lim_{x \rightarrow 0} e^x - e^{-x} = \lim_{x \rightarrow 0} \sin x = 0$, $\lim_{x \rightarrow 0} e^x + e^{-x} = 2$, $\lim_{x \rightarrow 0} \cos x = -1$, we use the L'Hospital's Rules to get

$$\lim_{x \rightarrow 0} \frac{e^x + e^{-x} - 2}{1 - \cos x} = \lim_{x \rightarrow 0} \frac{e^x - e^{-x}}{\sin x} = \lim_{x \rightarrow 0} \frac{e^x + e^{-x}}{\cos x} = 2.$$

(b) Using similar arguments as in (a) and the L'Hospital's Rules, it holds that

$$\lim_{x \rightarrow 0} \frac{x^2 - \sin^2 x}{x^4} = \lim_{x \rightarrow 0} \frac{2x - 2 \sin x \cos x}{4x^3} = \lim_{x \rightarrow 0} \frac{2 - 2 \cos^2 x + 2 \sin^2 x}{12x^2} = \lim_{x \rightarrow 0} \frac{\cos x \sin x}{3x} = \lim_{x \rightarrow 0} \frac{-\sin^2 x + \cos^2 x}{3} = \frac{1}{3}.$$

6.3.10. Consider $f(x) = \frac{e^x f(x)}{e^x}$. Then by the L'Hospital's Rules,

$$\lim_{x \rightarrow \infty} f(x) = \lim_{x \rightarrow \infty} \frac{e^x f(x) + e^x f'(x)}{e^x} = L.$$

Then

$$\lim_{x \rightarrow \infty} f'(x) = \lim_{x \rightarrow \infty} (f(x) + f'(x)) - \lim_{x \rightarrow \infty} f(x) = 0.$$

6.3.10. We use the L'Hospital's Rules to get

$$\begin{aligned} & \lim_{x \rightarrow c} \frac{x^c - c^x}{x^x - c^c} \\ &= \lim_{x \rightarrow c} \frac{cx^{c-1} - c^x \ln c}{x^x(\ln x + 1)} \\ &= \frac{1 - \ln c}{\ln c + 1}. \end{aligned}$$