

(c) Bernoulli's inequality

If $\alpha > 1$, then $(1+x)^\alpha \geq 1 + \alpha x$, $\forall x > -1$.

with "equality" $\Leftrightarrow x=0$.

Pf: Consider $f(x) = (1+x)^\alpha$ on $(-1, +\infty)$.

($1+x > 0 \Rightarrow$ taking root of $1+x$ is well-defined for $\alpha \neq$ integer)

Then $f'(x) = \alpha(1+x)^{\alpha-1}$ on $(-1, +\infty)$

(We've proved this in eg 6.1.10(d) for rational α . The case of irrational α will be proved in § 8.3)

If $x > 0$, applying MVT to $f(x)$ on $[0, x]$, we have

$c \in (0, x)$ such that

$$f(x) - f(0) = f'(c)(x-0).$$

That is

$$(1+x)^\alpha - 1 = \alpha(1+c)^{\alpha-1} x.$$

Since $c > 0$ & $\alpha-1 > 0$, we have $(1+c)^{\alpha-1} > 1$.

$\therefore (1+x)^\alpha > 1 + \alpha x$ (The ineq. is strict!)

If $-1 < x < 0$, then applying MVT to $f(x)$ on $[x, 0]$,

we have $c \in (x, 0)$ such that $f(0) - f(x) = f'(c)(0-x)$

That is

$$1 - (1+x)^\alpha = \alpha(1+x)^{\alpha-1}(-x)$$

Since $-1 < x < 0$, we have $0 < 1+x < 1$

$$\Rightarrow (1+x)^{\alpha-1} < 1 \quad (\alpha-1 > 0)$$

$$\therefore 1 - (1+x)^\alpha < \alpha(-x) \quad (\text{as } -x > 0)$$

That is $(1+x)^\alpha > 1 + \alpha x$ (ineq. is strict!)

Clearly $(1+x)^\alpha = 1 + \alpha x$ for $x=0$.

Therefore $(1+x)^\alpha \geq 1 + \alpha x$, $\forall x \in [-1, +\infty)$ and

"equality $\Leftrightarrow x=0$ ". $\#$

(d) If $0 < \alpha < 1$, then $\forall a > 0 \& b > 0$, we have

$$a^\alpha b^{1-\alpha} \leq \alpha a + (1-\alpha)b.$$


with "equality $\Leftrightarrow a=b$ ". (Example of application of 1st derivative test)

(Note: for $\alpha = \frac{1}{2}$, we have $\sqrt{ab} \leq \frac{a+b}{2}$)

Pf: Consider $g(x) = \alpha x - x^\alpha$ for $x \geq 0$.

Then $g'(x) = \alpha - \alpha x^{\alpha-1} = \alpha(1 - x^{-(1-\alpha)})$ ($0 < \alpha < 1$)

$$\Rightarrow g'(x) \begin{cases} < 0 & \text{for } 0 < x < 1 \\ > 0 & \text{for } 1 < x \end{cases}$$



Hence $g(x) \geq g(1)$, $\forall x \geq 0$ and

$$g(x) = g(1) \Leftrightarrow x = 1.$$

That is, $\alpha x - x^\alpha \geq \alpha - 1$ or

$$x^\alpha \leq \alpha x + (1 - \alpha), \quad \forall x \geq 0$$

with "equality $\Leftrightarrow x = 1$ ".

Now for $a > 0, b > 0$, put $x = \frac{a}{b} > 0$ into the inequality,
we have

$$\frac{a^\alpha}{b^\alpha} \leq \frac{\alpha a}{b} + (1 - \alpha)$$

$$\Rightarrow a^\alpha b^{1-\alpha} \leq \alpha a + (1 - \alpha)b \quad \times$$

Intermediate Value Property of Derivatives (Darboux's Thm)


Lemma 6.2.11 Let I be an interval and $c \in I$.

$f: I \rightarrow \mathbb{R}$ and $f'(c)$ exists.

Then


(a) If $f'(c) > 0$, then $\exists \delta > 0$ s.t.

$$f(x) > f(c) \quad \forall x \in (c, c+\delta) \cap I$$



(b) If $f'(c) < 0$, then $\exists \delta > 0$ s.t.

$$f(x) > f(c) \quad \forall x \in (c-\delta, c) \cap I$$



Pf: (a) Since $\lim_{x \rightarrow c} \frac{f(x) - f(c)}{x - c} = f'(c) > 0$, (Thm 4.2.9 of the textbook, MATH2050)

$$\exists \delta > 0 \text{ s.t. } \frac{f(x) - f(c)}{x - c} > 0, \quad \forall x \in (c - \delta, c + \delta) \cap I$$

$$\therefore f(x) - f(c) > 0, \quad \forall x \in (c, c + \delta) \cap I.$$

(b) Similarly

$$\exists \delta > 0 \text{ s.t. } \frac{f(x) - f(c)}{x - c} < 0, \quad \forall x \in (c - \delta, c + \delta) \cap I$$

$$\therefore f(x) - f(c) > 0, \quad \forall x \in (c - \delta, c) \cap I.$$

#

Thm 6.2.12 (Darboux's Thm)

If • f is differentiable on $[a, b]$

• k is a number between $f'(a)$ and $f'(b)$, ($f'(a) \neq f'(b)$)

then $\exists c \in (a, b)$ such that

$$f'(c) = k.$$

(Remark: No continuity of f' is assumed. Hence the usual Intermediate Value Thm of continuous function doesn't apply.)

Pf: Suppose $f'(a) < f'(b)$ and $f'(a) < k < f'(b)$.

Define $g(x) = kx - f(x)$, $\forall x \in [a, b]$.

Then f differentiable \Rightarrow

g is differentiable & hence continuous on $[a, b]$

In particular, g attains a maximum value on $[a, b]$.

Note that $g'(a) = k - f'(a) > 0$.

By Lemma 6.2.11, $\exists \delta > 0$ s.t.

$$g(x) > g(a), \quad \forall x \in (a, a+\delta) \cap [a, b].$$

$\therefore a$ is not the maximum of g

Also $g'(b) = k - f'(b) < 0$, lemma 6.2.11 implies

$\exists \delta > 0$ s.t. $g(x) > g(b)$, $\forall x \in (b-\delta, b) \cap [a, b]$.

$\therefore b$ is not the maximum of g .

Together $\Rightarrow g$ attains its maximum at an

interior point $c \in (a, b)$.

Then Interior Extremum Thm (Thm 6.2.1) implies

$$0 = g'(c) = k - f'(c).$$

If $f'(b) < f'(a)$, consider $(-f)$ and we can find

similarly a $c \in (a, b)$ s.t. $f'(c) = k$. ~~✗~~

Eg 6.2.13 The signum function $g(x) = \text{sgn}(x)$ restricted on $[-1, 1]$:

$$g(x) = \begin{cases} 1, & 0 < x \leq 1 \\ 0, & x = 0 \\ -1, & -1 \leq x < 0 \end{cases}$$

doesn't satisfy the intermediate value property,

$(1 = g(1), -1 = g(-1), \& -1 < \frac{1}{2} < 1, \text{ but no } x \in (-1, 1) \text{ s.t. } g(x) = \frac{1}{2})$

Therefore $g(x) \neq f'(x)$ for any differentiable function f on $[-1, 1]$.

(i.e. The differential eq $\frac{df}{dx} = g$ has no solution on $[-1, 1]$)