

# Fourier Analysis

Mar 14, 2024

## Review.

- A cts but nowhere diff function on  $\mathbb{R}$ .

Let  $\alpha \in (0, 1)$ . set

$$f_\alpha(x) = \sum_{n=0}^{\infty} 2^{-n\alpha} e^{i2^n x}, \quad x \in \mathbb{R}.$$


Then  $f_\alpha$  is cts but nowhere differentiable.

- By modifying the proof of  $f_\alpha$  being nowhere diff slightly, one can show the real and imaginary parts of  $f_\alpha$  are also nowhere diff. That is,

$$\sum_{n=0}^{\infty} 2^{-n\alpha} \cos(2^n x), \quad \sum_{n=0}^{\infty} 2^{-n\alpha} \sin(2^n x)$$

are nowhere diff. (check the outline proof in the text book).

## §4.4 Heat equation on the circle.



heat conduction on the circle.

$$\theta \in [0, 2\pi)$$

$$\theta = 2\pi x, \quad x \in [0, 1).$$

Let  $U = U(x, t)$  be the temperature at point  $x$  and time  $t$ .

Then  $u$  satisfies

$$\frac{\partial u}{\partial t} = c \cdot \frac{\partial^2 u}{\partial x^2}, \quad x \in [0, 1], \quad t > 0.$$

By scaling the time  $t$  if necessary, we may obtain a standard heat equation

$$\frac{\partial u}{\partial t} = \frac{\partial^2 u}{\partial x^2}, \quad x \in [0, 1], \quad t > 0. \quad (*)$$

The function  $u = U(x, t)$  can be extended to a function on  $\mathbb{R} \times (0, \infty)$ , which is 1-periodic in  $x$ .

Here we would like to find a solution  <sup>$u$</sup>  of  $(*)$ , in the mean time, we ask  $u$  to satisfy an initial condition

$$u(x, 0) = f(x), \quad (**)$$

We first find <sup>some</sup> special solutions  $U(x, t) = A(x)B(t)$  of  $(*)$ .

Plugging  $U = A(x)B(t)$  into  $(*)$ , we obtain

$$B'(t) A(x) = A''(x) B(t)$$

So

$$\frac{A''(x)}{A(x)} = \frac{B'(t)}{B(t)} = \lambda.$$

From  $A''(x) - \lambda A(x) = 0$ , we obtain

$$A(x) = \begin{cases} c_1 e^{\sqrt{\lambda} x} + c_2 e^{-\sqrt{\lambda} x} & \text{if } \lambda > 0 \\ c_1 x + c_2 & \text{if } \lambda = 0 \\ c_1 e^{i\sqrt{-\lambda} x} + c_2 e^{-i\sqrt{-\lambda} x} & \text{if } \lambda < 0 \end{cases}$$

To obtain a 1-periodic solution  $A(x)$ , we must

have  $A(x)$  be a const or

$$A(x) = c_1 e^{in2\pi x} + c_2 \cdot e^{-in2\pi x} \quad \text{for some } n \in \mathbb{Z}$$

corresponding to  $\lambda = -(2\pi n)^2 = -4\pi^2 n^2$ .

$$\frac{B'(t)}{B(t)} = -4\pi^2 n^2 \Rightarrow B(t) = c \cdot e^{-4\pi^2 n^2 t}$$

Hence our special solution should be of <sup>the</sup> form

$$\begin{aligned} u(x,t) &= (c_1 e^{i2\pi n x} + c_2 e^{-i2\pi n x}) \cdot c \cdot e^{-4\pi^2 n^2 t} \\ &= (a_n e^{i2\pi n x} + a_{-n} e^{-i2\pi n x}) e^{-4\pi^2 n^2 t} \end{aligned}$$

Using the superposition of these special solutions

we would like to find  $a_n$  ( $n \in \mathbb{Z}$ ) such that

$$u(x,t) = \sum_{n=-\infty}^{\infty} a_n e^{i2\pi n x} \cdot e^{-4\pi^2 n^2 t} \quad (***)$$