

# Introductory Probability

23-09-07

## Chapter 2 Axioms of probability

### 1. Introduction.

- Probability is a math area dealing with random behaviors.
- It has a history of more than 300 years in the study.
- It came from gambling in the early stage, and gamings of chance.

### 2. Random experiments, outcomes, sample space, events.

Random experiments / outcomes.

Example: ① Toss a coin to get a head or a tail.

② Roll a dice to see the number of the top face.

③ Measure the height of a randomly chosen student in the campus.

Def. (sample space). The set of all <sup>possible</sup> outcomes of an experiment is called the sample space of the experiment.

Usually, We use  $S$  to denote the sample space.

Example ① Toss a coin once.

$$S = \{H, T\}.$$

Toss a coin twice.

$$S = \{HH, HT, TH, TT\}$$

② Roll a dice once

$$S = \{1, 2, 3, 4, 5, 6\}.$$

Roll a dice 3 times.

$$S = \{(i, j, k) : i, j, k \in \{1, 2, 3, 4, 5, 6\}\}.$$

③ height of a randomly chosen student (in meters)

$$S = \{0 < x < \infty\} = (0, \infty)$$

Def (event) Let  $S$  be the sample space of an experiment.

Every subset  $E$  of  $S$  is called an event.

If an outcome of the experiment is contained in the event  $E$ , then we say that  $E$  has occurred.

- Basic operations on events.

Union:  $E \cup F$


Intersection:  $E \cap F$

Complement  $E^c = S \setminus E$


- $\emptyset$  Null event.

We say two events  $E, F$  are mutually exclusive if  $E \cap F = \emptyset$ .


- Venn diagram:



$E \cap F$



$E^c$



$(E \cap F) \cup (F \cap G)$

• Laws.

$$(i) \quad E \cup F = F \cup E, \quad E \cap F = F \cap E \quad \text{commutative laws}$$

$$E \cap (F \cup G) = (E \cap F) \cup (E \cap G) \quad \text{distributive law}$$

$$\left. \begin{aligned} E \cup (F \cap G) &= (E \cup F) \cap G \\ E \cap (F \cup G) &= (E \cap F) \cup G. \end{aligned} \right\} \text{associative laws}$$

(ii) De Morgan's laws

$$\left( \bigcup_{n=1}^{\infty} E_n \right)^c = \bigcap_{n=1}^{\infty} E_n^c$$

$$\left( \bigcap_{n=1}^{\infty} E_n \right)^c = \bigcup_{n=1}^{\infty} E_n^c.$$

Pf. Let us prove the first equality in (ii)

$$x \in \left( \bigcup_{n=1}^{\infty} E_n \right)^c$$

$$\Leftrightarrow x \in S, \quad x \notin \bigcup_{n=1}^{\infty} E_n$$

$$\Leftrightarrow x \in S, \quad x \notin E_n \text{ for } n=1, 2, \dots$$

$$\Leftrightarrow x \in E_n^c \text{ for } n=1, 2, \dots$$

$$\Leftrightarrow x \in \bigcap_{n=1}^{\infty} E_n^c$$

$$\text{Hence } \left( \bigcup_{n=1}^{\infty} E_n \right)^c = \bigcap_{n=1}^{\infty} E_n^c. \quad \square$$

## §2.3. Axioms of probability.

Q: How can we define the prob. of an event?

An intuitive approach:

repeat the random experiment  $n$  times.

let  $n(E)$  be the times that an event  $E$  occurs

$$\text{Let } p(E) = \lim_{n \rightarrow \infty} \frac{n(E)}{n}.$$

Drawbacks : ① why does the limit exist?

② Even if the limit exist, why is it independent of the experiments?

The axiomatic approach to prob. (by Kolmogorov)

Def. (Prob. of an event).

Let  $S$  be the sample space of a random experiment.

A probability  $P$  on  $S$  is a function that assigns a value to each event  $E$  such that the following 3 axioms hold:

Axiom 1:  $0 \leq P(E) \leq 1$ ,  $\forall$  event  $E$ .

Axiom 2:  $P(S) = 1$ .

Axiom 3: If  $E_1, E_2, \dots$  are a sequence

of events which are mutually exclusive,

then

$$P\left(\bigcup_{n=1}^{\infty} E_n\right) = \sum_{n=1}^{\infty} P(E_n).$$

(Countable additivity of prob.)

§ 2.4. Some properties of probability.

Prop 1.  $P(\emptyset) = 0$ .

Pf. Let  $E_1 = S$ , and  $E_n = \emptyset$  for  $n=2, 3, \dots$

Then  $E_1, E_2, \dots$ , are mutually exclusive.

By Axiom 3,

$$\begin{aligned} P\left(\bigcup_{n=1}^{\infty} E_n\right) &= \sum_{n=1}^{\infty} P(E_n) \\ &= P(E_1) + P(E_2) + \dots \\ &= P(S) + P(\emptyset) + P(\emptyset) + \dots \end{aligned}$$

LHS  $\leq 1$ , RHS  $\leq 1$  only occurs when  $P(\emptyset) = 0$ .  $\square$

Prop 2. (finite additivity)

Let  $E_1, E_2, \dots, E_n$  be mutually exclusive events.

Then

$$P\left(\bigcup_{k=1}^n E_k\right) = \sum_{k=1}^n P(E_k)$$

Pf. Define  $E_j = \emptyset$  for  $j = n+1, n+2, \dots$

By Axiom 3,

$$\begin{aligned} P\left(\bigcup_{k=1}^{\infty} E_k\right) &= \sum_{k=1}^{\infty} P(E_k) \\ &= \sum_{k=1}^n P(E_k) + \sum_{k=n+1}^{\infty} P(E_k) \\ &= \sum_{k=1}^n P(E_k) \quad (\text{since } P(E_{n+1}) \\ &\quad = P(E_{n+2}) = \dots = 0 \\ &\quad \text{by Prop 1}) \end{aligned}$$

Now the proposition follows from

$$\bigcup_{k=1}^n E_k = \bigcup_{k=1}^{\infty} E_k.$$





Prop 3.  $P(E^c) = 1 - P(E)$ .

Pf. Notice that

$$S = E^c \cup E \cup \emptyset \cup \emptyset \dots$$

By Axiom 3 and Prop 1,  
Axiom 2

$$1 = P(S) = P(E^c) + P(E).$$

□

Prop 4 Let  $E, F$  be two events. Then

$$P(E \cup F) = P(E) + P(F) - P(E \cap F).$$

Pf.

$$E \cup F = E \cup (F \setminus E)$$


since  $E \cap (F \setminus E) = \emptyset$ , so by Axiom 3,

$$P(E \cup F) = P(E) + P(F \setminus E). \quad \textcircled{1}$$

Now we consider  $P(F|E)$ .

Notice that

$$F = (F \setminus E) \cup (E \cap F)$$



red  $\leftrightarrow E \cap F$

blue  $\leftrightarrow F \setminus E$ .

Using Axiom 3 again,

$$P(F) = P(F \setminus E) + P(E \cap F)$$

hence

$$P(F|E) = P(F) - P(E \cap F) \quad (2)$$

Plugging (2) into (1) yields the desired identity.  $\square$


Prop. 5. Suppose  $F \subset E$ . Then

$$P(F) \leq P(E).$$

Pf. Since  $F \subset E$ ,

$$E = F \cup (E \setminus F)$$

(disjoint).



By Prop 2,

$$P(E) = P(F) + P(E \setminus F).$$

Since  $P(E \setminus F) \geq 0$  by Axiom 1, it follows that

$$P(E) \geq P(F).$$



Prop 6. Let  $E_1, E_2, \dots$  be a sequence of events.

Then

$$P\left(\bigcup_{n=1}^{\infty} E_n\right) \leq \sum_{n=1}^{\infty} P(E_n).$$

(Countable sub-additivity of prob.)

Proof. First we write  $\bigcup_{n=1}^{\infty} E_n$  as the union of some disjoint events. To do so,

write

$$F_1 = E_1$$

$$F_2 = E_2 \setminus E_1$$

$$F_3 = E_3 \setminus (E_1 \cup E_2),$$

...

$$F_n = E_n \setminus \left(\bigcup_{i=1}^{n-1} E_i\right),$$

...

Then the following properties hold:

$$(1) F_n \subset E_n, \quad n=1, \dots,$$

(2)  $F_1, F_2, \dots$  are mutually exclusive.

$$(3) \bigcup_{i=1}^n F_i = \bigcup_{i=1}^n E_i$$

$$(4) \bigcup_{i=1}^{\infty} F_i = \bigcup_{i=1}^{\infty} E_i$$

(1) and (2) are easy to see. Below we only prove (4).  
The proof of (3) is similar.

To show (4), recall that  $F_i \subset E_i$  so

$$\bigcup_{i=1}^{\infty} F_i \subset \bigcup_{i=1}^{\infty} E_i.$$

To prove  $\bigcup_{i=1}^{\infty} F_i \supset \bigcup_{i=1}^{\infty} E_i$ ,

let  $x \in \bigcup_{i=1}^{\infty} E_i$ . Then  $x \in E_i$  for some  $i$ .

Let  $i_0$  be the smallest integer such that

$$\begin{array}{c} x \in E_{i_0} \\ \text{Then } x \in E_{i_0} \setminus \bigcup_{j=1}^{i_0-1} E_j = F_{i_0}. \end{array}$$

It follows that

$$\bigcup_{i=1}^{\infty} E_i \subset \bigcup_{i=1}^{\infty} F_i,$$

which proves (\*)

Now using Axiom 3 to  $P\left(\bigcup_{n=1}^{\infty} F_n\right)$

we have

$$\begin{aligned} P\left(\bigcup_{n=1}^{\infty} F_n\right) &= \sum_{n=1}^{\infty} P(F_n) \\ &\leq \sum_{n=1}^{\infty} P(E_n), \end{aligned}$$

and we are done since

$$P\left(\bigcup_{n=1}^{\infty} F_n\right) = P\left(\bigcup_{n=1}^{\infty} E_n\right). \quad \square$$