

## Exercise 8

Standard notations are in force. Many problems are taken from [R].

(1)

$$\Phi(t) = \int_X |f + tg|^p d\mu$$

is differentiable at  $t = 0$  and

$$\Phi'(0) = p \int_X |f|^{p-2} fg d\mu.$$

Hint: Use the convexity of  $t \mapsto |f + tg|^p$  to get

$$|f + tg|^p - |f|^p \leq t(|f + g|^p - |f|^p), \quad t > 0$$

and a similar estimate for  $t < 0$ .

(2) Suppose  $f$  is a measurable function on  $X$ ,  $\mu$  is a positive measure on  $X$ , and

$$\varphi(p) = \int_X |f|^p d\mu = \|f\|_p^p \quad (0 < p < \infty).$$

Let  $E = \{p : \varphi(p) < \infty\}$ . Assume  $\|f\|_\infty > 0$ .

- (a) If  $r < p < s$ ,  $r \in E$ , and  $s \in E$ , prove that  $p \in E$ .
- (b) Prove that  $\log \varphi$  is convex in the interior of  $E$  and that  $\varphi$  is continuous on  $E$ .
- (c) By (a),  $E$  is connected. Is  $E$  necessarily open? Closed? Can  $E$  consist of a single point? Can  $E$  be any connected subset of  $(0, \infty)$ ?
- (d) If  $r < p < s$ , prove that  $\|f\|_p \leq \max(\|f\|_r, \|f\|_s)$ . Show that this implies the inclusion  $L^r(\mu) \cap L^s(\mu) \subset L^p(\mu)$ .
- (e) Assume that  $\|f\|_r < \infty$  for some  $r < \infty$  and prove that

$$\|f\|_p \rightarrow \|f\|_\infty \quad \text{as } p \rightarrow \infty.$$

(3) Assume, in addition to the hypothesis of the previous problem, that

$$\mu(X) = 1.$$

- (a) Prove that  $\|f\|_r \leq \|f\|_s$  if  $0 < r < s \leq \infty$ .

- (b) Under what conditions does it happen that  $0 < r < s \leq \infty$  and  $\|f\|_r = \|f\|_s < \infty$  ?
- (c) Prove that  $L^r(\mu) \supset L^s(\mu)$  if  $0 < r < s$ . Under what conditions do these two spaces contain the same functions?
- (d) Assume that  $\|f\|_r < \infty$  for some  $r > 0$ , and prove that

$$\lim_{p \rightarrow 0} \|f\|_p = \exp \left\{ \int_X \log |f| d\mu \right\}$$

if  $\exp\{-\infty\}$  is defined to be 0.

- (4) For some measures, the relation  $r < s$  implies  $L^r(\mu) \subset L^s(\mu)$ ; for others, the inclusion is reversed; and there are some for which  $L^r(\mu)$  does not contain  $L^s(\mu)$  if  $r \neq s$ . Give examples of these situations, and find conditions on  $\mu$  under which these situations will occur.
- (5) Suppose  $\mu(\Omega) = 1$ , and suppose  $f$  and  $g$  are positive measurable functions on  $\Omega$  such that  $fg \geq 1$ . Prove that

$$\int_{\Omega} f d\mu \cdot \int_{\Omega} g d\mu \geq 1.$$

- (6) Suppose  $\mu(\Omega) = 1$  and  $h : \Omega \rightarrow [0, \infty]$  is measurable. If

$$A = \int_{\Omega} h d\mu,$$

prove that

$$\sqrt{1 + A^2} \leq \int_{\Omega} \sqrt{1 + h^2} d\mu \leq 1 + A.$$

If  $\mu$  is Lebesgue measure on  $[0, 1]$  and if  $h$  is continuous,  $h = f'$ , the above inequalities have a simple geometric interpretation. From this, conjecture (for general  $\Omega$ ) under what conditions on  $h$  equality can hold in either of the above inequalities, and prove your conjecture.

- (7) Optional. Suppose  $1 < p < \infty$ ,  $f \in L^p = L^p((0, \infty))$ , relative to Lebesgue measure, and

$$F(x) = \frac{1}{x} \int_0^x f(t) dt \quad (0 < x < \infty).$$

- (a) Prove Hardy's inequality

$$\|F\|_p \leq \frac{p}{p-1} \|f\|_p$$

which shows that the mapping  $f \rightarrow F$  carries  $L^p$  into  $L^p$ .

- (b) Prove that equality holds only if  $f = 0$  a.e.

- (c) Prove that the constant  $\frac{p}{p-1}$  cannot be replaced by a smaller one.
- (d) If  $f > 0$  and  $f \in L^1$ , prove that  $F \notin L^1$ .

Suggestions: (a) Assume first that  $f \geq 0$  and  $f \in C_c((0, \infty))$ . Integration by parts gives

$$\int_0^\infty F^p(x) dx = -p \int_0^\infty F^{p-1}(x) x F'(x) dx.$$

Note that  $x F' = f - F$ , and apply Hölder's inequality to  $\int F^{p-1} f$ . Then derive the general case.

- (c) Take  $f(x) = x^{-1/p}$  on  $[1, A]$ ,  $f(x) = 0$  elsewhere, for large  $A$ . See also Exercise 14, Chap. 8 in [R].