

13.5 Calculating Gradients

In Exercises 1–6, find the gradient of the function at the given point. Then sketch the gradient, together with the level curve that passes through the point.

1. $f(x, y) = y - x$, $(2, 1)$ 2. $f(x, y) = \ln(x^2 + y^2)$, $(1, 1)$

3. $g(x, y) = xy^2$, $(2, -1)$ 4. $g(x, y) = \frac{x^2}{2} - \frac{y^2}{2}$, $(\sqrt{2}, 1)$


Sol'n: $\left. \frac{\partial g}{\partial x} \right|_{(2, -1)} = y^2 \Big|_{(2, -1)} = 1$, $\left. \frac{\partial g}{\partial y} \right|_{(2, -1)} = 2xy \Big|_{(2, -1)} = -4$

So $\vec{\nabla} g(2, -1) = (1, -4)$.

$g(2, -1) = 2(-1)^2 = 2$.

So level curve that passes through the point is

$xy^2 = 2 \Rightarrow x = \frac{2}{y^2}$



13.5

Finding Directional Derivatives

In Exercises 11–18, find the derivative of the function at P_0 in the direction of $\mathbf{v}$.

11. $f(x, y) = 2xy - 3y^2$, $P_0(5, 5)$, $\mathbf{v} = 4\mathbf{i} + 3\mathbf{j}$

12. $f(x, y) = 2x^2 + y^2$, $P_0(-1, 1)$, $\mathbf{v} = 3\mathbf{i} - 4\mathbf{j}$

13. $g(x, y) = \frac{x-y}{xy+2}$, $P_0(1, -1)$, $\mathbf{v} = 12\mathbf{i} + 5\mathbf{j}$

Sol'n: We first need to take $\vec{u} = \frac{\vec{v}}{\|\vec{v}\|}$.

$$\|\vec{v}\| = \sqrt{12^2 + 5^2} = 13 \quad \text{So } \vec{u} = \frac{1}{13}(12, 5).$$

Then $D_{\vec{u}}f(P_0) = \vec{\nabla}f(P_0) \cdot \vec{u}$.

$$\left. \frac{\partial f}{\partial x} \right|_{(1, -1)} = \left. \frac{(1)(xy+2) - (x-y)(y)}{(xy+2)^2} \right|_{(1, -1)} = \left. \frac{xy+2 - xy + y^2}{(xy+2)^2} \right|_{(1, -1)}$$

$$= \frac{2 + (-1)^2}{(1(-1)+2)^2} = 3$$

$$\begin{aligned} \frac{\partial f}{\partial y} \Big|_{(1,-1)} &= \frac{(-1)(xy+2) - (x-y)(x)}{(xy+2)^2} \Big|_{(1,-1)} = \frac{-xy-2-x^2+xy}{(xy+2)^2} \Big|_{(1,-1)} \\ &= \frac{-2-(1)^2}{((1)(-1)+2)^2} = -3 \end{aligned}$$

$$\text{So } D_{\vec{u}} f(P_0) = \vec{\nabla} f(P_0) \cdot \vec{u} = (3, -3) \cdot \frac{1}{13}(12, 5) = \boxed{\frac{21}{13}}.$$

13.5

Theory and Examples

29. Let $f(x, y) = x^2 - xy + y^2 - y$. Find the directions $\mathbf{u}$ and the values of $D_{\mathbf{u}}f(1, -1)$ for which

- a. $D_{\mathbf{u}}f(1, -1)$ is largest b. $D_{\mathbf{u}}f(1, -1)$ is smallest
 c. $D_{\mathbf{u}}f(1, -1) = 0$ d. $D_{\mathbf{u}}f(1, -1) = 4$
 e. $D_{\mathbf{u}}f(1, -1) = -3$

Sol'n: a) As discussed in lecture $-\|\vec{\nabla}f(\vec{a})\| \leq D_{\vec{u}}f(\vec{a}) \leq \|\vec{\nabla}f(\vec{a})\|$
 with equality when $\mathbf{u} = \pm \frac{\vec{\nabla}f(\vec{a})}{\|\vec{\nabla}f(\vec{a})\|}$ resp. for a unit vector.

compute $\vec{\nabla}f(1, -1) = \left[\frac{\partial f}{\partial x} \Big|_{(1, -1)} \quad \frac{\partial f}{\partial y} \Big|_{(1, -1)} \right]$

$$\frac{\partial f}{\partial x} \Big|_{(1, -1)} = 2x - y \Big|_{(1, -1)} = 2 - (-1) = 3, \quad \frac{\partial f}{\partial y} \Big|_{(1, -1)} = -x + 2y - 1 \Big|_{(1, -1)} = -4.$$

so $\vec{\nabla}f(1, -1) = [3 \ -4]$.

so $\|\vec{\nabla}f(1, -1)\| = \sqrt{3^2 + (-4)^2} = 5$

So $D_{\vec{u}}f(1,-1)$ is largest when $\boxed{\vec{u} = \left(\frac{3}{5}, -\frac{4}{5}\right)}$

b) $D_{\vec{u}}f(1,-1)$ is smallest when $\boxed{\vec{u} = \left(-\frac{3}{5}, \frac{4}{5}\right)}$

c) For $\vec{u} = (u_1, u_2)$ unit vector,

$$D_{\vec{u}}f(1,-1) = \vec{\nabla}f(1,-1) \cdot \vec{u} = (3, -4) \cdot (u_1, u_2) = 3u_1 - 4u_2$$

$$0 = 3u_1 - 4u_2$$

So can take $u_1 = \frac{4}{5}, u_2 = \frac{3}{5} \Rightarrow \boxed{\vec{u} = \left(\frac{4}{5}, \frac{3}{5}\right)}$

$u_1 = -\frac{4}{5}, u_2 = -\frac{3}{5} \Rightarrow \boxed{\vec{u} = \left(-\frac{4}{5}, -\frac{3}{5}\right)}$

d) $4 = 3u_1 - 4u_2$

$$3u_1 = 4 + 4u_2$$

$$u_1 = \frac{4}{3} + \frac{4}{3}u_2$$

$$1 = \sqrt{u_1^2 + u_2^2} \Rightarrow 1 = u_1^2 + u_2^2 \Rightarrow 1 = \left(\frac{4}{3} + \frac{4}{3}u_2\right)^2 + u_2^2$$

$$\Rightarrow \frac{25u_2^2}{9} + \frac{32u_2}{9} + \frac{7}{9} = 0 \Rightarrow u_2 = -1, -\frac{7}{25}$$

When $u_2 = -1$, we have $u_1 = 0 \Rightarrow \boxed{\vec{u} = (0, -1)}$

$u_2 = -\frac{7}{25}$, we have $4 = 3u_1 - 4\left(-\frac{7}{25}\right) \Rightarrow 3u_1 = \frac{72}{25}$
 $\Rightarrow u_1 = \frac{24}{25}$.

$\Rightarrow \boxed{\vec{u} = \left(\frac{24}{25}, -\frac{7}{25}\right)}$

e) $-3 = 3u_1 - 4u_2$.

$1 = u_1^2 + u_2^2 \Rightarrow 1 = u_1^2 + \left(\frac{3}{4}u_1 + \frac{3}{4}\right)^2 = u_1^2 + \frac{9u_1^2}{16} + \frac{9u_1}{8} + \frac{9}{16}$

$4u_2 = 3u_1 + 3$.

$u_2 = \frac{3}{4}u_1 + \frac{3}{4}$.

$= \frac{25u_1^2}{16} + \frac{9u_1}{8} + \frac{9}{16}$

$\Rightarrow \frac{25u_1^2}{16} + \frac{9u_1}{8} - \frac{7}{16} = 0$

$\Rightarrow u_1 = -1, \frac{7}{25}$.

When $u_1 = -1$, $u_2 = 0 \Rightarrow \boxed{\vec{u} = (-1, 0)}$

$u_1 = \frac{7}{25}$, $u_2 = \frac{3}{4}\left(\frac{7}{25}\right) + \frac{3}{4} = \frac{24}{25} \Rightarrow \boxed{\vec{u} = \left(\frac{7}{25}, \frac{24}{25}\right)}$

23. By about how much will

13.6

$$g(x, y, z) = x + x \cos z - y \sin z + y$$

change if the point $P(x, y, z)$ moves from $P_0(2, -1, 0)$ a distance of $ds = 0.2$ unit toward the point $P_1(0, 1, 2)$?

Sol'n: $dg = \sum_{i=1}^3 \frac{\partial g}{\partial x_i}(P_0) dx_i = (\nabla g|_{P_0} \cdot \vec{u}) ds$

$$\vec{u} = \frac{P_1 - P_0}{\|P_1 - P_0\|} = \frac{(0, 1, 2) - (2, -1, 0)}{\|(0, 1, 2) - (2, -1, 0)\|} = \frac{(-2, 2, 2)}{\sqrt{(-2)^2 + 2^2 + 2^2}} = \left(\frac{-1}{\sqrt{3}}, \frac{1}{\sqrt{3}}, \frac{1}{\sqrt{3}}\right)$$

$$\frac{\partial g}{\partial x} \Big|_{(2, -1, 0)} = (1 + \cos z) \Big|_{(2, -1, 0)} = 1 + \cos 0 = 2.$$

$$\frac{\partial g}{\partial y} \Big|_{(2, -1, 0)} = (-\sin z + 1) \Big|_{(2, -1, 0)} = 1$$

$$\frac{\partial g}{\partial z} \Big|_{(2, -1, 0)} = (-x \sin z - y \cos z) \Big|_{(2, -1, 0)} = -2 \overset{0}{\sin 0} + 1 \cos 0 = 1.$$

$$\text{So } dg = \left((2, 1, 1) \cdot \left(\frac{-1}{\sqrt{3}}, \frac{1}{\sqrt{3}}, \frac{1}{\sqrt{3}}\right) \right) 0.2 = \left(\frac{-2}{\sqrt{3}} + \frac{1}{\sqrt{3}} + \frac{1}{\sqrt{3}}\right) 0.2 = 0 \cdot 0.2 = \boxed{0}.$$

13.6

Linearizations for Three Variables

Find the linearizations $L(x, y, z)$ of the functions in Exercises 41–46 at the given points.

45. $f(x, y, z) = e^x + \cos(y + z)$ at

a. $(0, 0, 0)$

b. $(0, \frac{\pi}{2}, 0)$

c. $(0, \frac{\pi}{4}, \frac{\pi}{4})$

Sol'n: Linearization: $L(\vec{w}) = f(\vec{a}) + \vec{\nabla}f(\vec{a}) \cdot (\vec{w} - \vec{a})$.

$$\vec{\nabla}f = \left[\frac{\partial f}{\partial x} \quad \frac{\partial f}{\partial y} \quad \frac{\partial f}{\partial z} \right] = [e^x \quad -\sin(y+z) \quad -\sin(y+z)]$$

a) $f(\vec{a}) = f(0, 0, 0) = e^0 + \cos(0+0) = 1 + 1 = 2$.

$$\vec{\nabla}f(0, 0, 0) = [e^x \quad -\sin(y+z) \quad -\sin(y+z)]|_{(0,0,0)}$$

$$= [e^0 \quad -\sin(0+0) \quad -\sin(0+0)]$$

$$= [1 \quad 0 \quad 0]$$

$$\vec{w} = (x, y, z).$$

$$\text{So } L(\vec{w}) = 2 + [1 \ 0 \ 0] \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \boxed{2+x}$$

$$b) f(\vec{w}) = f(0, \frac{\pi}{2}, 0) = e^0 + \cos(\frac{\pi}{2} + 0) = 1 + \cos(\frac{\pi}{2}) = 1.$$

$$\vec{\nabla} f(0, \frac{\pi}{2}, 0) = [e^0 \quad -\sin(\frac{\pi}{2} + 0) \quad -\sin(\frac{\pi}{2} + 0)] = [1 \quad -1 \quad -1].$$

$$\vec{\nabla} f(0, \frac{\pi}{2}, 0) \cdot (x, y - \frac{\pi}{2}, z) = x - (y - \frac{\pi}{2}) - z = x - y - z + \frac{\pi}{2}$$

$$\text{So } L(\vec{w}) = \boxed{x - y - z + \frac{\pi}{2} + 1}$$

$$c) f(\vec{w}) = f(0, \frac{\pi}{4}, \frac{\pi}{4}) = e^0 + \cos(\frac{\pi}{4} + \frac{\pi}{4}) = 1.$$

$$\vec{\nabla} f(0, \frac{\pi}{4}, \frac{\pi}{4}) = [e^0 \quad -\sin(\frac{\pi}{4} + \frac{\pi}{4}) \quad -\sin(\frac{\pi}{4} + \frac{\pi}{4})] = [1 \quad -1 \quad -1].$$

$$\vec{\nabla} f(0, \frac{\pi}{4}, \frac{\pi}{4}) \cdot (x, y - \frac{\pi}{4}, z - \frac{\pi}{4}) = x + \frac{\pi}{4} - y + \frac{\pi}{4} - z,$$

$$\text{So } L(\vec{w}) = \boxed{x - y - z + \frac{\pi}{2} + 1}$$