

**THE CHINESE UNIVERSITY OF HONG KONG**  
**Department of Mathematics**  
**MATH 3030 Abstract Algebra 2024-25**  
**Homework 9 solutions**

**Compulsory Part**

1. Let  $R$  be a commutative ring and  $I$  an ideal of  $R$ . Show that the set  $\sqrt{I}$  of all  $a \in R$ , such that  $a^n \in I$  for some  $n \in \mathbb{Z}^+$ , is an ideal of  $R$ , called the **radical** of  $I$ .

**Answer.** Clearly,  $0^1 = 0 \in I$ , so  $0 \in \sqrt{I}$ .

Let  $x, y \in \sqrt{I}$ . Then  $x^m, y^n \in I$  for some  $m, n > 0$ . Then  $(x+y)^{m+n} = \sum_{k=0}^{m+n} \binom{m+n}{k} x^k y^{m+n-k}$ . But for any  $0 \leq k \leq m+n$ , either  $k \geq m$  or  $m+n-k \geq n$ . Hence either  $x^k \in I$  or  $y^{m+n-k} \in I$ . It follows that each summand and hence  $(x+y)^{m+n}$  is in  $I$ . Then  $x+y \in \sqrt{I}$ .

Let  $a \in R, x \in \sqrt{I}$ . Then  $x^m \in I$  for some  $m > 0$ . Then  $(ax)^m = a^m x^m \in I$ . Therefore,  $ax \in \sqrt{I}$ .

It follows that  $\sqrt{I}$  is an ideal.

2. Show by examples that for proper ideals  $I$  of a commutative ring  $R$ ,

(a)  $\sqrt{I}$  need not equal  $I$ .

(b)  $\sqrt{I}$  may equal  $I$ .

**Answer.** (a) Take  $R = \mathbb{Z}[x]/\langle x^2 \rangle$  and  $I = 0$ . Then  $x \in \sqrt{I} - I$  because  $x^2 \in \langle x^2 \rangle$ .

(b) Take  $R = \mathbb{Z}$  and  $I = 0$ . Then  $\sqrt{I} = I$  because  $\mathbb{Z}$  is an integral domain.

3. Prove that  $\mathbb{Z}[x]$  is not a PID by showing that the ideal  $\langle 2, x \rangle$  is not principal.

**Answer.** Suppose the ideal  $\langle 2, x \rangle$  is principal ideal  $\langle p(x) \rangle$ . Since  $2 \in \langle p(x) \rangle$ ,  $2 = p(x)q(x)$  for some  $q(x) \in \mathbb{Z}[x]$ . Since  $\mathbb{Z}$  is an integral domain, we have  $\deg(p(x)q(x)) = \deg(p(x)) + \deg(q(x))$ . Thus, both  $p(x)$  and  $q(x)$  must be constant. The only possible options for  $p(x)$  are  $\{\pm 1, \pm 2\}$ . However, This ideals obviously either contain units or not contain  $x$ .

4. Let  $D$  be an integral domain. Show that, for  $k = 1, \dots, n$ , the ideal  $\langle x_1, \dots, x_k \rangle$  is prime in  $D[x_1, \dots, x_n]$ .

**Answer.** Note that we have the isomorphism:  $D[x_1, \dots, x_n]/\langle x_1, \dots, x_k \rangle \simeq D[x_{k+1}, \dots, x_n]$ . We conclude that  $\langle x_1, \dots, x_n \rangle$  is a prime ideal since  $D[x_{k+1}, \dots, x_n]$  is an integral domain.

5. Let  $\varphi : R \rightarrow S$  be a homomorphism of commutative rings, and let  $I \subset S$  be an ideal. Prove that if  $I$  is a prime ideal in  $S$ , then  $\varphi^{-1}(I)$  is a prime ideal in  $R$ . Show by giving an example that, however,  $\varphi^{-1}(I)$  is not necessarily maximal when  $I$  is maximal.

**Answer.** Let  $\varphi : R \rightarrow S$  be a homomorphism of commutative rings, and let  $I \subset S$  be an ideal. Suppose  $I$  is prime in  $S$ . Let  $x, y \in R$  with  $xy \in \varphi^{-1}(I)$ . Then  $\varphi(x)\varphi(y) = \varphi(xy) \in I$ . Since  $I$  is prime,  $\varphi(x) \in I$  or  $\varphi(y) \in I$ . Therefore,  $x \in \varphi^{-1}(I)$  or  $y \in \varphi^{-1}(I)$ . It follows that  $\varphi^{-1}(I)$  is a prime ideal in  $R$ .

Consider the embedding  $\mathbb{Z} \rightarrow \mathbb{Q}$ ,  $\{0\}$  is a maximal ideal in  $\mathbb{Q}$ , but its preimage  $\{0\}$  is not a maximal ideal in  $\mathbb{Z}$ .

6. Show that every prime ideal in a *finite* commutative ring  $R$  is a maximal ideal.

**Answer.** Let  $R$  be a finite commutative ring. Let  $P$  be a prime ideal in  $R$ . Then  $R/P$  is a finite integral domain. Let  $x \in R/P$  be a nonzero element. Let  $m_x : R/P \rightarrow R/P$  be defined by  $m_x(y) = xy$ . If  $m_x(y) = m_x(y')$ , then  $x(y - y') = m_x(y - y') = 0$ . Then  $y - y' = 0$  since  $R/P$  is an integral domain. Therefore,  $m_x$  is injective. Since  $R/P$  is finite,  $m_x$  is surjective. Therefore,  $xy = m_x(y) = 1$  for some  $y \in R/P$ . Then  $x$  is a unit in  $R/P$ . It follows that  $R/P$  is a field. Therefore,  $P$  is a maximal ideal in  $R$ .

### Optional Part

1. Let  $R$  be a commutative ring, and let  $P$  be a prime ideal of  $R$ . Suppose that 0 is the only zero-divisor of  $R$  contained in  $P$ . Show that  $R$  is an integral domain.

**Answer.** Let  $x, y \in R$ . Suppose  $xy = 0$ . Then  $xy \in P$ , and so  $x \in P$  or  $y \in P$ . Without loss of generality, we assume that  $x \in P$ . If  $y \neq 0$ , then  $x$  is a zero-divisor of  $R$  contained in  $P$ . Then  $x = 0$ . It follows that  $y = 0$  or  $x = 0$ . Therefore,  $R$  is an integral domain.

2. An element  $a$  of a ring  $R$  is **nilpotent** if  $a^n = 0$  for some  $n \in \mathbb{Z}^+$ . Show that the collection  $N$  of all nilpotent elements in a commutative ring  $R$  is an ideal, called the **nilradical** of  $R$ .

**Answer.** Note that it is just the set  $\sqrt{\{0\}}$ . By Ex.1 in the compulsory part, it is an ideal.

3. Show that the nilradical  $N$  of a commutative ring  $R$  is contained in every prime ideal of  $R$ .

**Answer.** By definition of prime ideals,  $x^k = 0 \in P \Rightarrow x \in P$  for any prime ideal  $P$ . Thus the nilradical is contained in each prime ideal.

4. What is the relationship between the radical  $\sqrt{I}$  of an ideal  $I$  in a commutative ring  $R$  and the nilradical of the quotient ring  $R/I$ ? Explain your answer carefully.

**Answer.** Let  $\pi : R \rightarrow R/I$  be the natural projection. Then  $\pi(\sqrt{I}) = \sqrt{0_{R/I}}$ . **Proof:** For any  $x \in \sqrt{I}$ ,  $x^n \in I$  for some  $n > 0$ . Then  $\pi(x)^n \in 0_{R/I}$ , and so  $\pi(x) \in \sqrt{0_{R/I}}$ . Therefore,  $\pi(\sqrt{I}) \subseteq \sqrt{0_{R/I}}$ .

Conversely, let  $y \in \sqrt{0_{R/I}}$ , then  $y^n = 0_{R/I}$  for some  $n > 0$ . Since  $\pi$  is surjective,  $\pi(x) = y$  for some  $x \in R$ . Then  $\pi(x^n) = y^n = 0_{R/I}$ . Then  $x^n \in I$ , and so  $x \in \sqrt{I}$ . Then  $y = \pi(x) \in \pi(\sqrt{I})$ . It follows that  $\pi(\sqrt{I}) = \sqrt{0_{R/I}}$ .

Since  $\sqrt{I} \supseteq I = \ker(\pi)$ ,  $\sqrt{I}$  is the ideal corresponding to  $\sqrt{0_{R/I}}$  via  $\pi$ .

5. Let  $F$  be a subfield of a field  $E$ .

(a) For  $\alpha_1, \dots, \alpha_n \in E$ , define the *evaluation map*

$$\phi_{\alpha_1, \dots, \alpha_n} : F[x_1, \dots, x_n] \rightarrow E$$

by sending  $f(x_1, \dots, x_n)$  to  $f(\alpha_1, \dots, \alpha_n)$ . Show that  $\phi_{\alpha_1, \dots, \alpha_n}$  is a ring homomorphism. We say that  $(\alpha_1, \dots, \alpha_n) \in F^n$  is a zero of  $f = f(x_1, \dots, x_n)$  if  $f(\alpha_1, \dots, \alpha_n) = 0$ , or equivalently, if  $\phi_{\alpha_1, \dots, \alpha_n}(f) = 0$ .

(b) Given a subset  $V \subset F^n$ , show that the set of polynomials  $f \in F[x_1, \dots, x_n]$  such that every element in  $V$  is a zero of  $f$  forms an ideal of  $F[x_1, \dots, x_n]$ .

**Answer.** (a) It is the multi-variable evaluation homomorphism, and can be realized as  $(ev_{\alpha_1} \circ ev_{\alpha_2} \circ \dots \circ ev_{\alpha_n})|_{F[x_1, \dots, x_n]}$ , where each  $ev_{\alpha_r} : E[x_1, \dots, x_r] = E[x_1, \dots, x_{r-1}][x_r] \rightarrow E[x_1, \dots, x_{r-1}]$  is the evaluation homomorphism sending  $x_r$  to  $\alpha_r$ .

(b) That set is  $\bigcap_{(\alpha_1, \dots, \alpha_n) \in V} \ker(\phi_{\alpha_1, \dots, \alpha_n})$ , and is an ideal of  $F[x_1, \dots, x_n]$

6. Prove the *equivalence* of the following two statements:

**Fundamental Theorem of Algebra:** Every nonconstant polynomial in  $\mathbb{C}[x]$  has a zero in  $\mathbb{C}$ .

**Nullstellensatz for  $\mathbb{C}[x]$ :** Let  $f_1(x), \dots, f_r(x) \in \mathbb{C}[x]$  and suppose that every  $\alpha \in \mathbb{C}$  that is a zero of all  $r$  of these polynomials is also a zero of a polynomial  $g(x)$  in  $\mathbb{C}[x]$ . Then some power of  $g(x)$  is in the smallest ideal of  $\mathbb{C}[x]$  that contains the  $r$  polynomials  $f_1(x), \dots, f_r(x)$ .

**Answer.** ( $FTA \implies N$ ) Suppose we have FTA. Then every nonconstant polynomial in  $\mathbb{C}[x]$  factors into  $c(x - z_1)^{k_1} \dots (x - z_r)^{k_r}$  for some  $c, z_1, \dots, z_r \in \mathbb{C}, k_1, \dots, k_r \in \mathbb{Z}_{>0}$ .

Let  $f_1(x), \dots, f_r(x) \in \mathbb{C}[x]$  and suppose that every  $\alpha \in \mathbb{C}$  that is a zero of all  $r$  of these polynomials is also a zero of a polynomial  $g(x)$  in  $\mathbb{C}[x]$ . Let  $I = (f_1, \dots, f_r)$ . Since  $\mathbb{C}[x]$  is a PID,  $I = (h)$  for some  $h \in \mathbb{C}[x]$ .

Case 1.  $h$  is not a constant polynomial. Write  $h = c(x - z_1)^{k_1} \dots (x - z_s)^{k_s}$  as above. Note that each  $f_j \in (h)$ , so  $h | f_j$  for each  $j$ . Then each  $z_i$  is a root of  $h$ , hence a root of  $f_j$  for each  $j$ . By assumption  $g(z_i) = 0$  for each  $i = 1, 2, \dots, s$ . Then  $(x - z_1) \dots (x - z_s) \mid g$ . Then  $g^{\max(k_1, \dots, k_s)} \in (h) = I$ .

Case 2.  $h$  is a constant. If  $h \neq 0$ , then  $(h) = (1)$ , and  $g \in (h)$ . If  $h = 0$ , then each  $f_i = 0$ , so  $g$  is zero everywhere on  $\mathbb{C}$ , and so  $g = 0 \in (h)$ .

( $FTA \Leftarrow N$ ) Conversely, suppose we have  $N$ . Let  $f$  be a nonconstant polynomial in  $\mathbb{C}[x]$ . Suppose  $f$  has no root in  $\mathbb{C}$ . Let  $g = 1$ , then any root of  $f$  is a root of  $g$ , and by  $N$ ,  $g^n \in (f)$ . But then  $1 \in (f)$ . This is absurd because  $\deg(f) > 0$ . Therefore,  $f$  has a root in  $\mathbb{C}$ .