

Exercise 4

1. We continue our study of the Lebesgue measure beginning in Ex 3. Show that

(a) \mathcal{L}^n is a Borel measure.

(b) For every set E , there exists a sequence of open sets $\{G_k\}$ satisfying $E \subset G_k$ and

$$\mathcal{L}^n(E) = \lim_{k \rightarrow \infty} \mathcal{L}^n(G_k) .$$

(c) For every measurable set A , there exists a sequence of compact sets $\{K_j\}$ satisfying $K_j \subset A$ and

$$\mathcal{L}^n(A) = \lim_{j \rightarrow \infty} \mathcal{L}^n(K_j) .$$

Hint: First assume A is bounded.

2. Let $(\mathbb{R}^n, \mathcal{B}, \mu)$ be a measure space where \mathcal{B} is the Borel σ -algebra on \mathbb{R}^n . Suppose that μ is translational invariant, i.e., $\mu(E+x) = \mu(E)$, $\forall x \in \mathbb{R}^n$, $E \in \mathcal{B}$, and that μ is non-trivial in the sense that $0 < \mu([0, 1]^n) < \infty$. Show that μ is a constant multiple of the Lebesgue measure on \mathbb{R}^n when restricted to \mathcal{B} .

3. Let X be a metric space and \mathcal{C} be a subset of \mathcal{P}_X containing the empty set and X . Assume that there is a function $\rho : \mathcal{C} \rightarrow [0, \infty]$ satisfying $\rho(\emptyset) = 0$. For each $\delta > 0$, show that (a)

$$\mu_\delta(E) = \inf \left\{ \sum_k \rho(C_k) : E \subset \bigcup_k C_k, \text{ diameter}(C_k) \leq \delta \right\}$$

is an outer measure on X , and (b) $\mu(E) = \lim_{\delta \rightarrow 0} \mu_\delta(E)$ exists and is also an outer measure on X .

4. Consider in the previous problem the Euclidean space \mathbb{R}^n , $\mathcal{C} = \mathcal{P}_X$ and $s \in [0, \infty)$. Let

$$\rho(C) = (\text{diam}(C))^s ,$$

where the diameter of C is given by $\sup_{x,y \in C} |x-y|$. Show that the resulting outer measures are Borel measures.

5. Let X be a metric space and $C(X)$ the collection of all continuous real-valued functions in X . Let \mathcal{A} consist of all sets of the form $f^{-1}(G)$ which $f \in C(X)$ and G is open in \mathbb{R} .

The “Baire σ -algebra” is the σ -algebra generated by \mathcal{A} . Show that the Baire σ -algebra coincides with the Borel σ -algebra \mathcal{B} .

6. Identify the Riesz measures corresponding to the following positive functionals ($X = \mathbb{R}$):

(a) $\Lambda_1 f = \int_a^b f dx$, and

(b) $\Lambda_2 f = f(0)$.

7. Let c be the counting measure on \mathbb{R} ,

$$c(A) = \begin{cases} \#A, & A \neq \phi, \\ 0, & A = \phi. \end{cases}$$

Is there a positive functional

$$\Lambda f = \int f dc \quad ?$$