

Homework 5 for MATH5070

Topology of Manifolds

Due Wednesday, Nov. 13

1. Denote by $(x_1, \dots, x_n, y_1, \dots, y_n)$ the coordinate functions on \mathbb{R}^{2n} . Let

$$\omega = \sum_{i=1}^n dx_i \wedge dy_i.$$

- (i) What is $d\omega$?
 - (ii) What is $\omega^n = \omega \wedge \dots \wedge \omega$ (wedge n times)?
 - (iii) Let $X = \frac{\partial}{\partial x_1} + \dots + \frac{\partial}{\partial x_k}$ where $k \leq n$. What is $\iota_X \omega$?
 - (iv) Let $\iota : \mathbb{R}^{2n-2} \hookrightarrow \mathbb{R}^{2n}$ be the embedded submanifold of \mathbb{R}^{2n} defined by $x_n = y_n = 0$. What is $\iota^* \omega$?
 - (v) Let $\iota : \mathbb{R}^n \hookrightarrow \mathbb{R}^{2n}$ be the embedded submanifold of \mathbb{R}^{2n} defined by $y_1 = \dots = y_n = 0$. What is $\iota^* \omega$?
 - (vi) Let $\iota : \mathbb{T}^n \hookrightarrow \mathbb{R}^{2n}$ be the embedded submanifold of \mathbb{R}^{2n} defined by $x_i^2 + y_i^2 = 1$ for $1 \leq i \leq n$. What is $\iota^* \omega$?
2. Recall that S^n is a smooth submanifold of \mathbb{R}^{n+1} . For each $p \in S^n$, one can think of the tangent space $T_p S^n$ as the plane in \mathbb{R}^{n+1} that contains p and tangents to S^n . For each $a > 0$, denote $S^n(a)$ the sphere in \mathbb{R}^{n+1} of radius a , centered at the origin.

- (i) Assume X is a smooth vector field on S^n so that $\|X_p\| = 1$ for all $p \in S^n$. Consider the map

$$f_t : S^n \rightarrow \mathbb{R}^{n+1}, \quad p \mapsto p + tX_p.$$

Prove that $\text{Image}(f_t) \subset S^n(\sqrt{1+t^2})$. In what follows we regard f_t as a map from S^n to $S^n(\sqrt{1+t^2})$.

- (ii) Show that f_t is an orientation-preserving diffeomorphism for sufficiently small t .
- (iii) Let ω be an n -form on \mathbb{R}^{n+1} defined as

$$\omega = \sum_{i=1}^{n+1} (-1)^{i-1} x_i dx_1 \wedge \dots \wedge \widehat{dx_i} \wedge \dots \wedge dx_{n+1}.$$

Use (ii) to show that the function $I(t) = \int_{S^n(\sqrt{1+t^2})} \omega$ is a polynomial of t .

- (iv) Apply Stokes' theorem to show that $I(t)$ is a polynomial of t if and only if n is odd.
- (v) Conclude that S^n admits a nowhere-vanishing vector field if and only if n is odd; and there is no Lie group structure on S^{2k} .

3. In 3-dimensional vector calculus, the *divergence theorem* claims that for a region V with boundary S ,

$$\iiint_V (\nabla \cdot \vec{F}) dV = \iint_S \vec{F} \cdot \hat{n} dS;$$

and the *Stokes' theorem* claims that for a surface S with boundary C ,

$$\iint_S (\nabla \times \vec{F}) \cdot \hat{n} dS = \oint_C \vec{F} \cdot d\vec{r}.$$

Derive the two theorems as special cases of the Stokes' theorem for differential forms.

4. Suppose $f \in C^\infty(M)$, $\omega \in \Omega^k(M)$, and X, X_i 's are smooth vector fields on M . Prove:

- (i) $\mathcal{L}_f X \omega = f \mathcal{L}_X \omega + df \wedge \iota_X \omega$
- (ii) $\iota_{[X_1, X_2]} \omega = \mathcal{L}_{X_1} \iota_{X_2} \omega - \iota_{X_2} \mathcal{L}_{X_1} \omega.$
- (iii) $\mathcal{L}_{[X_1, X_2]} \omega = \mathcal{L}_{X_1} \mathcal{L}_{X_2} \omega - \mathcal{L}_{X_2} \mathcal{L}_{X_1} \omega.$
- (iv) $(\mathcal{L}_X \omega)(X_1, \dots, X_k) = \mathcal{L}_X(\omega(X_1, \dots, X_k)) - \sum_{i=1}^k \omega(X_1, \dots, \mathcal{L}_X X_i, \dots, X_k).$

—END—