

2.3 Curve curvature

Definition 2.3.1 (Unit tangent and normal vector). Let $\mathbf{r}(t)$ be a regular parametrized curve.

1. The **unit tangent vector** to the curve at $\mathbf{r}(t)$ is defined by

$$\mathbf{T}(t) = \frac{\mathbf{r}'(t)}{\|\mathbf{r}'(t)\|}.$$

In particular if $\mathbf{r}(s)$ is an arc length parametrization, then

$$\mathbf{T}(s) = \mathbf{r}'(s).$$

2. Suppose $\mathbf{T}'(t) \neq 0$. We define the **unit normal vector** to the curve at $\mathbf{r}(t)$ by

$$\mathbf{N}(t) = \frac{\mathbf{T}'(t)}{\|\mathbf{T}'(t)\|}.$$

In particular if $\mathbf{r}(s)$ is an arc length parametrization, then

$$\mathbf{N}(s) = \frac{\mathbf{T}'(s)}{\|\mathbf{T}'(s)\|} = \frac{\mathbf{r}''(s)}{\|\mathbf{r}''(s)\|}.$$

Prop $\mathbf{T}(t) \perp \mathbf{N}(t)$

$$\begin{aligned} \langle \mathbf{T}(t), \mathbf{N}(t) \rangle &= \left\langle \mathbf{T}(t), \frac{\mathbf{T}'(t)}{\|\mathbf{T}'(t)\|} \right\rangle \\ &= \frac{1}{\|\mathbf{T}'(t)\|} \langle \mathbf{T}(t), \mathbf{T}'(t) \rangle \\ &\stackrel{(*)}{=} 0 \end{aligned}$$

$$(*) \quad \|\mathbf{T}\| \equiv 1$$

$$\begin{aligned} \frac{d}{dt} \langle \mathbf{T}, \mathbf{T} \rangle &= \frac{d}{dt} \|\mathbf{T}\|^2 \equiv 0 \\ \langle \mathbf{T}', \mathbf{T} \rangle + \langle \mathbf{T}, \mathbf{T}' \rangle &= 0 \Rightarrow \langle \mathbf{T}, \mathbf{T}' \rangle = 0 \end{aligned}$$




Figure 7: Unit tangent and unit normal vector

Proposition 2.3.2. Let $\mathbf{r}(t)$ be a regular parametrized curve and $\mathbf{N}(t)$ be the unit normal vector. We have

$$1. \frac{d}{dt} \|\mathbf{r}'\| = \frac{\langle \mathbf{r}', \mathbf{r}'' \rangle}{\|\mathbf{r}'\|}$$

$$2. \mathbf{T}' = \frac{\mathbf{r}''}{\|\mathbf{r}'\|} - \frac{\langle \mathbf{r}', \mathbf{r}'' \rangle}{\|\mathbf{r}'\|^3} \mathbf{r}'$$

Pf ① $\frac{d}{dt} \|\mathbf{r}'\| = \frac{d}{dt} \sqrt{\langle \mathbf{r}', \mathbf{r}' \rangle}$

$$= \frac{\langle \mathbf{r}'', \mathbf{r}' \rangle + \langle \mathbf{r}', \mathbf{r}'' \rangle}{2\sqrt{\langle \mathbf{r}', \mathbf{r}' \rangle}}$$

$$= \frac{\langle \mathbf{r}', \mathbf{r}'' \rangle}{\sqrt{\langle \mathbf{r}', \mathbf{r}' \rangle}}$$

② $\mathbf{T} = \frac{\mathbf{r}'}{\|\mathbf{r}'\|}$

$$\mathbf{T}' = \frac{\|\mathbf{r}'\| \mathbf{r}'' - \|\mathbf{r}'\|' \mathbf{r}'}{\|\mathbf{r}'\|^2}$$

$$= \frac{\|\mathbf{r}'\| \mathbf{r}'' - \frac{\langle \mathbf{r}', \mathbf{r}'' \rangle}{\|\mathbf{r}'\|} \mathbf{r}'}{\|\mathbf{r}'\|^2} = \text{RHS.}$$

Definition 2.3.3 (Curve curvature). Let $\mathbf{r}(t)$ be a regular parametrized curve and $\mathbf{T}(t)$ be the unit tangent to the curve at $\mathbf{r}(t)$. Then the **curvature** of the curve at $\mathbf{r}(t)$ is

$$\kappa(t) = \frac{\|\mathbf{T}'(t)\|}{\|\mathbf{r}'(t)\|}.$$

$\mathbf{T}'(t) = \frac{d\mathbf{T}}{dt}$ = rate of change of direction

$\mathbf{r}'(t) = \frac{d\mathbf{r}}{dt}$ = rate of change of displacement

In particular if $\mathbf{r}(s)$ is an arc length parametrized curve, the curvature is

$$\kappa(s) = \|\mathbf{T}'(s)\|$$

$\mathbf{T}'(s) = \frac{d\mathbf{T}}{ds}$ = change of direction relative to arclength

Proposition 2.3.4. Let $\mathbf{r}(t)$ be a regular parametrized curve. Then the curvature satisfies $\kappa(t) = 0$ for any $a \leq t \leq b$ if and only if $\mathbf{r}(t)$ is a straight line segment joining \mathbf{r}_0 and \mathbf{r}_1 , where $\vec{r}_0 = \vec{r}(a)$, $\vec{r}_1 = \vec{r}(b)$.

$\kappa(t) \equiv 0 \iff$ straight line

Pf (\Rightarrow) Suppose $\kappa(t) \equiv 0$

$$\Rightarrow \frac{\|\tau'(t)\|}{\|\mathbf{r}'(t)\|} \equiv 0 \Rightarrow \tau'(t) \equiv 0$$

$\Rightarrow \tau(t) \equiv \vec{u}$ for some constant unit vector \vec{u}

$$\Rightarrow \frac{\mathbf{r}'(t)}{\|\mathbf{r}'(t)\|} \equiv \vec{u}$$

$$\Rightarrow \mathbf{r}'(t) = \|\mathbf{r}'(t)\| \vec{u}$$

$$\Rightarrow \mathbf{r}(t) = \left(\int_a^t \|\mathbf{r}'(x)\| dx \right) \vec{u} + \vec{v}$$



(\Leftarrow) Suppose $\mathbf{r}(t)$ is straight line

Then $\mathbf{r}(t) = \vec{v} + \alpha(t) \vec{u}$, where

$\alpha(t)$ is increasing function

$$\mathbf{r}'(t) = \alpha'(t) \vec{u} \neq \vec{0} \Rightarrow \alpha' > 0$$

$$\tau = \frac{\mathbf{r}'}{\|\mathbf{r}'\|} = \frac{\alpha'(t) \vec{u}}{\|\alpha'(t) \vec{u}\|}$$

$$= \frac{\alpha'(t)}{|\alpha'(t)| \|\vec{u}\|} \vec{u} = \vec{u}$$

$$\kappa(t) = \frac{\|\tau'(t)\|}{\|\mathbf{r}'(t)\|} = \frac{\|\vec{0}\|}{\|\mathbf{r}'(t)\|} = 0$$

Proposition 2.3.5 (Formulas for curvature). Let $\mathbf{r}(t)$ be a regular parametrized curve.

1. Suppose $\mathbf{r}(t) = (x(t), y(t))$ is a plane curve. Then

$$\kappa(t) = \frac{|x'y'' - x''y'|}{(x'^2 + y'^2)^{\frac{3}{2}}}.$$

$$\frac{\left| \det \begin{bmatrix} x' & y' \\ x'' & y'' \end{bmatrix} \right|}{\|\mathbf{r}'\|^3}$$


2. Suppose $\mathbf{r}(t) = (x(t), y(t), z(t))$ is a space curve. Then

$$\kappa(t) = \frac{\|\mathbf{r}' \times \mathbf{r}''\|}{\|\mathbf{r}'\|^3}.$$

Rmk

① can be considered as a special case of ②

Regard $\mathbf{r}(t) = (x, y) = (x, y, 0)$



$$\kappa = \frac{\|\mathbf{T}'\|}{\|\mathbf{r}'\|} = \left\| \frac{\|\mathbf{r}'\|^2 \mathbf{r}'' - \langle \mathbf{r}', \mathbf{r}'' \rangle \mathbf{r}'}{\|\mathbf{r}'\|^4} \right\|$$

$$\frac{d}{dt} \|\mathbf{r}'\| = \frac{\langle \mathbf{r}', \mathbf{r}'' \rangle}{\|\mathbf{r}'\|}$$

$$\mathbf{T}' = \frac{\mathbf{r}''}{\|\mathbf{r}'\|} - \frac{\langle \mathbf{r}', \mathbf{r}'' \rangle}{\|\mathbf{r}'\|^3} \mathbf{r}'$$

1. Suppose $\mathbf{r}(t) = (x(t), y(t))$ is a plane curve. Then

$$\kappa(t) = \frac{|x'y'' - x''y'|}{(x'^2 + y'^2)^{\frac{3}{2}}}.$$

$$\kappa = \frac{\|\mathbf{T}'\|}{\|\mathbf{r}'\|} = \left\| \frac{\|\mathbf{r}'\|^2 \mathbf{r}'' - \langle \mathbf{r}', \mathbf{r}'' \rangle \mathbf{r}'}{\|\mathbf{r}'\|^4} \right\|$$

Pf $\mathbf{r}' = (x', y')$ $\mathbf{r}'' = (x'', y'')$

$$\langle \mathbf{r}', \mathbf{r}'' \rangle = x'x'' + y'y''$$

$$\|\mathbf{r}'\|^2 \mathbf{r}'' - \langle \mathbf{r}', \mathbf{r}'' \rangle \mathbf{r}' = (x'^2 + y'^2)(x'', y'') - (x'x'' + y'y'')(x', y')$$

$$= [\underbrace{(x')^2 x'' + (y')^2 x''}_{\text{red}}, \underbrace{(x')^2 x'' - x'y'y''}_{\text{red}}, \underbrace{(x'')^2 y'' + (y')^2 y''}_{\text{green}}, \underbrace{(x'')^2 y'' - x'x''y'}_{\text{green}}]$$

$$= [-y'(x'y'' - x''y'), x'(x'y'' - x''y'')]$$

$$= (x'y'' - x''y')(-y', x')$$

$$\kappa = \frac{1}{\|\mathbf{r}'\|^4} \|(x'y'' - x''y')(-y', x')\|$$

$$= \frac{1}{\|\mathbf{r}'\|^4} |x'y'' - x''y'| \underbrace{\sqrt{(-y')^2 + (x')^2}}_{\|\mathbf{r}'\|} = \frac{1}{\|\mathbf{r}'\|^3} |x'y'' - x''y'|$$

2. Suppose $\mathbf{r}(t) = (x(t), y(t), z(t))$ is a space curve. Then

$$\kappa(t) = \frac{\|\mathbf{r}' \times \mathbf{r}''\|}{\|\mathbf{r}'\|^3}.$$

$$\kappa = \frac{\|\mathbf{T}'\|}{\|\mathbf{r}'\|} = \left\| \frac{\|\mathbf{r}'\|^2 \mathbf{r}'' - \langle \mathbf{r}', \mathbf{r}'' \rangle \mathbf{r}'}{\|\mathbf{r}'\|^4} \right\|$$

pf Suppose $\mathbf{r}(t) = (x(t), y(t), z(t))$ is a space curve. Then

$$\begin{aligned} & \left\| \|\mathbf{r}'\|^2 \mathbf{r}'' - \langle \mathbf{r}', \mathbf{r}'' \rangle \mathbf{r}' \right\|^2 \\ = & \langle \|\mathbf{r}'\|^2 \mathbf{r}'' - \langle \mathbf{r}', \mathbf{r}'' \rangle \mathbf{r}', \|\mathbf{r}'\|^2 \mathbf{r}'' - \langle \mathbf{r}', \mathbf{r}'' \rangle \mathbf{r}' \rangle \\ = & \|\mathbf{r}'\|^4 \|\mathbf{r}''\|^2 - 2 \langle \mathbf{r}', \mathbf{r}'' \rangle^2 \|\mathbf{r}'\|^2 + \langle \mathbf{r}', \mathbf{r}'' \rangle^2 \|\mathbf{r}'\|^2 \\ = & \|\mathbf{r}'\|^4 \|\mathbf{r}''\|^2 - \langle \mathbf{r}', \mathbf{r}'' \rangle^2 \|\mathbf{r}'\|^2 \\ = & \|\mathbf{r}'\|^2 (\|\mathbf{r}'\|^2 \|\mathbf{r}''\|^2 - \langle \mathbf{r}', \mathbf{r}'' \rangle^2) \\ = & \|\mathbf{r}'\|^2 \|\mathbf{r}' \times \mathbf{r}''\|^2. \end{aligned}$$

$$\begin{aligned} \kappa &= \frac{\|\mathbf{r}'\| \|\mathbf{r}' \times \mathbf{r}''\|}{\|\mathbf{r}'\|^4} \\ &= \frac{\|\mathbf{r}' \times \mathbf{r}''\|}{\|\mathbf{r}'\|^3}. \end{aligned}$$

Theorem 2.3.6. Suppose $\mathbf{r}(s)$ is an arc length parametrized curve. Then

1. $\kappa(s) = \|\mathbf{r}''(s)\|$

2. $\mathbf{T}'(s) = \kappa(s)\mathbf{N}(s)$

Pf ① $\mathbf{T}(s) = \frac{\mathbf{r}'(s)}{\|\mathbf{r}'(s)\|} = \mathbf{r}'(s)$

$$\kappa(s) = \frac{\|\mathbf{T}'(s)\|}{\|\mathbf{r}'(s)\|} = \|\mathbf{T}'(s)\| = \|\mathbf{r}''(s)\|$$

② $\mathbf{N}(s) = \frac{\mathbf{T}'(s)}{\|\mathbf{T}'(s)\|}$

$$\mathbf{T}'(s) = \|\mathbf{T}'(s)\| \mathbf{N}(s)$$

$$= \kappa(s) \mathbf{N}(s)$$

Example 2.3.7 (Circle). Let $\mathbf{r}(\theta) = (r \overset{x}{\parallel} \cos \theta, r \overset{y}{\parallel} \sin \theta)$, $0 < \theta < 2\pi$, be the circle of radius $r > 0$ centered at the origin. Then

$$x(\theta) = x = r \cos \theta$$

$$y(\theta) = y = r \sin \theta$$

$$x' = -r \sin \theta$$

$$y' = r \cos \theta$$

$$x'' = -r \cos \theta$$


$$y'' = -r \sin \theta$$

$$\kappa(t) = \frac{|x'y'' - x''y'|}{(x'^2 + y'^2)^{\frac{3}{2}}}.$$

$$\kappa = \frac{|x'y'' - x''y'|}{((x')^2 + (y')^2)^{\frac{3}{2}}} = \frac{\left| \det \begin{bmatrix} x' & y' \\ x'' & y'' \end{bmatrix} \right|}{\|\mathbf{r}'\|^3}$$

$$= \frac{|r^2 \sin^2 \theta - (-r^2 \cos^2 \theta)|}{|r^2 (\sin^2 \theta + \cos^2 \theta)|^{\frac{3}{2}}}$$

$$= \frac{|r^2|}{|r^2|^{\frac{3}{2}}} = \frac{1}{r}$$




Example 2.3.8 (Cycloid). The **cycloid** is the curve parametrized by

$$\mathbf{r}(\theta) = (\theta - \sin \theta, 1 - \cos \theta), \text{ for } \theta \in (0, 2\pi).$$

$$\kappa(t) = \frac{|x'y'' - x''y'|}{(x'^2 + y'^2)^{\frac{3}{2}}}.$$

$$\begin{cases} \mathbf{r}'(\theta) = (1 - \cos \theta, \sin \theta) \\ \mathbf{r}''(\theta) = (\sin \theta, \cos \theta) \end{cases}$$

$$\begin{aligned} \kappa(\theta) &= \frac{|x'y'' - x''y'|}{(x'^2 + y'^2)^{\frac{3}{2}}} \\ &= \frac{|(1 - \cos \theta) \cos \theta - \sin \theta \sin \theta|}{((1 - \cos \theta)^2 + (-\sin \theta)^2)^{\frac{3}{2}}} \\ &= \frac{1 - \cos \theta}{(2 - 2 \cos \theta)^{\frac{3}{2}}} \\ &= \frac{1}{2^{\frac{3}{2}} \sqrt{1 - \cos \theta}}. \end{aligned}$$



Example 2.3.9 (Helix). Let $a, b > 0$ be constants. The space curve $\mathbf{r}(\theta) = (a \cos \theta, a \sin \theta, b\theta)$, $\theta \in \mathbb{R}$, is called a **helix**. Then

$$\mathbf{r}' = \mathbf{r}'(\theta) = (-a \sin \theta, a \cos \theta, b)$$

$$\mathbf{r}'' = (-a \cos \theta, -a \sin \theta, 0)$$

$$\mathbf{r}' \times \mathbf{r}'' = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ -a \sin \theta & a \cos \theta & b \\ -a \cos \theta & -a \sin \theta & 0 \end{vmatrix}$$

$$\begin{aligned} \|\mathbf{r}' \times \mathbf{r}''\| &= \sqrt{\begin{vmatrix} a \cos \theta & b \\ -a \sin \theta & 0 \end{vmatrix}^2 + \begin{vmatrix} -a \sin \theta & b \\ -a \cos \theta & 0 \end{vmatrix}^2 + \begin{vmatrix} -a \sin \theta & a \cos \theta \\ -a \cos \theta & -a \sin \theta \end{vmatrix}^2} \\ &= \sqrt{(ab \sin \theta)^2 + (ab \cos \theta)^2 + a^4} \\ &= \sqrt{a^4 + a^2 b^2} = a \sqrt{a^2 + b^2} \end{aligned}$$

$$\kappa(\theta) = \frac{\|\mathbf{r}' \times \mathbf{r}''\|}{\|\mathbf{r}'\|^3} = \frac{a \sqrt{a^2 + b^2}}{(a^2 + b^2)^{3/2}} = \frac{a}{a^2 + b^2}$$

$\mathbf{r}(t) = (x(t), y(t), z(t))$ is a space curve.

$$\kappa(t) = \frac{\|\mathbf{r}' \times \mathbf{r}''\|}{\|\mathbf{r}'\|^3}.$$




Figure 5: Helix

Proposition 2.3.10 (Curvature of graphs of functions).

1. (Rectangular coordinates): The curvature of the curve given by the graph of function $y = f(x)$ in rectangular coordinates is

$$\kappa(x) = \frac{|f''|}{(1 + f'^2)^{\frac{3}{2}}}.$$

2. (Polar coordinates): The curvature of the curve given by the graph of function $r = r(\theta)$ in polar coordinates is

$$\kappa(\theta) = \frac{|r^2 + 2r'^2 - rr''|}{(r^2 + r'^2)^{\frac{3}{2}}}.$$


$$\kappa(t) = \frac{|x'y'' - x''y'|}{(x'^2 + y'^2)^{\frac{3}{2}}}.$$

$$\textcircled{2} \quad x = r(\theta) \cos \theta$$


$$y = r(\theta) \sin \theta$$

Exercise / Note

Pf ① $x = t \quad y = f(t)$
 $x' = 1 \quad y' = f'$
 $x'' = 0 \quad y'' = f''$



$$\kappa = \frac{|x'y'' - x''y'|}{[(x')^2 + (y')^2]^{\frac{3}{2}}} = \frac{|f''|}{[1 + (f')^2]^{\frac{3}{2}}}$$



$$r = r(\theta) = r_0$$

$$r' = r'' = 0$$

$$\kappa = \frac{|r_0^2|}{|r_0^2|^{\frac{3}{2}}} = \frac{1}{r_0}$$

Example 2.3.11 (Catenary). The **catenary** is the curve given by the graph of the function $y = \cosh x$. Show that the curvature of the catenary is

$$\kappa = \frac{1}{\cosh^2 x}.$$



$$\kappa(x) = \frac{|f''|}{(1 + f'^2)^{\frac{3}{2}}}.$$

Proof. Observe that

$$\begin{cases} f'(x) = \sinh x, \\ f''(x) = \cosh x. \end{cases}$$

By Proposition [2.3.10](#), the curvature of the catenary is

$$\begin{aligned} \kappa &= \frac{|f''|}{(1 + f'^2)^{\frac{3}{2}}} \\ &= \frac{\cosh x}{(1 + \sinh^2 x)^{\frac{3}{2}}} \\ &= \frac{\cosh x}{(\cosh^2 x)^{\frac{3}{2}}} \\ &= \frac{1}{\cosh^2 x} \end{aligned}$$



Parametrized Curve	Arc length	Curvature
Plane curve $\mathbf{r}(t) = (x(t), y(t)),$ $a < t < b$	$\int_a^b \ \mathbf{r}'\ dt$	$\kappa(t) = \frac{ x'y'' - x''y' }{(x'^2 + y'^2)^{\frac{3}{2}}}$
Space curve $\mathbf{r}(t) = (x(t), y(t), z(t)),$ $a < t < b$	$\int_a^b \ \mathbf{r}'\ dt$	$\kappa(t) = \frac{\ \mathbf{r}' \times \mathbf{r}''\ }{\ \mathbf{r}'\ ^3}$
Arc length parametrized curve $\mathbf{r}(s)$ with $\ \mathbf{r}'(s)\ = 1$ $a < s < b$	$b - a$	$\kappa(s) = \ \mathbf{r}''(s)\ $
Circle $\mathbf{r}(\theta) = (r \cos \theta, r \sin \theta),$ $0 < \theta < 2\pi$	$2\pi r$	$\kappa = \frac{1}{r}$
Cycloid $\mathbf{r}(\theta) = (\theta - \sin \theta, \cos \theta),$ $\theta \in (0, 2\pi)$	8	$\frac{1}{2^{\frac{3}{2}} \sqrt{1 - \cos \theta}}$
Helix $\mathbf{r}(\theta) = (a \cos \theta, a \sin \theta, b\theta),$ $0 < \theta < 2\pi$	$2\pi \sqrt{a^2 + b^2}$	$\frac{a}{a^2 + b^2}$
Graph of function $y = f(x)$ in rectangular coordinates $\mathbf{r}(t) = (t, f(t)),$ $a < t < b$	$\int_a^b \sqrt{1 + f'^2} dx$	$\frac{ f'' }{(1 + f'^2)^{\frac{3}{2}}}$
Graph of function $r = r(\theta)$ in polar coordination $\mathbf{r}(\theta) = (r \cos \theta, r \sin \theta),$ $\alpha < \theta < \beta$	$\int_\alpha^\beta \sqrt{r^2 + r'^2} d\theta$	$\frac{ r^2 + 2r'^2 - rr'' }{(r^2 + r'^2)^{\frac{3}{2}}}$

Defn $\mathbf{T}(t) = \frac{\mathbf{r}'(t)}{\|\mathbf{r}'(t)\|}.$

$\mathbf{N}(t) = \frac{\mathbf{T}'(t)}{\|\mathbf{T}'(t)\|}$



$\frac{d}{dt} \|\mathbf{r}'\| = \frac{\langle \mathbf{r}', \mathbf{r}'' \rangle}{\|\mathbf{r}'\|}$

$\mathbf{T}' = \frac{\mathbf{r}''}{\|\mathbf{r}'\|} - \frac{\langle \mathbf{r}', \mathbf{r}'' \rangle}{\|\mathbf{r}'\|^3} \mathbf{r}'$



Proposition 2.3.12. Let $\mathbf{r}(s)$ be an arc length parametrized plane curve and $\theta(s)$ be the angle between \mathbf{T} and positive x -axis. Then

$$\kappa(s) = \left| \frac{d\theta}{ds} \right|.$$

$$x' = \cos \theta$$


$$y' = \sin \theta$$

$$x'' = -\sin \theta \frac{d\theta}{ds}$$

$$y'' = \cos \theta \frac{d\theta}{ds}$$

$$\kappa = \frac{|x y'' - x'' y|}{[(x')^2 + (y')^2]^{\frac{3}{2}}}$$

$$= \frac{\left| \cos^2 \theta \frac{d\theta}{ds} - (-\sin^2 \theta) \frac{d\theta}{ds} \right|}{(\cos^2 \theta + \sin^2 \theta)^{\frac{3}{2}}} = \left| \frac{d\theta}{ds} \right|$$




$$\kappa(t) = \frac{|x' y'' - x'' y'|}{(x'^2 + y'^2)^{\frac{3}{2}}}.$$

Definition 2.3.13 (Signed curvature). Let $\mathbf{r}(t) = (x(t), y(t))$ be a regular parametrized curve. The **signed curvature**, also denoted by κ , of \mathbf{r} is


$$\kappa(t) = \frac{d\theta}{ds} = \frac{x'y'' - y'x''}{(x'^2 + y'^2)^{\frac{3}{2}}}$$

where θ is the angle between the unit tangent vector \mathbf{T} and the positive x -axis so that $\mathbf{T} = (\cos \theta, \sin \theta)$.

Rmk Signed curvature is defined for curves in \mathbb{R}^2 , not \mathbb{R}^3




$$\frac{d\theta}{ds} > 0 \quad (\text{turn left})$$




$$\frac{d\theta}{ds} < 0 \quad (\text{turn right})$$

Definition 2.3.14 (Simple closed curve). A regular **simple closed curve** in \mathbb{R}^2 is a closed and bounded connected subset $C \subset \mathbb{R}^2$ such that for any point $p \in C$, we may find an open set $U_p \subset \mathbb{R}^2$ containing p such that $U_p \cap C$ is the image of a regular parametrized curve.




Rmk A simple closed curve is a loop with no self-intersection

Orientation of a parametrized simple closed curve




Positively oriented (inside = left hand side)




Negatively oriented (outside = left hand side)

Theorem 2.3.15. Let $\mathbf{r}(t)$, $a \leq t \leq b$, be a positively oriented regular parametrization of a regular simple closed curve C such that $\mathbf{r}(t)$ is injective on (a, b) and $\mathbf{r}(a) = \mathbf{r}(b)$. Let $\theta(t)$ be a continuous function such that $\theta(t)$ is the angle between the unit tangent vector $\mathbf{T}(t)$ and the positive x -axis so that $\mathbf{T} = (\cos \theta, \sin \theta)$. Then $\theta(b) - \theta(a) = 2\pi$.



eg



eg




eg




$$\frac{d\theta}{ds} \equiv \frac{2\pi - 0}{2\pi r} = \frac{1}{r}$$

signed curvature



θ discontinuous at vertices
 $5\pi/2 - \pi/2 = 2\pi$




$$3\pi - \pi = 2\pi$$

$$\theta(b) - \theta(a) = 2\pi$$

Why?

Deform a simple closed curve to a circle





$$d = \theta(b) - \theta(a) : 2k_1\pi \rightarrow 2k_2\pi \rightarrow 2\pi \text{ continuously} \Rightarrow k_1 = k_2 = 1$$

Signed curvature of a simple closed curve can be considered as the continuous version of exterior angles of a polygon. The following theorem is the continuous version of the theorem for sum of exterior angles of polygon.

Theorem 2.3.16. *Let C be a simple closed curve and κ be the signed curvature defined by positively oriented parametrization. Then*

$$\int_C \kappa ds = 2\pi.$$

Meaning of $\int_C \kappa ds$:



Pf $\int_C \kappa ds = \int_C \frac{d\theta}{ds} ds = \int_C d\theta = [\theta]_a^b = \theta(b) - \theta(a) = 2\pi$

Proposition 2.3.17. Let $\mathbf{r}(t)$ be a regular parametrized curve. Then

$$\mathbf{a} = \mathbf{r}'' = \frac{dv}{dt}\mathbf{T} + \kappa v^2\mathbf{N}$$

where $\overset{\text{speed}}{v} = \|\mathbf{v}\| = \|\mathbf{r}'\|$.

Proof. First, we have


$$\mathbf{r}'(t) = v(t)\mathbf{T}(t).$$

Let s be an arc length parameter, that means $s(t)$ is a function such that $\frac{ds}{dt} = \|\mathbf{r}'(t)\|$. Then $\frac{d}{ds}\mathbf{T} = \kappa\mathbf{N}$ by Theorem 2.3.6 and we have

$$\begin{aligned}\mathbf{r}'' &= \frac{dv}{dt}\mathbf{T} + v\frac{d}{dt}\mathbf{T} \\ &= \frac{dv}{dt}\mathbf{T} + v\frac{ds}{dt}\frac{d}{ds}\mathbf{T} \\ &= \frac{dv}{dt}\mathbf{T} + \kappa v^2\mathbf{N}.\end{aligned}$$

Corollary $\kappa(t) = \frac{\langle \mathbf{r}''(t), \mathbf{N} \rangle}{\|\mathbf{r}'(t)\|^2}.$

There is one more way to interpret the curvature of a curve. When we consider $\mathbf{r}(t)$ as the displacement of a moving particle, we try to find a circle which is closest to the trajectory of the particle at a certain point on the curve. Then the curvature of the curve at that point can be interpreted as the reciprocal of the radius of that circle.



Proposition 2.3.18. *Let $\mathbf{r}(t)$ be a regular parametrized curve. Let $s(t)$ be an arc length parameter, that is, $\frac{ds}{dt} = \|\mathbf{r}'(t)\|$ or equivalently $\left\|\frac{d\mathbf{r}}{ds}\right\| = 1$. Let \mathbf{T} and \mathbf{N} be the unit tangent and normal vectors, which can be considered as vector valued functions of t or s , respectively. The curvature κ of the curve is characterized by any of the following conditions.*

1.

$$\kappa(t) = \frac{\|\mathbf{T}'(t)\|}{\|\mathbf{r}'(t)\|}$$

2.

$$\frac{d\mathbf{T}}{ds} = \kappa\mathbf{N}$$

3. If $\mathbf{r} = (x, y)$ is a plane curve, we have

$$\kappa = \frac{|x'y'' - x''y'|}{(x'^2 + y'^2)^{\frac{3}{2}}}.$$

4. If $\mathbf{r} = (x, y, z)$ is a space curve, we have

$$\kappa = \frac{\|\mathbf{r}' \times \mathbf{r}''\|}{\|\mathbf{r}'\|^3}.$$

5.

$$\kappa = \left\|\frac{d^2\mathbf{r}}{ds^2}\right\|$$

6. If $\mathbf{r} = (x, y)$ is a plane curve and θ is the angle between \mathbf{T} and the positive x -axis, that is, $\mathbf{T} = (\cos \theta, \sin \theta)$, then we have

$$\kappa = \frac{d\theta}{ds}.$$

7.

$$\mathbf{r}'' = \frac{dv}{dt}\mathbf{T} + \kappa v^2\mathbf{N}, \text{ where } v = \|\mathbf{r}'(t)\|.$$