



Department of Mathematics
The Institute of Mathematical Sciences
The Chinese University of Hong Kong

數學系
數學科學研究所
香港中文大學

Phone: (852) 3943 7988 • Fax: (852) 2603 5154 • Email: dept@math.cuhk.edu.hk (Math. Dept.)
Room 220, Lady Shaw Building, The Chinese University of Hong Kong, Shatin, N.T., Hong Kong

Joint Geometric Analysis Seminar

(Part of MIST program)

Non-positively curved Ricci surfaces with catenoidal ends

*Dr. Yiming Zang
Institut lie Cartan de Lorraine*

Abstract

A Ricci surface is defined to be a Riemannian surface (M, g_M) whose Gauss curvature K satisfies the differential equation $K\Delta K + g_M(\nabla K, \nabla K) + 4K^3 = 0$. These surfaces were firstly studied by A. Moroianu and S. Moroianu. They have proved that the Gauss curvature does not change sign on a connected Ricci surface. Moreover, they also showed that every non-positively curved Ricci surface admits locally isometric minimal immersions into R^3 . From a famous theorem of Huber, we know that a complete non-positively curved surface with finite total curvature must be biholomorphic to a compact Riemann surface with a finite number of punctured points. This fact inspires us to define the catenoidal ends for complete non-positively curved Ricci surfaces. This talk is divided into two parts:

1. By using an analogue of the Weierstrass representation, we obtain some classification results for Ricci surfaces of genus zero with catenoidal ends.
2. With the help of a recent work of G. Mondello and D. Panov, we get an existence result for Ricci surfaces with positive genus and arbitrary number of catenoidal ends.

Date: February 3, 2023 (Friday)

Time: 3:00pm-4:00pm (Hong Kong time)

ZOOM link: <https://cuhk.zoom.us/j/91805734715>

All are Welcome