

MATH 1510 Chapter 8

8.1 Power Series

Roughly speaking, a power series is a polynomial with degree ∞ .

Definition 8.1 (Power series). A **power series** is a function of the form

$$\begin{aligned} f(x) &= \sum_{k=0}^{\infty} c_k (x - a)^k \\ &= c_0 + c_1(x - a) + c_2(x - a)^2 + c_3(x - a)^3 + \cdots \end{aligned}$$

where a, c_0, c_1, c_2, \dots , are real numbers. We call a the **center** of the series.

The implied domain of a power series is the set of x such that it converges.

It is known that any power series either converges everywhere, or there is a number R such that $f(x)$ converges for all $x \in (a - R, a + R)$ and diverges for all x such that $|x - a| > R$.

This number R is called the **radius of convergence** of the series. For convenience, if $f(x)$ converges everywhere, we say that its radius of convergence is $R = \infty$.

It is also known that

$$R = \lim_{k \rightarrow \infty} \left| \frac{c_k}{c_{k+1}} \right|$$

if the limit exists or is equal to infinity.

(If the limit does not exist, it does not mean R does not exist. It just means that one would have to use some other method to find R .)

Example 8.2. Consider the power series

$$f(x) = \sum_{k=0}^{\infty} x^k = 1 + x + x^2 + x^3 + \cdots$$

Clearly, $a = 0$, $c_k = 1$ and so,

$$R = \lim_{k \rightarrow \infty} \left| \frac{c_k}{c_{k+1}} \right| = 1 \implies R = 1.$$

Since

$$\begin{aligned} f\left(\frac{1}{2}\right) &= 1 + \left(\frac{1}{2}\right) + \left(\frac{1}{2}\right)^2 + \left(\frac{1}{2}\right)^3 + \cdots \text{ converges (geometric series)} \\ f(2) &= 1 + 2 + 2^2 + 2^3 + \cdots \text{ diverges,} \end{aligned}$$

we know that $\frac{1}{2} \in D_f$ but $2 \notin D_f$.

Proposition 8.3 (Interval of convergence). *The implied domain of any power series $f(x)$ with center a and radius of convergence R is an interval of the form:*

$$(a - R, a + R), (a - R, a + R], [a - R, a + R) \text{ or } [a - R, a + R].$$

Proof of Interval of convergence. Let us handle the case when $R \neq 0, \infty$. The cases when $R = 0$ and $R = \infty$ follow from similar arguments. Suppose $|x - a| <$

R . Then, for the series $f(x) = \sum_{k=0}^{\infty} c_k(x - a)^k$,

$$\lim_{k \rightarrow \infty} \left| \frac{(k+1)\text{-th term}}{k\text{-th term}} \right| = \lim_{k \rightarrow \infty} \left| \frac{c_{k+1}(x - a)^{k+1}}{c_k(x - a)^k} \right| = \frac{|x - a|}{R} < 1$$

Therefore, by ratio test, the series converges. On the other hand, if $|x - a| > R$, then

$$\lim_{k \rightarrow \infty} \left| \frac{(k+1)\text{-th term}}{k\text{-th term}} \right| = \lim_{k \rightarrow \infty} \left| \frac{c_{k+1}(x - a)^{k+1}}{c_k(x - a)^k} \right| = \frac{|x - a|}{R} > 1$$

Again, by ratio test, the series diverges. Hence,

$$(a - R, a + R) \subseteq D_f \subseteq [a - R, a + R]$$

and the result follows. □

Thus, we call the implied domain of a power series its **interval of convergence**.

Example 8.4. As in Example 8.2,

$$f(x) = \sum_{k=0}^{\infty} x^k = 1 + x + x^2 + x^3 + \cdots .$$

Since its center and radius of convergence are 0, 1 respectively, we can conclude that the interval of convergence of $f(x)$ is either

$$(-1, 1), (-1, 1], [-1, 1) \text{ or } [-1, 1].$$

That means $f(x)$ converges whenever $x \in (-1, 1)$. In fact, for any $x \in (-1, 1)$,

$$f(x) = \sum_{k=0}^{\infty} x^k = 1 + x + x^2 + x^3 + \cdots = \frac{1}{1-x}.$$

Example 8.5. For the following power series, find its center and radius of convergence. For what x does the series converge?

•

$$f(x) = \sum_{k=0}^{\infty} (k!)(x-1)^k$$

•

$$f(x) = \sum_{k=0}^{\infty} (-1)^k (\sin 2^{-k}) x^k$$

8.2 Taylor Series

While a calculator can only perform basic arithmetic: $+$, $-$, \times , \div , how does it compute something like $\sin 1$ or e^π ? The answer is **Taylor series**.

Definition 8.6 (Taylor series). We say that a function $f(x)$ is **smooth** (or **infinitely differentiable**) over an interval I if $f^{(n)}(x)$ is differentiable over I for any $n \geq 0$. The Taylor series of a smooth function $f(x)$ at a point $x = a$ is:

$$\begin{aligned} T(x) &= \sum_{k=0}^{\infty} c_k (x-a)^k \quad \text{where } c_k = \frac{f^{(k)}(a)}{k!} \\ &= f(a) + f'(a)(x-a) + \frac{f''(a)}{2!}(x-a)^2 + \frac{f^{(3)}(a)}{3!}(x-a)^3 + \cdots \end{aligned}$$

The **Maclaurin series** of $f(x)$ is its Taylor series with center $a = 0$:

$$T(x) = \sum_{k=0}^{\infty} \frac{f^{(k)}(0)}{k!} x^k = f(0) + f'(0)x + \frac{f''(0)}{2!}x^2 + \frac{f^{(3)}(0)}{3!}x^3 + \dots .$$

The **Taylor polynomial of order n** of $f(x)$ at a point $x = a$ is:

$$T_n(x) = \sum_{k=0}^n \frac{f^{(k)}(a)}{k!} (x-a)^k = f(a) + f'(a)(x-a) + \frac{f''(a)}{2!}(x-a)^2 + \dots + \frac{f^{(n)}(a)}{n!}(x-a)^n$$

(Expanded up to order n) The **Maclaurin polynomial of order n** of $f(x)$ is its Taylor polynomial of order n with center $a = 0$:

$$T_n(x) = \sum_{k=0}^n \frac{f^{(k)}(0)}{k!} x^k = f(0) + f'(0)x + \frac{f''(0)}{2!}x^2 + \dots + \frac{f^{(n)}(0)}{n!}x^n.$$

Remark. Observe that the Taylor polynomial $T_n(x)$ of $f(x)$ at $x = a$ is the unique polynomial which satisfies the condition:

$$T_n^{(k)}(a) = f^{(k)}(a), \quad 0 \leq k \leq n.$$

Example 8.7. Consider the function $f(x) = e^x$. It's clearly smooth over \mathbb{R} . Moreover,

$$f^{(n)}(x) = e^x \implies f^{(n)}(0) = 1 \text{ and } f^{(n)}(1) = e.$$


Therefore,

$$\text{Maclaurin series of } f(x) = \sum_{k=0}^{\infty} \frac{1}{k!} x^k = 1 + x + \frac{1}{2!}x^2 + \dots$$

$$\text{Taylor series of } f(x) \text{ about } x = 1 \text{ is } \sum_{k=0}^{\infty} \frac{e}{k!} (x-1)^k = e + e(x-1) + \frac{e}{2!}(x-1)^2 + \dots .$$

By definition, the Maclaurin polynomials of $f(x)$ of orders 0, 1, 2 are

$$1, 1 + x, 1 + x + \frac{1}{2}x^2 \text{ respectively.}$$



From Example 8.7, we can see that Taylor polynomial of order n can be regarded as a degree n polynomial approximation of f around the center a . In particular,

$$T_1(x) = f(a) + f'(a)(x - a)$$

is the linearization of f at a .

- Taylor polynomials of $f(x) = \sin x$ centered at $a = 0$.
- Taylor polynomials of $f(x) = \sin x$ centered at $a = \pi/2$.

The following are some basic Taylor series:

Proposition 8.8.

$$e^x = \sum_{k=0}^{\infty} \frac{1}{k!} x^k = 1 + x + \frac{1}{2!} x^2 + \frac{1}{3!} x^3 + \dots$$

converges for all $x \in \mathbb{R}$

$$\sin x = \sum_{k=0}^{\infty} (-1)^k \frac{1}{(2k+1)!} x^{2k+1} = x - \frac{1}{3!} x^3 + \frac{1}{5!} x^5 - \frac{1}{7!} x^7 + \dots$$

converges for all $x \in \mathbb{R}$

$$\cos x = \sum_{k=0}^{\infty} (-1)^k \frac{1}{(2k)!} x^{2k} = 1 - \frac{1}{2!} x^2 + \frac{1}{4!} x^4 - \frac{1}{6!} x^6 + \dots$$

converges for all $x \in \mathbb{R}$

$$\frac{1}{1-x} = \sum_{k=0}^{\infty} x^k = 1 + x + x^2 + x^3 + \dots$$

converges for all $x \in (-1, 1)$

$$\ln(1+x) = \sum_{k=1}^{\infty} (-1)^{k+1} \frac{1}{k} x^k = x - \frac{1}{2}x^2 + \frac{1}{3}x^3 - \frac{1}{4}x^4 + \dots$$

converges for all $x \in (-1, 1]$

Remark. In general, a function and its Taylor series are not necessarily equal to each other as functions.

For instance, the domain of the Maclaurin series of $\frac{1}{1-x}$ is $(-1, 1)$, while the domain of $\frac{1}{1-x}$ is $(-\infty, 1) \cup (1, \infty)$.

8.3 Operations on Taylor Series

It is known that if a power series $\sum_{k=0}^{\infty} c_k(x-a)^k$ converges to a given function $f(x)$ on an open interval centered at $x = a$, then that power series *is* the Taylor series of $f(x)$ at $x = a$:

$$c_k = \frac{f^{(k)}(a)}{k!}$$

This implies in particular that the power series centered at $x = a$ converging to the function $f(x)$ on an open interval is *unique*: There cannot be another power series with the same center which also converges to the same function on an open interval.

This fact offers a "shortcut" to find the Taylor series of various functions based on known Taylor series.

Suppose

$$f(x) = \sin x, \quad \text{and} \quad g(x) = \cos x.$$

(For the following Taylor series, the centers are assumed to be $a = 0$.)

Taylor series of $(f(x) + g(x))$ = Taylor series of $f(x)$ + Taylor series of $g(x)$

$$= 1 + x - \frac{x^2}{2!} - \frac{x^3}{3!} + \frac{x^4}{4!} + \dots$$

$$\begin{aligned}\text{Taylor series of } (f(x) - g(x)) &= \text{Taylor series of } f(x) - \text{Taylor series of } g(x) \\ &= -1 + x + \frac{x^2}{2!} - \frac{x^3}{3!} - \frac{x^4}{4!} + \cdots\end{aligned}$$

$$\begin{aligned}\text{Taylor series of } (f(x) \cdot g(x)) &= (\text{Taylor series of } f(x)) \cdot (\text{Taylor series of } g(x)) \\ &= x - \frac{2x^3}{3} + \cdots\end{aligned}$$

$$\begin{aligned}\text{Taylor series of } g(f(x)) &= \text{Taylor series of } g(y) \text{ with } y = \text{Taylor series of } f(x) \\ &= 1 - \frac{x^2}{2!} + \frac{5x^4}{4!} + \cdots\end{aligned}$$

$$\begin{aligned}\text{Taylor series of } f'(x) &= \text{Differentiating Taylor series of } f(x) \text{ term by term} \\ &= 1 - \frac{x^2}{2!} + \frac{x^4}{4!} + \cdots\end{aligned}$$

(Notice that this coincides with the Taylor series of $g(x) = \cos x$.)

$$\begin{aligned}\text{Taylor series of } \int_0^x f(t) dt &= \text{Integrating Taylor series of } f(t) \text{ term by term} \\ &= \frac{x^2}{2!} - \frac{x^4}{4!} + \cdots\end{aligned}$$

(Notice that this coincides with the Taylor series of $1 - \cos x$.)

To find the Taylor series of $\frac{f(x)}{g(x)}$ where $g(a) \neq 0$, we start by letting:

$$\text{Taylor series of } \frac{f(x)}{g(x)} = c_0 + c_1x + c_2x^2 + c_3x^3 + c_4x^4 + \cdots$$

Then,

$$\text{Taylor series of } f(x) = \left(\text{Taylor series of } \frac{f(x)}{g(x)} \right) \cdot (\text{Taylor series of } g(x))$$

$$\begin{aligned}x - \frac{1}{6}x^3 + \cdots \\ = (c_0 + c_1x + c_2x^2 + c_3x^3 + c_4x^4 + \cdots) \left(1 - \frac{1}{2}x^2 + \frac{1}{24}x^4 + \cdots \right)\end{aligned}$$

Hence, by comparing the coefficients, we have

$$\begin{aligned}
 x^0 \text{ term: } \quad 0 &= c_0(1) && \implies c_0 = 0 \\
 x^1 \text{ term: } \quad 1 &= c_0(0) + c_1(1) && \implies c_1 = 1 \\
 x^2 \text{ term: } \quad 0 &= c_0\left(-\frac{1}{2}\right) + c_1(0) + c_2(1) && \implies c_2 = 0 \\
 x^3 \text{ term: } \quad -\frac{1}{6} &= c_0(0) + c_1\left(-\frac{1}{2}\right) + c_2(0) + c_3(1) && \implies c_3 = \frac{1}{3}
 \end{aligned}$$

$$\begin{aligned}
 x^4 \text{ term: } \quad 0 &= c_0\left(\frac{1}{24}\right) + c_1(0) + c_2\left(-\frac{1}{2}\right) + c_3(0) + c_4(1) \\
 &&& \implies c_4 = 0
 \end{aligned}$$

and we can conclude that:

$$\text{Taylor series of } \frac{f(x)}{g(x)} = x + \frac{1}{3}x^3 + \dots$$

Example 8.9. • Find the Maclaurin series of $f(x) = \sin^2 x$.

• Hence, find $f^{(10)}(0)$ and $f^{(11)}(0)$.

Example 8.10. Find the Maclaurin series of $f(x) = \sqrt{1+x^2}$.

Example 8.11. Find the Maclaurin series of $f(x) = \frac{x}{1-x^3}$.

Example 8.12. Find the Maclaurin series of $f(x) = \arctan x$.

Example 8.13. Find the Taylor series of $f(x) = \frac{x}{x+1}$ with center $a = 1$.

Example 8.14. Find the Maclaurin polynomial of order 3 of $f(x) = e^{\cos x}$.

8.4 Lagrange Form of Remainder

Although Taylor series is powerful, no machine can really perform an infinite sum. So in practice, a calculator computes a finite sum with acceptable error instead. That means we need to control the error.

Theorem 8.15 (Taylors Theorem). *Suppose $f(x)$ is $(n+1)$ -times differentiable over the interval $[a, x]$ (or $[x, a]$). Then,*

$$f(x) = T_n(x) + R_n(x)$$

where

$$R_n(x) = \frac{f^{(n+1)}(c)}{(n+1)!}(x-a)^{n+1}$$

for some $c \in (a, x)$ (or (x, a)). $R_n(x)$ is called the **Lagrange form of remainder**

Remark. Be careful: $R_n(x)$ looks similar to the $(x-a)^{n+1}$ term of the Taylor series, but is not the same.

Proof of Taylor's Theorem. Let:

$$F(t) = f(t) + f'(t)(x-t) + \frac{f''(t)}{2!}(x-t)^2 + \cdots + \frac{f^{(n)}(t)}{n!}(x-t)^n$$

and $G(t) = (t-x)^{n+1}$. Then $F(t)$ is differentiable over $[a, x]$ (or $[x, a]$) and $G'(t) \neq 0$ over (a, x) (or (x, a)). Notice that $F(x) = f(x)$, $F(a) = T_n(x)$ and

$$\begin{aligned} F'(t) &= \frac{d}{dt} \left(f(t) + f'(t)(x-t) + \frac{f''(t)}{2!}(x-t)^2 + \cdots + \frac{f^{(n)}(t)}{n!}(x-t)^n \right) \\ &= f'(t) + (f''(t)(x-t) - f'(t)) + \frac{1}{2!}(f'''(t)(x-t)^2 - 2f''(t)(x-t)) \\ &\quad + \cdots + \frac{1}{n!}(f^{(n+1)}(t)(x-t)^n - nf^{(n)}(t)(x-t)^{n-1}) \\ &= \frac{1}{n!}f^{(n+1)}(t)(x-t)^n, \end{aligned}$$

Therefore, by Theorem 5.9 (Cauchy's Mean Value Theorem), there exists $c \in (a, x)$ (or (x, a)) such that

$$\begin{aligned} \frac{F'(c)}{G'(c)} &= \frac{F(x) - F(a)}{G(x) - G(a)} \\ \frac{\frac{1}{n!}f^{(n+1)}(c)(x-c)^n}{(n+1)(c-x)^n} &= \frac{f(x) - T_n(x)}{-(a-x)^{n+1}} \\ \frac{f^{(n+1)}(c)}{(n+1)!}(-1)^n(-1)(a-x)^{n+1} &= f(x) - T_n(x) \end{aligned}$$

Hence, $f(x) = T_n(x) + R_n(x)$ as desired. □

Alternatively,

Proof of Taylor's Theorem. Recall that $T_n^{(k)}(a) = f^{(k)}(a)$ for $k = 0, 1, 2, \dots, n$. Moreover, observe that $T_n^{(k)} = 0$ for $k > n$, since T_n is a polynomial of degree at most n .

Let:

$$F(x) = f(x) - T_n(x), \quad G(x) = (x - a)^{n+1}.$$

Then, $F(a) = G(a) = 0$, and by Theorem 5.9 (Cauchy's Mean Value Theorem), we have:

$$\begin{aligned} \frac{f(x) - T_n(x)}{(x - a)^{n+1}} &= \frac{F(x) - F(a)}{G(x) - G(a)} \\ &= \frac{F'(x_1)}{G'(x_1)} \\ &= \frac{f'(x_1) - T'_n(x_1)}{(n+1)(x_1 - a)^n} \end{aligned}$$

for some x_1 between a and x .

Now let:

$$\begin{aligned} F_1(x) &= F'(x) = f'(x) - T'_n(x), \\ G_1(x) &= G'(x) = (n+1)(x - a)^n. \end{aligned}$$

Repeating the same procedure carried out before, we have:

$$\frac{f'(x_1) - T'_n(x_1)}{(n+1)(x_1 - a)^n} = \frac{F_1(x)}{G_1(x)} = \frac{f^{(2)}(x_2) - T_n^{(2)}(x_2)}{(n+1)n(x_2 - a)^{n-1}}$$

for some x_2 between a and x_1 . Repeating this process $n+1$ times, we have:

$$\begin{aligned} \frac{f(x) - T_n(x)}{(x - a)^{n+1}} &= \frac{f'(x_1) - T'_n(x_1)}{(n+1)(x_1 - a)^n} \\ &= \frac{f^{(2)}(x_2) - T_n^{(2)}(x_2)}{(n+1)n(x_2 - a)^{n-1}} \\ &\vdots \\ &= \frac{f^{(n)}(x_n) - T_n^{(n)}(x_n)}{(n+1)n(n-1) \cdots 2(x_n - a)} \\ &= \frac{f^{(n+1)}(x_{n+1}) - 0}{(n+1)!} \end{aligned}$$

for some x_{n+1} between a and x . Letting $c = x_{n+1}$, the theorem follows. \square

Remark. If we apply the Taylor's Theorem with $n = 0$, we have

$$f(x) = T_0(x) + R_0(x) = f(a) + f'(c)(x - a)$$

$$\implies \frac{f(x) - f(a)}{x - a} = f'(c)$$

Thus, Taylor's Theorem can be regarded as a generalization of Lagrange's MVT.

Example 8.16. For any $x > 0$,

$$e^x = T_3(x) + R_3(x) = 1 + x + \frac{1}{2!}x^2 + \frac{1}{3!}x^3 + \frac{e^c}{4!}x^4$$

for some $c \in (0, x)$. In particular, when $x = 1$,

$$e = 1 + 1 + \frac{1}{2} + \frac{1}{6} + \frac{e^c}{24}$$

with $c \approx 0.214114 \in (0, 1)$.

Example 8.17. To approximate the value of $\sin 1$, we apply Taylor's Theorem on $\sin x$:

$$\sin x = T_4(x) + R_4(x) = x - \frac{1}{3!}x^3 + \frac{\cos c}{5!}x^5$$

where $c \in (0, x)$. By putting $x = 1$, we have:

$$\frac{5}{6} - \frac{1}{120} \leq \sin 1 = \frac{5}{6} + \frac{\cos c}{120} \leq \frac{5}{6} + \frac{1}{120}$$

$$0.825 \leq \sin 1 \leq 0.8416667$$

(In fact, $\sin 1 \approx 0.841471$)

Example 8.18. Let's try to approximate

$$\int_0^1 \cos(x^2) dx$$

with an error < 0.001 . First of all, we apply Taylor's Theorem on $\cos t$:

$$\begin{aligned} \cos t &= T_n(t) + R_n(t), \quad \text{where } n = 2m \text{ is even,} \\ &= 1 - \frac{1}{2!}t^2 + \cdots + (-1)^m \frac{1}{(2m)!}t^{2m} + \frac{(-1)^{m+1} \sin c}{(2m+1)!}t^{2m+1} \end{aligned}$$

for some $c \in (0, t)$. By putting $t = x^2$, we have

$$\begin{aligned}
 \text{Exact value} &= \int_0^1 \cos(x^2) dx \\
 &= \int_0^1 \left(1 - \frac{1}{2!}x^4 + \cdots + (-1)^m \frac{1}{(2m)!}x^{4m} \right) dx \\
 &\quad + \int_0^1 (-1)^{m+1} \frac{\sin c}{(2m+1)!} x^{4m+2} dx \\
 &= \text{Approximation} + \text{Error}
 \end{aligned}$$

So, we can see that:

$$\begin{aligned}
 |\text{Error}| &= \left| \int_0^1 (-1)^{m+1} \frac{\sin c}{(2m+1)!} x^{4m+2} dx \right| \leq \int_0^1 \frac{1}{(2m+1)!} x^{4m+2} dx \\
 &= \frac{1}{(2m+1)!(4m+3)},
 \end{aligned}$$

which would be < 0.001 when $m = 2$. Hence, with $m = 2$,

$$\text{Approximation} = \int_0^1 \left(1 - \frac{1}{2!}x^4 + \frac{1}{4!}x^8 \right) dx \approx 0.9046296.$$

(In fact, $\int_0^1 \cos(x^2) dx \approx 0.9045242$.)

Example 8.19. Find the exact value of

$$\frac{1}{1!} - \frac{1}{2!} + \frac{1}{3!} - \frac{1}{4!} + \cdots$$